

COVID-19 **MD**



# COVID-19

Clinical Guidelines Dentistry

| **Update**

Scientific paper drawn up by the COVID-19 MD group aiming the dissemination of the knowledge about the clinical practice in Dentistry in the context of COVID-19.

The expression *Clinical Guidelines* refers to a scientific methodology and type of publication arising from it. This document should not, thus, be understood as legislation nor as any type of imposition of regulatory or legal nature. It is a scientific contribution to the broadening of the knowledge about the professional practice in the context of COVID-19, hopefully serving its recipients.

<b>Authors</b>	<b>10</b>
<b>Advisory Board</b>	<b>14</b>
<b>List of Abbreviations</b>	<b>18</b>
<b>Preliminary remark</b>	<b>19</b>
Objectives	21
Scientific Relevance	21
Topic/Disease	22
Category	22
Target Population	22
Recipients of the Guidelines	22
<b>Introduction</b>	<b>23</b>
Origin of the Infection	23
Epidemiology	23
Virus Structure and Physiopathology	24
Clinical Manifestations and Forms of Presentation of the Disease	25
Routes of Contagion	29
Treatment and Vaccine	30
Context of the consultation of Dentistry and the need for these Clinical Guidelines	33
<b>Materials and Methods</b>	<b>35</b>
Questions	35
Bibliographic Research	35
Inclusion/Exclusion Criteria	36
Selection and Assessment of Evidence	37
Ranking of Evidence and Class of Recommendation	37
Research Results	38
<b>The Pre-Intervention Stages</b>	<b>39</b>
The Schedule	39
The management of the appointments	39
1.1. How can you manage the schedule?	39
1.2. Should the patient be alerted to be on time for the appointment?	39
1.3. Should the patient be alerted to avoid bringing personal ornaments to the appointment?	40
1.4. Should the patient be alerted to avoid paying the consultation in cash?	40
1.5. What are you supposed to do if a patient asks for a teleconsultation?	40
<b>Questionnaire to be completed</b>	<b>42</b>
2.1. Which questions must be asked while making an appointment by telephone?	42
2.2. If the patient answers in the affirmative to any of the questions included in the questionnaire referred to in 2.1, but refers absence of fever, how is scheduling supposed to be handled?	42
2.3. While scheduling an appointment for a patient having a professional profile of risk, as is the case of healthcare workers, or hospital environment, day care center or retirement homes employees, should there be any particular care?	43
2.4. The patient refers having been infected with SARS-CoV-2 but he/she have already tested negative since. How is one expected to proceed?	43
2.5. How should one tackle the situation of a patient having a clinical history of COVID-19 and who is in home isolation?	43

2.6. A patient goes spontaneously to the clinic to schedule an appointment without having previously called. What is the receptionist supposed to do?	44
2.7. Should the accompanying person be advised not to come, whenever possible, and/or avoid his/her permanence in any place of the clinic?	44
2.8. If the patient comes accompanied, is it necessary for the accompanying person to go through the triage for COVID-19?	44
<b>Reception, Waiting Room and Sanitary Facilities</b>	<b>46</b>
<b>Organization of space and furniture</b>	<b>46</b>
3.1. How can you make means of disinfection and protection available to those entering the clinic?	46
3.2. Which special measures must be applied in this customer service area?	46
3.3. How should you organize the waiting room?	46
3.4. How many people can remain in the waiting room? What is the advised capacity?	47
3.5. Which furniture can there be in the waiting room?	47
3.6. Should magazines or newspapers be available?	47
3.7. Should the TV and air conditioning remote controls be available?	47
3.8. Should the water dispenser or the coffee machine be available?	47
3.9. Which changes must be made in the sanitary facilities?	48
3.10. Which means of decontamination must be available at the sanitary facilities?	48
<b>Physical Barriers and Signage</b>	<b>50</b>
4.1. Should there be physical barriers between the reception and the user?	50
4.2. Which signposting can indicate the organization of the waiting spaces or the interdiction of crossing without permission?	50
4.3. Which informative billboards on COVID-19 must be posted and where?	50
<b>Useful Information</b>	<b>52</b>
5.1. What is the rationale behind hand washing?	52
5.2. Which products must be used to sanitize the hands?	53
5.3. Which products must be used to sanitize the hands?	53
5.4. Which are the methods that keep hands' skin healthy?	54
5.5. Should the patients be allowed to brush their teeth, prostheses or removable braces before stepping inside the consultation office?	55
<b>Cleaning the Waiting Room and the Sanitary Facilities</b>	<b>57</b>
6.1. How often should you clean?	57
6.2. How should you clean surfaces?	57
6.3. How should you do this cleaning?	57
6.4. Which cleaning/disinfection products must be used?	57
6.5. What are the most critical surfaces?	58
6.6. How often should the most critical areas or surfaces be cleaned?	58
6.7. What are the materials you must use while cleaning the most critical areas?	58
6.8. How often must you clean the sanitary facilities?	58
6.9. How should you clean the sanitary facilities?	58
6.10. Which are the most critical surfaces in the sanitary facilities?	59
6.11. Which products must be used to clean the sanitary facilities?	59
6.12. What are the materials that you must use to clean the sanitary facilities?	60
<b>People flow</b>	<b>62</b>
7.1. How can the receptionist keep a social distancing, minimizing the fear that this measure might arouse on the patient?	62
7.2. What should you do after finishing serving a patient/accompanying person and before serving the following one?	62
7.3. Should there be separate entrances for users and professionals?	62

7.4. What is the minimum mandatory recommendation of personal protective equipment at the dental clinic?	62
<b>Access to the Clinic</b>	<b>65</b>
<b>Dentists and Staff</b>	<b>65</b>
8.1. Am I able to go to work?	65
8.2. I have a relative (who is part of my household) who is sick with COVID-19. Can I go to work?	65
8.3. I am pregnant or more than 65 years old, I suffer from related systemic pathologies of higher risk. Can I go to work?	65
8.4. How is it possible to control if the staff are infected?	65
<b>Patients and Accompanying People</b>	<b>67</b>
9.1. What shall be checked as soon as the patient enters the clinic?	67
9.2. The patient is feverish or reports having had fever and cough or sore throat and muscular pains over the last 24h. What should you do?	67
9.3. The patient is not feverish. What should you do next?	67
9.4. Can the patient come accompanied?	67
9.5. Should the accompanying person wear a mask too?	68
9.6. In case the accompanying person walks into the clinical office, where should he/she remain and how should you proceed?	68
9.7. In case the accompanying person does not enter the clinical office, where should he/she wait?	68
9.8. What should you do with the patient's belongings (and the accompanying person's as well, if he/she happens to walk into the consultation office) before putting on the appropriate PPE?	68
<b>Maintenance Technicians</b>	<b>70</b>
10.1. How should maintenance technicians be welcome?	70
<b>Delivery men</b>	<b>70</b>
11.1. How should delivery men be welcome?	70
<b>Waste Collectors</b>	<b>70</b>
12.1. How should waste collection employees be welcome?	70
<b>Means of Protection</b>	<b>72</b>
<b>Reception/administrative staff</b>	<b>72</b>
13.1. What is the appropriate protective equipment?	72
<b>Cleaning Staff</b>	<b>73</b>
14.1. What is the appropriate protective equipment?	73
<b>The Stages of Peri-Intervention or Peri-Intervention Stages?</b>	<b>75</b>
<b>Definition of Risk of the Procedure</b>	<b>75</b>
15.1. Should risk areas be identified or marked at the clinic?	75
15.2. What are the differentiated recommendations for the different risk areas?	75
15.3. How should the general procedures in a clinic be ranked in terms of risk?	76
15.4. How should the different clinical procedures be ranked in terms of risk?	77
<b>Preparing the office and individual protection</b>	<b>79</b>
<b>The space and the equipment</b>	<b>79</b>
16.1. What is the stability of the virus in the atmosphere?	79
16.2. What is the reach of contaminated aerosols released in normal breathing in a closed environment?	79
16.3. Which distance may reach the aerosols generated during the operative procedures in Dentistry?	80
16.4. How should dental equipment and the surrounding surfaces be protected?	81
16.5. How should the disinfection of the several components and infrastructures of the consultation office be carried out?	82
16.6. What is the most suitable ventilation system in a clinical office?	83
16.7. Is the use of additional systems of purification of biological contaminants in the air of the clinical office justified?	84

16.8. What must the acclimatization conditions at the clinical office be?	85
16.9. Is it advisable to use means of microbiological decontamination in the water supply systems and water reservoirs of the equipment?	85
16.10. What is the effectiveness of suction systems in the reduction of the generation of aerosols at the source of contamination?	86
16.11. Does the use of antimicrobials in the irrigation system of the ultrasonic devices reduce the risk of contamination during aerosolization?	86
<b>Personal Protective Equipment (PPE)</b>	<b>89</b>
<b>PPE donning</b>	<b>89</b>
17.1. Must the PPE to be worn be defined by the risk involved in the procedure?	89
17.2. What is the appropriate PPE according to the levels of protection?	89
17.3. Which type of masks are there? And what is the indication for its clinical use?	90
17.4. What should you do before putting the PPE on?	91
17.5. How are you supposed to put the PPE on?	91
17.6. What care should be taken over the use of PPE?	92
17.7. What are you supposed to do if the first pair of gloves (inner pair in degree 3 protection PPE) touches a potentially contaminated area?	93
<b>Treatment</b>	<b>95</b>
18.1. How can you reduce microbiological burden in intraoperative stage?	95
18.2. Should a perioral decontamination be performed? How should it be performed?	96
18.3. What is the importance of using a rubber dam?	96
18.4. Should priority be given to the use of single-dose materials?	97
<b>Rotary Instruments</b>	<b>98</b>
19.1. What is the impact of operating instruments on aerosol generation?	98
19.2. Which specifications should cutting/preparation operating instruments have? Are non-return systems better?	99
<b>After-Intervention Stages</b>	<b>101</b>
<b>End of the Consultation and Disinfection of the Medical Office</b>	<b>101</b>
20.1. I have finished the operative procedure. What comes next?	101
20.2. Which are the critical surfaces?	101
20.3. How can you minimize contamination and perform the cleaning of critical areas?	102
20.4. Should protection barriers in screen surfaces, keyboards and mice be removed between patients?	102
20.5. What care should I take with the casing of the digital imaging sensor and the intraoral x-ray films?	102
20.6. How should X-ray positioners be disinfected?	102
20.7. How should digital imaging sensors, intraoral cameras and scanners be protected and disinfected?	103
20.8. Should the purge of rotary instruments be performed before being removed to be sterilized?	103
20.9. Is chemical external disinfection of turbines, counter-angles and scalers enough against SARS-CoV2 or should they be sterilized between patients?	103
20.10. What is the sequence for physical barriers and rotary instruments removal?	103
20.11. How should physical barriers to protect the equipment be eliminated?	104
20.12. Regarding surface disinfection (equipment and workbenches), which product is more efficient and for how long should contact time be? be the contact time?	104
20.13. Is UV radiation an effective sterilization method?	104
20.14. Regarding wrapping and packaging of the consumables used, which is the most suitable disinfection product and what is its contact time?	105
20.15. Is it necessary to make the suitable aeration of the clinical office before starting the following consultation?	105
20.16. How often should the clinical office's floor be cleaned and disinfected?	105
20.17. Regarding the clinical office's floor, which product should be used and for how long should it be left in contact to achieve a suitable disinfection?	105

20.18. How should the clinical office cleaning be performed?	106
20.19. Which cleaning/disinfection products should be used?	106
20.20. Which types of presentation should cleaning products have?	106
20.21. What is the most suitable disinfectant to be used in the form of wipes?	106
20.22. How should surfaces be cleaned in the event they are contaminated with blood or other organic products?	107
<b>Materials and devices to be sent to the prosthetics laboratory</b>	<b>109</b>
21.1. What is the protocol to disinfect dental impressions?	109
21.2. What is the protocol to disinfect dental stone models?	109
21.3. What is the protocol for try-in devices (copings, registration waxes, preliminary crowns/bridges and removable prosthesis) disinfection?	109
21.4. What is the protocol to disinfect pre-existing prosthesis?	109
<b>Removal of PPE</b>	<b>111</b>
22.1. What is the sequence to remove PPE in each one of the levels (1, 2 and 3)?	111
22.2. Where should disposable PPE be placed after being removed? In what type of container?	112
22.3. What is the protocol to remove gloves?	113
22.4. When and how should the dentist remove the surgical mask?	113
22.5. When should the assistant remove PPE?	114
22.6. Can FFP2 (N95) and FFP3 respirators be reused?	114
22.7. In which place should PPE be removed? In the clinical office or somewhere else?	114
22.8. Where should the assistant place PPE after being removed?	114
22.9. How should the face shield and the goggles be washed and disinfected?	115
22.10. Can uniforms be worn again the following day without being washed?	115
22.11. What kind of treatment should be given to non-critical fabric materials: uniforms?	115
22.12. After a day at work how should you behave when you get home?	115
<b>Patient care after treatment</b>	<b>117</b>
23.1. Does the use of a surgical mask by the patient inside the clinic reduce the risk of cross contamination with SARS-CoV-2?	117
23.2. Should patients' movement inside the clinical office be limited after the operative procedures?	117
23.3. Should additional information be provided to the patient in digital form instead of paper form?	117
23.4. How should hand antisepsis be performed before leaving the office, regarding the dentist, the assistant, the patient/ accompanying person?	117
<b>Patient's return to the Reception</b>	<b>119</b>
24.1. Should the patient keep social distance at the reception and means of protection after consultation?	119
24.2. Should the patient keep the mask on in the reception area after the consultation?	119
24.3. Which important additional information should be provided to the patient by the receptionist, in the context of COVID-19 infection, in the post-consultation stage?	119
24.4. Which information should be confirmed to the patient by the receptionist before leaving the clinic?	119
<b>Payment methods</b>	<b>122</b>
25.1. Which payment methods and receipt issuing methods should be recommended?	122
<b>Sterilizing</b>	<b>124</b>
<b>Material Cleaning, Disinfection and Sterilization Standards</b>	<b>124</b>
26.1. Does contamination with SARS-CoV-2 imply a change of the procedures used to clean, decontaminate and sterilize materials and instruments?	124
26.2. Which kind of bag is indicated to pack group III waste?	125
26.3. Should the placement of group III waste in the corresponding container be performed in the clinical office or in the sterilizing room?	125
26.4. Does contamination with SARS-CoV-2 imply a change of the procedures used to handle and process waste?	125

26.5. Should the material/instruments handled by the dentist be considered as semi-critical or critical?	125
26.6. Should the material/instruments handled by the dental assistant be considered as semi-critical or critical?	126
26.7. Is it possible to disinfect FFP2 (N95) and FFP3 respirators?	126
<b>Accidental Exposure</b>	<b>128</b>
How to act in the event of Accidental Exposure	128
27.1. Is the oral cavity a tropism location for SARS-CoV-2?	128
27.2. In the event of an accidental exposure to biological contaminants from a COVID-19 patient, which procedure should be adopted?	128
<b>Conclusion</b>	<b>131</b>
Future Perspectives	131
Updating	132
Acknowledge	132
<b>Appendix I</b>	<b>134</b>
Advisory Board	135
<b>Appendix II</b>	<b>137</b>
Section 1   Questions	138
Section 2   Bibliographic Research	146
Section 3   Articles Included	148
Section 4   Articles Excluded	164
Section 5   Level of Evidence and Classes of Recommendation	174
<b>Appendix III</b>	<b>176</b>
Section 1   Hand hygiene technique using an alcohol-base antiseptic solution (ABAS) or water and soap[84]	177
Section 2   Sequence to put on and remove PPE	178
Section 3   Disinfectant Solutions	180
<b>Bibliography</b>	<b>182</b>





# Authors

## Fernando Guerra

Dentist

Full Professor at the Faculty of Medicine of the University of Coimbra

Director of the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0001-8780-9379>

## António Duarte Mata

Dentist

Full Professor at the Faculty of Dentistry of the University of Lisbon

Coordinator of the LIBPhys-UL Research Unit-UID/FIS/04559/2013, Faculty of Dentistry, University of Lisbon

Main Researcher of the Research Group in Oral Biochemistry and Biology, GIBBO FMDUL

<https://orcid.org/0000-0003-1474-2681>

## Alexandra Vinagre

Dentist

Guest Assistant at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0001-6929-3807>

## Ana Luísa Costa

Dentist

Assistant Professor at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0002-4102-1719>

## Andreia Luís Vieira

Dentist

Guest Assistant at the Faculty of Dentistry of the University of Lisbon

Researcher of the Research Group in Oral Biochemistry and Biology, GIBBO FMDUL

<https://orcid.org/0000-0002-3271-8645>

## Carlos Falcão

Dentist

Assistant Professor at the Faculty of Health Sciences of the Fernando Pessoa University

<https://orcid.org/0000-0001-8031-6176>

## Carlota Duarte de Mendonça

Dentist

Researcher of the Research Group in Oral Biochemistry and Biology, GIBBO FMDUL

<https://orcid.org/0000-0001-6133-1995>

## Catarina Chaves

Doctor

Hospital Assistant in Clinical Pathology

Head of the Laboratory of General Bacteriology

Service of Clinical Pathology, Medical Teaching Hospital of Coimbra

<https://orcid.org/0000-0003-0131-5728>

## David Herrera

Odontologist

Professor at the Faculty of Dentistry of the Complutense University of Madrid

Vice-Dean at the Faculty of Dentistry of Madrid, Spain

<https://orcid.org/0000-0002-5554-2777>

## Isabel Poiares Baptista

Dentist

Assistant Professor at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0003-4503-5365>

## Ivan Toro

Dentist

<https://orcid.org/0000-0002-0437-4152>

## João Braga

Dentist

Assistant Professor at the Faculty of Dentistry of the University of Oporto

<https://orcid.org/0000-0002-0895-1009>

## João Carlos Ramos

Dentist

Assistant Professor at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0003-1965-1092>

## João Miguel Santos

Dentist

Assistant Professor at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0002-2865-9689>

## João Miguel Silveira

Dentist

Assistant Professor at the Faculty of Dentistry of the University of Lisbon

Researcher at the LIBPhys-UL Research Unit-UID/FIS/04559/2013, Faculty of Dentistry, University of Lisbon

Researcher of the Research Group in Oral Biochemistry and Biology, GIBBO FMDUL

<https://orcid.org/0000-0001-9366-0662>

## João Paulo Tondela

Dentist

Assistant Professor at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0003-0787-7713>

## Paulo Palma

Dentist

Guest Assistant Professor at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0003-4730-8072>

## Pedro Nicolau

Dentist

Associate Professor at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0001-7900-4482>

## Pedro Sousa Gomes

Dentist

Associate Professor at the Faculty of Dentistry of the University of Oporto

Researcher at the Center of Research Bone Lab of the Faculty of Dentistry of the University of Oporto

<https://orcid.org/0000-0001-5365-2123>

## Ricardo Dias

Dentist

Guest Assistant at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0002-0824-9754>

## Salomão Rocha

Dentist

Guest Assistant at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0002-5702-333X>

## Sérgio Matos

Dentist

Assistant Professor at the Faculty of Medicine of the University of Coimbra

Researcher at the Center of Research and Innovation in Dental Sciences of the Faculty of Medicine of the University of Coimbra

<https://orcid.org/0000-0002-2548-4326>

Version 1.0.1 | May, 10<sup>th</sup> of 2020. All rights reserved.

# Advisory Board

## *Doctor Ana Lúcia de Pereira Neves Messias*

Guest Teaching Assistant at the Faculty of Medicine of the University of Coimbra

## *Professor Professor Aníbal Diógenes*

Associate Professor at the University of Texas, *Health Center of San Antonio*, United States of America

## *Doctor and Professor António Vaz Carneiro*

Full Professor of Medicine at the Faculty of Medicine of the University of Lisbon and Director of Cochrane Portugal

## *Doctor and Professor Carlos Robalo Cordeiro*

Director of the Faculty of Medicine of the University of Coimbra

## *Doctor and Professor Duarte Marques*

Representative of the Portuguese Society of Stomatology and Dental Medicine

## *Doctor and Professor Gil Alcoforado*

Full Professor

## *Doctor and Professor Ion Zabalegui*

Guest Professor at the University *Complutense* of Madrid, Spain

## *Doctor José Manuel da Costa Rodrigues de Oliveira*

Dentist

## *Doctor José Frias Bulhosa*

Master's degree in public health from the Faculty of Medicine of the University of Coimbra; Representative of the Portuguese Association of Public Service Dentists (APOMED-SP)

## *Doctor Luís António Silva Ferreira*

Dentist

## *Doctor and Professor Luís Pires Lopes*

Head of the Faculty of Dental Medicine of the University of Lisbon

*Doctor and Manuel Gameiro*

Full Professor at the Mechanical Engineering Department of the University of Coimbra

*Doctor and Professor Maria Helena Raposo Fernandes*

Full Professor at the Faculty of Dental Medicine of the University of Porto

*Doctor and Professor Mário Bernardo*

Coordinator of the Integrated Master Degree in Dental Medicine at the Faculty of Dental Medicine at the University of Lisbon

*Doctor and Professor Mariano Sanz*

Full Professor at the University *Complutense* of Madrid, Spain

*Doctor and Professor Marlene Barros*

Head of the Faculty of Dental Medicine of the Catholic University of Portugal

*Doctor Paulo Manuel de Oliveira Nunes*

Dentist

*Doctor Nuno Ventura*

Dentist

*Doctor Ricardo Jorge Martins*

Dentist

*Doctor and Professor Sandra Gavinha*

Head of the Faculty of Health Sciences of Fernando Pessoa University





This document consists of a partial update of the full document COVID-19 | Clinical Guidelines for Dental Medicine, published in May 2020<sup>[1]</sup>. Updating a document of this magnitude may involve replacing it in its entirety or maintaining the matrix with the relevant changes (partial update). The previously published recommendations that justified changes are duly presented below, having been the subject of discussion and deliberation by the COVID-19 MD group.

It should be noted that the process of updating a Clinical Guidance Standard is governed by the following purposes<sup>[2,3]</sup>:

- Alteration or addition of bibliographic references;
- Alteration of the clinical recommendation, in order to reflect the context of current practice; this change can alter the cost-benefit of a given recommendation;
- Important changes in results;
- Changes in health service settings; changes in available resources;
- Ensure that the recommendations take into account the latest government policies;
- Reflect people's involvement in care decisions;
- Patient-centered language.

A quality clinical decision rests on three assumptions:

- The patient's interests; if the clinical decision doesn't satisfy him/her, it will have failed miserably its main goal;
- The practitioner's experience as a key element unifying his/her acumen, knowledge and even intuition, the so-called human factor;
- The scientific evidence as a basis of reliable, objective and essential information.

The Clinical Guidelines presented here intend to be that basis of information and knowledge, objectives which are key to support the clinical decision.

Every theatre of clinical practice has its own specificities. In this context, these Clinical Guidelines do not constitute hermetic orders to abide by, but only a robust body of knowledge allowing the practitioner to set up the implementing protocol which best suits the characteristics of each clinic.

It should also be added that the adoption of these guidelines doesn't guarantee zero risk of acquisition of the infection.

## List of Abbreviations

ADA = American Dental Association

CDC = Centers for Disease Control and Prevention

CHX = Chlorhexidine gluconate

CPC = Cetylpyridinium chloride

ACE2 = Angiotensin-Converting Enzyme 2

IL = Interleukin

IPVD = Povidone-iodine

PXH = Hydrogen peroxide

rt-PCR = Reverse transcription polymerase chain reaction

## Preliminary remark

Given the fact that the COVID-19 disease is an extremely recent one and that, consequently, there is a natural gap in the amount and quality of scientific evidence on this subject, the drafting of some classical Clinical Guidelines is not possible. In this context, these Guidelines were drawn up according to the protocol recently made available by the English National Institute for Health and Care Excellence (NICE), namely the Interim process and methods for developing rapid guidelines on COVID-19 ([nice.org.uk](https://www.nice.org.uk)<sup>[4]</sup>).

These Clinical Guidelines are the result of the work developed by a committee of dentists.

There was no public consultation of representatives of every potential stakeholder about this topic. The goal is to produce Clinical Guidelines which might reduce the entropy and promote the consistency of intervention. However, an advisory board composed of different figures was created so as to review the Clinical Guidelines and proceed to its assessment, the composition of this board being listed in Appendix I.

We promote the identification of the key issues applicable to the scenarios upon return to the clinical practice in dentistry.

No systematic reviews of the literature were conducted. Even though searches were made in the most widely used databases, the main information contained in these Clinical Guidelines stems from the use of guiding documents previously published concerning pandemics that have already occurred.

As expected, there is very scarce literature regarding COVID-19, wherefore a great deal of the alternative basis of evidence we resorted to refers to *Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV)*, *Swine Pandemic (AH1N1) Influenza A* and to *Middle East Respiratory Syndrome Coronavirus (MERS-CoV)*

The databases of choice were:

1. Guides of the World Health Organization (WHO) on COVID-19
2. Recommendations of the Portuguese Health Authority (DGS)
3. NICE and SIGN
4. Cochrane collaboration
5. MEDLINE (via PubMed)
6. International and national scientific societies
7. The available evidence was simply ranked
  - The strategy of search is reproduced in Appendix II, section 2
  - Priority has been given to systematic reviews and meta-analysis
  - Followed by the search for randomized clinical trials
  - Followed by observational studies
  - Whenever it proved impossible, we searched for publications arising from the observation of experts in previous experiences
  - In those cases where no kind of evidence could be found, the experts who are the authors of these guidelines drew up the recommendations through discussion and consensus based primarily on biological plausibility.

We have made a summary critical assessment of the literature but we have not formally assessed the risk of the presence of biases resorting to specific scales.

However, we always briefly and simply indicated the kind of information underlying the clinical recommendations that were produced.

## Objectives

Identify and classify the risks of acquisition and spread of SARS-CoV-2 in the context of pre-, peri-, and post-intervention in oral health.

Produce recommendations on how to act in pre-, peri-, and post-intervention in oral health for every element present at the clinical theatre aiming to minimize the risk of acquisition and/or spread of SARS-CoV-2.

## Scientific Relevance

Given that:

Over the course of the actual pandemic of COVID-19 several sources have identified the activity of dentistry as being at the top of the pyramid as far as the risk of acquisition and spread of the virus SARS-CoV-2 is concerned due to the enormous generation of aerosols which characterize it, the Portuguese government having determined the provisional closure of the clinics of dentistry and stomatology.

Currently, there are no national Clinical Guidelines which convey recommendations on how to act on every step of pre-, peri-, and post-intervention in dentistry aiming the prevention of the acquisition and spread of the virus SARS-CoV-2.

There is, therefore, the need to create these Clinical Guidelines as a set of recommendations on how to act, allowing the return to the activity of dentistry in safe conditions of those involved at the clinical theater, whether they are professionals, partners of the sector or the user population in general.

It should be noted that the Clinical Guidelines intend to support the clinical decision. It is not a document which might replace the clinical decision, but rather one which the practitioner can lean on whenever he/she must choose the best course of action considering his/her experience, the patient's specific interests and the very clinical theater in which the practice takes place. It is not, therefore, a document which aims the tyrannization of evidence over the choice, but one which only allows to decide taking into consideration the greatest robustness of evidence possible.

## **Topic/Disease**

COVID-19 - Minimization of the risk of acquisition and spread of the disease in dentistry in the context of the pandemic.

## **Category**

These are Clinical Guidelines on clinical governance advice, best practice and clinical safety.

## **Target Population**

Patients of dentistry and members of the oral health staff.

## **Recipients of the Guidelines**

All the parties concerned, namely: dentists, stomatologists, odontologists, pharmacists, nurses, oral hygienists, dental prosthetic technicians, clinic's support auxiliary personnel, receptionists, administrative staff, diagnosis and therapy technicians, senior technicians of safety and hygiene at work, technicians of clinical analysis, suppliers of dental material and equipment, medical sales representatives, maintenance technicians, waste collection technicians, professional associations, scientific societies, education institutions, students of dentistry, hospitals, insurance companies and conventions, commercial and industrial partners, policy makers, Portuguese Dental Association, Portuguese Health Authority, sectoral regulatory bodies and patients.


# Introduction

## Origin of the Infection

Throughout history mankind has been faced with cyclical microbiological outbreaks (Figure 1). Among the most recent ones we can point out the Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV), which originated in China, in 2002; the Swine Pandemic (A/H1N1) Influenza A, which shook Mexico in 2009; and the Middle East Respiratory Syndrome Coronavirus (MERS-CoV), embryonic in Saudi Arabia in 2012<sup>[5]</sup>.

In December 2019, a series of cases involving a new virus responsible for an acute respiratory syndrome was detected in Wuhan, China. It is a zoonosis-like infection (originating in a bat) whose causative agent is a new strain of coronavirus, SARS-CoV-2. This virus, which was isolated on 7th January 2020, presents 50% of genetic homology with MERS-CoV, 79% with SARS-CoV-1 and is the causative agent of a new disease, COVID-19.

Figure 1 - Epidemiology. Biological Outbreaks. Adapted from Jiuyang Xu, et al. 2020<sup>[6]</sup>.



## Epidemiology

Over a short period of time, COVID-19 has grown to pandemic proportions, being present in every continent to the exception of Antarctica. Around the time when this text is being written there are 2.626.321 cases all over the world and 181.938 confirmed fatalities (data supplied by the WHO on 24th April 2020). In Portugal, 23.392 cases and 880 fatalities (data supplied by the DGS on 25th April 2020) are reported. However, by the time the readers have access to this document, all these figures will have increased.

The WHO, the American Centers for Disease Control and Prevention (CDC) and modern databases keep an updated interactive epidemiologic map.

There are few indications about the actual prevalence of the disease among dentists. The scarce existing sources indicate that the dental class, even though they are at the top of the chain of potential susceptibility and transmissibility due to the high generation of aerosols, presents a very low prevalence of the disease comparing to other professions in the area of health.

It is not possible, at this stage, to state whether the low prevalence of cases of COVID-19 among oral health professionals is due to the early closure of the care units in consequence of the generalized confinement, or if it results, in effect, from the use of strict protocols of asepsis and prevention of cross infection implemented long ago and resulting from the pandemics by AIDS and Hepatitis B and C viruses.

It should, however, be noted that one of the few studies available in these matters was conducted in Wuhan, during the epidemic stage, in a large clinic housing more than 1.000 professionals, the prevalence of COVID-19 having reached 0,9%.

## Virus Structure and Physiopathology

The physiopathology of this new entity is not fully known. Up till now several hypotheses have been outlined which lack confirmation by different kinds of studies<sup>[7]</sup>.

It is known that SARS-CoV-2 (60-90 nm) belongs to the genus *Betacoronavirus*, subgenus *Sarbecovirus*, *Coronaviridae* family, related to the SARS<sup>[5,7-11]</sup>. It is a positive simple chain RNA (ribonucleic acid) virus with the potential to infect vertebrates<sup>[7,10]</sup>. Besides sharing 79,5% of its genomic sequence with SARS-CoV and the preservation of eight out of the fourteen residues of amino acids of SARS-CoV in the new coronavirus, the mechanism of cellular invasion in the host is also shared by these two viruses: through the receptor and the angiotensin-converting enzyme 2 (ACE2 - a membrane receptor widely diffused by the cells of the organism, including the oral cavity which may be a gateway for the virus). Notwithstanding, the affinity for the receptor is 10 to 20 times higher as far as the new coronavirus is concerned<sup>[5-7,9,12,13]</sup>. Eight out of the fourteen residues of amino acids of SARS-CoV are preserved in the new coronavirus<sup>[7]</sup>.

One of the challenges of this virus has to do with the genomic mutation which occurred during the worldwide dissemination<sup>[12]</sup>.



# Clinical Manifestations and Forms of Presentation of the Disease

The diagnosis of a patient infected with SARS-CoV-2 includes the screening for the signs and clinical symptoms, and the assessment of the epidemiologic risk<sup>[7]</sup>.

The most common signs and symptoms are<sup>[8,9,14]</sup>:

- Fever, in 82% of patients (confidence interval (CI) 95%: 56-99%;  $n = 4410$ );
- Cough, in 61% of patients (CI 95%: 39-81%;  $n = 3985$ );
- Muscular pains/fatigue, in 36% of patients (CI 95%: 18-55%;  $n = 3778$ );
- Dyspnea, in 36% of patients (CI 95%: 18-55%;  $n = 3778$ );
- Sore throat, in 10% of patients (CI 95%: 5-17%;  $n = 1387$ );
- Headaches, in 12% of patients (CI 95%: 4-23%;  $n = 3598$ );
- Diarrhea and vomiting, in 12% of patients (CI 95%: 4-23%;  $n = 3598$ );

The symptoms may appear 3 to 6 days after the exposure to the agent<sup>[9]</sup>. Additionally, the patient may present dysgeusia, however rare, which justifies the tropism of the new coronavirus compared to the well-known SARS-CoV, MERS-CoV and A/ H1N1<sup>[11]</sup>.

Regarding the epidemiologic risk, the following aspects are taken into account (for every point below, consider 14 days following the beginning of the symptoms)<sup>[7]</sup>:

- History of international trips;
- Close contact (<2 m distance) with patients infected with SARS-CoV-2 (confirmed by lab test);
- Healthcare professionals having worked in an environment where patients suffering from acute respiratory infection were present;
- Quick deterioration of the health condition upon returning home;
- Presence in hospitals or health units in countries where the infection by SARS-CoV-2 was reported.

In imaging terms, the X-rays display bilateral opacities (similar to ground-glass) and a thickening of the pulmonary texture. In CAT scans it is possible to observe a thickening of the septum, ground-glass opacities and pulmonary consolidation<sup>[9,11,14]</sup>.

The affection of the respiratory system is directly related to its vulnerability to the inoculation of pathogens, associated to the contact of infected droplets with the oral, nasal or ocular mucosa, or even directly through the external airway<sup>[5]</sup>.

The difficulty of diagnosing this disease focuses on the fact that the clinical condition presents unspecific symptoms and, very often, indistinguishable from other respiratory pathologies. Symptoms like headaches, sore throat and gastrointestinal discomfort must not be neglected<sup>[14]</sup>.

The clinical condition of a patient infected with SARS-CoV-2 may range from the presence of a mild upper respiratory infection to SARS<sup>[9,13]</sup>. The severity of the clinical condition is predictive of a bad prognosis. The presentation of the disease is divided into not serious and serious cases. The serious cases are associated to old age, underlying comorbidities (high blood pressure, chronic obstructive pulmonary disease, diabetes), change of the laboratory values (leukopenia, lymphopenia, thrombocytopenia, high levels of C-reactive protein and lactate dehydrogenase, prolonged prothrombin time) and occurrence of complications (pneumonia, acute respiratory failure syndrome, acute respiratory distress syndrome, acute kidney and/or myocardium injury, secondary bacterial or fungal infections, septic shock and multiple organ failure)<sup>[8,9,11]</sup>.

In geographical areas where the virus of COVID-19 is disseminated throughout the population in general, a simple algorithm may be adopted, for instance, the triage by rRT-PCR of a simple target of a genetic sequence being enough. One or more negative results do not rule out the possibility of infection. There is a variety of factors which may lead to a false negative result: poor quality of the sample, containing scarce genetic material; collection at a very early stage of the infection; sample badly conditioned after its collection; specific reasons inherent in the technique, such as mutations of the virus or presence of inhibitors of the PCR. If a result is negative and, even so, there is a high suspicion of COVID-19, other samples must be collected and tested. The serological tests may prove useful in the aftermath of outbreaks and also in those cases where the tests by rRT-PCR come back negative but there is a strong epidemiologic connection to COVID-19. The samples collected at the acute stage of the disease and at the recovery stage may confirm the diagnosis (note that the production of immunoglobulins - IgG and IgM - starts days or weeks after the infection with the virus and the immune response depends on several factors, including age, nutritional status, severity of the disease and immunosuppression)<sup>[15-17]</sup>.

## **Oral manifestations**

### **Periodontal Disease**

Periodontal disease can be compared to a silent pandemic which pathophysiology is complex and multifactorial<sup>[5,6]</sup>. COVID-19 is classified as a pandemic, where the development of symptoms may be associated with a “cytokine burst”. This event involves an increase in the levels of inflammatory factors, namely interleukins (IL), gamma interferon and tumor necrosis factors <sup>[18,19]</sup>. In patients diagnosed with periodontal disease, there is an increase in the levels of IL-6 and IL-17, produced at the level of gingival tissues and with repercussions at the systemic level<sup>[18,19]</sup>. This “cytokine burst”, present in both pathologies, suggests a possible relationship between both, in order to understand whether the presence of one of them is a risk or predictive factor. That said, the need and importance of periodontal support consultations is reinforced<sup>[18,20,21]</sup>.

In Periodontal Disease, the supporting tissues are destroyed by bacterial infection, with the transition from a healthy sulcus to a periodontal pocket. When it comes to periodontal pocket, viral species are detected (herpes simplex 26-78%; epstein-barr 46-58%; cytomegalovirus 42-58%). This suggests that periodontal pockets provide an environment compatible with viral infection and survival<sup>[21,22]</sup>. In addition, gingival crevicular fluid can also constitute a reservoir for SARS-CoV-2<sup>[22,23]</sup>.

## Oral Injuries

The spread of COVID-19 correspond to the detection of a wide range of oral manifestations<sup>[24-27]</sup>. On this basis, concerns were raised about the possible causes of these injuries: whether caused directly by the SARS-CoV-2 virus or indirectly by the failure of the infected host's immune system<sup>[24-27]</sup>. Oral signs and symptoms associated with COVID-19 include changes in taste, unspecific oral ulcerations, desquamative gingivitis, petechiae, and opportunistic infections, such as infection by candida albicans. Oral lesions have different clinical aspects, varying in location, size, appearance of color and quantity<sup>[24]</sup>.

Abanoub Riad<sup>[28]</sup> and his co-workers published several cases, in which they suggest that the presence of oral mucositis in positive SARS-CoV-2 patients may be related to direct damage to cells infected by the virus or as an opportunistic infection resulting from immune dysregulation.

The name “COVID tongue” emerged in an attempt to categorize clinical characteristics associated with infection by SARS-CoV-2, a virus that is still unknown. However, the oral manifestations detected in the tongue are nonspecific and relatively common in the general population, namely glossitis, aphthous stomatitis, irregular depapillation (associated with lesions of the geographical tongue), therefore, this label should be interpreted cautiously in order to avoid false diagnoses<sup>[29-30]</sup>.

## Saliva

The functional receptor for SARS-CoV-2 is known to be the receptor for the angiotensin II-converting enzyme (ECA2). This receptor was found in the ducts of the epithelial cells of the salivary glands, especially in the minor glands. Therefore, the early contamination of the salivary gland ducts produces infected saliva, which can culminate in the appearance of oral symptoms prior to lung injuries<sup>[31-33]</sup>. In addition to this route of entry of the virus into the oral cavity, there are two others, the SARS-CoV-2 from the upper respiratory tract or bloodstream<sup>[34-35]</sup>.

Recent research has put saliva as a potential diagnostic tool for SARS-CoV-2 infection due to its easy access and interconnection with systemic diseases<sup>[32,36-38]</sup>.

Regarding the diagnosis of infection by SARS-CoV-2 via salivary samples, there are several advantages over the collection of nasopharyngeal samples. These include the fact that the technique is minimally invasive, easy, inexpensive, and the harvest can be performed by the individual himself (not requiring specialized personnel) and with a minimal risk of cross-infection<sup>[32-34, 37, 39-41]</sup>. In addition, the collection of saliva provides a sample of sufficient size to test different biomarkers<sup>[35]</sup>. However, the interpretation of the results published in the literature has to be performed with caution, since results with different sensitivity and detection rates have been revealed in symptomatic and asymptomatic patients. In another words, in asymptomatic patients, salivary samples are more sensitive and have a higher SARS-CoV-2 virus detection rate than nasopharyngeal samples. In symptomatic individuals the results are exactly the opposite<sup>[32]</sup>.

The collection of saliva can be performed in the posterior oropharyngeal region (the individual encourages the production of sputum, including the

collection of pharyngeal and not exclusively oral secretions); or collection of oral saliva from the salivary glands exclusively<sup>[40]</sup>.

Haya Altawalrah and her co-workers<sup>[42]</sup> conducted a cross-sectional study, whose objective was to evaluate the potential of salivary detection of SARS-CoV-2 by rt-PCR as an alternative to nasopharyngeal samples. The results suggest a high sensitivity (83.43%) and specificity (96.71%) for the salivary rt-PCR test when compared to the nasopharyngeal standard sample. These results corroborate those of previous publications. The kappa coefficient points to strong concordance analyzes for the diagnosis of SARS-CoV-2 via nasopharyngeal and salivary samples<sup>[42]</sup>.

László Márk Czumbel and his co-workers<sup>[43]</sup> in a meta-analysis, emphasize the need for standardization of procedures related to the collection and processing of salivary samples as a means of diagnosing SARS-CoV-2 infection, despite highlighting saliva as a promising method.

Another issue to be addressed in future investigations is related to the technique of collecting the salivary sample and whether different collection methods influence the sensitivity of the test for the detection of SARS-CoV-2 infection<sup>[34]</sup>. It should be noted that the technique of extracting viral RNA, detected viral load, the number of crops, which saliva to collect and when to collect it must also be standardized. For example, it is known that the viral load in saliva is higher in the first days of infection and that it tends to decrease exponentially when the virus already has pulmonary involvement<sup>[35]</sup>. There are studies that are beginning to highlight the potential diagnosis of samples of gingival crevicular fluid<sup>[44]</sup>.

## Routes of Contagion

The introduction into the human community occurred via wild animals, but currently the human to human in close proximity transmission path is the predominant one<sup>[7,8,12]</sup>.

It still hasn't been established for any respiratory virus the specific degree of infectiousness in terms of the proportion of the several transmission paths. There are multiple individual factors and environmental contexts which might contribute to the importance of each transmission path<sup>[45]</sup>. Up to date there is no evidence about the causality of the environmental presence of the virus (or viral RNA) and the risk of transmission to humans. This fact is partially related to the ignorance about the dose of

infectiousness of the virus. One of this virus' notorious features is the possibility of transmission by asymptomatic carriers (during the virus' incubation period, 2-14 days) or presenting a mild symptomatology<sup>[7,8,46-48]</sup>. The routes of contagion include direct contact with respiratory secretions, ingestion, inhalation of respiratory droplets (>5 µm, cough/sneeze), aerosols (through clusters of droplets <5 µm generated during the exhalation), fomites or orofecal<sup>[9,47,49-54]</sup>. The close contact increases the risk of transmission<sup>[8]</sup>. The possibility of transmission through aerosols exists, particularly, in enclosed spaces with exposure to high concentrations of aerosols for an extended period of time, something which potentially happens while practicing the profession in the sector of dentistry<sup>[50]</sup>.

## Treatment and Vaccine

Currently there are no pharmacological therapies of choice, nor any vaccine for immunization against the infection by SARS-CoV-2<sup>[7-9,12]</sup>.

The key to the treatment of the pandemic by SARS-CoV-2 is focused on the early diagnosis, isolation and monitoring<sup>[11]</sup>.

The containment measures include the restriction of the movement of persons (confinement to the home space), social distancing, interruption of labor activities in the several sectors, use of masks and frequent hand hygiene when there is social contact<sup>[12]</sup>.

The treatments are oriented towards the support, and include pharmacological therapy, additional oxygenation, non-invasive ventilation, invasive ventilation and extracorporeal membrane oxygenation<sup>[8,12]</sup>. The first line of intervention is to prescribe antipyretics (paracetamol) and expectorants (guaifenesin). In cases of acute respiratory infection, acute respiratory distress syndrome, hypoxemia and septic shock, the therapy is to administer oxygen [5 L/min], until the values of oxygen saturation in the blood are higher than 90% (children and adults) or than 92-95% (pregnant women)]. In conditions of acute renal failure, the monitoring of renal function and hydroelectrolytic balance are key. In cases of secondary bacterial and/or fungal infections the prescription of broad spectrum antibiotics is recommended<sup>[7]</sup>.

In April 2020, efforts to develop a SARS-CoV-2 vaccine grew exponentially and culminated in more than 100 proposals under study<sup>[55]</sup>.

Currently, the European Medicines Agency has approved the use of vaccines from BioNTech/Pfizer®, Moderna's® and AstraZeneca®. The Janssen vaccine is in the process of approval and Sanofi vaccines - GSK® and CureVac® have not yet started the authorization process.

The formulation of each type of vaccine depends on how the virus's genetic material is used - recombinant proteins, replicative or non-replicating viral vectors, DNA, mRNA, live attenuated vaccines or with the virus inactive<sup>[55-57]</sup>.

Vaccines provoke an immune response through specific antibodies in the long term, although existing studies are still insufficient to attest to this claim<sup>[58]</sup>.

It is important to note that within the concentrations of viral particles in the vaccine that can confer immunity, side effects must be taken into account. Thus, higher concentrations tend to have more pronounced side effects, and the effectiveness is maintained<sup>[55]</sup>. The most common side effects are fatigue, flu-like syndrome, vomiting, nausea, fever, arthralgia, headache, anaphylaxis - stressed the importance of vaccination<sup>[59-61]</sup>.

For vaccines already approved, their prescription consists of two doses at intervals of 21 days, 28 days or 12 weeks respectively for BioNTech/Pfizer<sup>®</sup><sup>[59]</sup>, Moderna's<sup>®</sup><sup>[60]</sup> and AstraZeneca<sup>®</sup><sup>[61]</sup>. The effectiveness of the three vaccines is very similar. The BioNTech/Pfizer<sup>®</sup> and Moderna's<sup>®</sup> vaccines suggest an efficacy greater than 90% after the second dose<sup>[62]</sup>, while the AstraZeneca<sup>®</sup> vaccine suggested a 70% efficacy in the initial studies. However, new publications show a greater effectiveness of this vaccine when applied with an interval of 12 weeks instead of 4 weeks<sup>[63]</sup>.

It is not correct to say that there is a vaccine that overlaps another, since the three approved vaccines have led to a decrease in severe cases of hospitalization and in cases of death by COVID-19.

However, it is necessary to maintain the preventive rules for compliance with social distance, respiratory etiquette and the use of a mask.

## **Impact of COVID-19 on the Dental sector**

The panorama of dental medicine has experienced major changes, not only economically, but also in the psychological field of the dental team. And, if before the pandemic, the dental medicine consultation was a safe place, with the adaptations introduced today, it is even more<sup>[64,65]</sup>.

Economically, this impact translates into an increase in the cost of personal protective equipment<sup>[66]</sup> and a decrease in the number of consultations given. Since then, the appointment times were increased, in order to consider a period of

disinfection and ventilation of the office<sup>[67]</sup>. On the other hand, the expectation of a possible economic crisis often culminates in postponing treatments<sup>[68]</sup>.

The challenges of this new era have forced dentistry and medicine to reinvent itself. This adaptation includes the introduction of new technologies, so that the interests of patients are protected and safeguarded. Telemedicine is an example of this reinvention.

## Telemedicine

Telehealth consists of using electronic and technological strategies to provide long-distance clinical support and promote the education of patients and professionals<sup>[69]</sup>. Telemedicine is defined as the use of video conferencing for diagnosis and medical advice about a certain pathology<sup>[69,70]</sup>. These concepts are not new and date back to 1997. Nevertheless, the lockdown has forced the world to, as a result of the pandemic by COVID-19, allow the rapid development of this medical approach<sup>[69]</sup>. The word “tele” means distance, so telemedicine presupposes compliance with one of the most important measures to prevent the spread of the virus: social distance<sup>[71]</sup>.

There are several advantages in the area of Telemedicine, namely: Teleconsultation; Telediagnosis, Teletriaging and Telemonitoring<sup>[71]</sup>.

COVID-19 has a real impact on oral health, not only because of the role that salivary glands play as potential reservoirs of the SARS-CoV-2 virus, but also because of the injuries that can manifest in the oral cavity of infected individuals. In the latest publications that relate oral health to infected individuals, vesicle-bullous and maculopapular lesions are described<sup>[72,73]</sup>. So, the early detection of these lesions by an oral medicine specialist can play a key role in screening individuals suspected of being infected with SARS-CoV-2<sup>[11,72]</sup>.

The systems available for the application of Telemedicine are still restricted, but they represent an easily accessible, viable and useful tool for both healthcare professionals and patients. This communication can be via messaging applications (WhatsApp, Instagram, SMS, Messenger) or video calling systems (WhatsApp, Skype, FaceTime)<sup>[72]</sup>. Petruzzi et al.<sup>[74]</sup> confirmed the 82% agreement between the diagnosis of cases by teleconsultation and clinical and pathological examination.



One of the specialties that will most benefit from this remote system is Orthodontics, especially in appliance control appointments<sup>[75]</sup>. Still, it is an area with limitations as it is often dependent on the quality of the image that patients provide. For a correct diagnosis, the images must be representative of the entire region of the mouth to be analyzed, as well as without any filter<sup>[72]</sup>.

It should be noted that Telemedicine is not a substitute for face-to-face appointment because it has a palliative approach<sup>[72]</sup>. However, once explored and well implemented, it is a complement to the practice of conventional dentistry<sup>[66]</sup>. In pandemic situations, Telemedicine prevents the spread of the virus among healthcare professionals and patients<sup>[16]</sup>.

## **Context of the consultation of Dentistry and the need for these Clinical Guidelines**

The dentists are, in countless situations, the first line of diagnosis<sup>[6]</sup>. Moreover, they represent the most exposed professional class and the one at greater risk of infection by SARS-CoV-2 since the majority of the procedures they perform imply the generation of aerosols<sup>[12]</sup>.

The saliva contains a very high viral load of SARS-CoV-2 in infected patients and evidence shows that it can survive up to 9 days on surfaces and items that have come into contact with contaminated oral fluids<sup>[8-12]</sup>.

The limitation of the sector's activity represents not only a significant economic impact, but it also affects oral health and, subsequently, the general health of the community<sup>[12]</sup>.

The first guidelines made public well into the pandemic restricted the dentists' activity to urgent and immediate consultations referring that, in each case, the solution for the problem should involve the non-generation of aerosols.

In spite of the countless efforts to publish and disseminate the guidelines of post-pandemic intervention, the information concerning SARS-CoV-2 is very volatile and dynamic, wherefore it is the health professionals' duty to keep updated.

It is a challenge to the community, and not only in the context of dentistry, where every measure is oriented towards the prevention and control of the virus' dissemination, their main goal being the protection of each and every one involved in the different

sectors (patients, receptionists, dental assistants, dental prosthetic technicians, dentists, administrative staff)<sup>[6-12]</sup>.

The need for this paper arises from the inexistence, nationwide, of Clinical Guidelines based on the evidence regarding this problem.

# Materials and Methods

## Questions

(Appendix II, section 1)

## Bibliographic Research

A systematic research of the *MEDLINE* (via *PubMed*), *Cochrane Library* and *Trip Database* databases was conducted, with known technical limitations, during the period ranging between March and April 2020. The search included the following keywords individually and in different combinations (Appendix II, section 2): “SARS”; “MERS”; “SARS-CoV-2”; “COVID-19 OR Covid-19 OR coronavirus\*”; “influenza”; “dentistry”; [“ventilator” AND (“oral hygiene” OR “dentistry”)]; [(“aerosol\*” OR air) AND “SARS”]. In a second phase we cross-checked the results of the first research with the algorithm created by the *National Institute for Health and Care Excellence* (NICE)<sup>[1]</sup>.

Additionally, we conducted a manual research on websites (*researchgate.net*, *Dental Blog*, *Evidentia Medica*, *thenew.dentist*) e online newspapers (*Influenza Journal*, *ABSA – The Association for Biosafety and Biosecurity*, *American Society for Microbiology*, *International Journal of Oral Science*, *Clinical Infectious Diseases*, *Nature*, *The Lancet*, *JAMA Network*, *Frontiers of Oral and Maxillofacial Medicine*, *International Journal of Environmental Research and Public Health*, *Science*, *Interface Focus*, *The Centre for Evidence-Based Medicine*, *British Microbiology Research Journal*, *The Chronicle of Higher Education*, *British Dental Journal*).

In parallel to this, the authors of the included studies were contacted, and the lists of references were reviewed, with the purpose of looking into the publication of more articles within the framework of the Clinical Guidelines.

In spite of the systematic research, the main information stems from the use of guiding documents previously published regarding pandemics already occurred (SARS-CoV, AH1N1 and MERS-CoV).

The databases of choice were:

1. Guides of the World Health Organization (WHO) on COVID-19
2. Recommendations of the Portuguese Health Authority (DGS)
3. NICE e SIGN

4. Cochrane collaboration
5. MEDLINE (via PubMed)
6. International and national scientific societies
7. The available evidence, obtained through the search strategy that is reproduced in Appendix II, was simply ranked
  - Priority has been given to systematic reviews and meta-analysis
  - Followed by the search for randomized clinical trials
  - Followed by observational studies
  - Whenever it proved impossible, we searched for publications arising from the observation of experts in previous experiences
  - In those cases where no kind of evidence can be found, the experts who are the authors of these guidelines drew up the recommendations through discussion and consensus based primarily on biological plausibility.

## Inclusion/Exclusion Criteria

The following have been defined as inclusion criteria (Appendix II, section 3):

**Patients (P):** patients infected or not with the virus SARS-CoV-2.

**Intervention (I):** any type of dental procedure, with or without the generation of aerosols, in a context of pandemic by SARS-CoV-2.

**Comparison (C):** any type of dental procedure, with or without the generation of aerosols, before the pandemic by SARS-CoV-2.

**Outcomes (O):** control of SARS-CoV-2 virus transmission; monitoring of patients infected with SARS-CoV-2; prevention measures; disinfection measures; risk of the procedures carried out in dentistry in the contraction of the infection by SARS-CoV-2; training for the health professionals and administrative staff for the new post-pandemic situation; interpersonal contact at times of pandemic by SARS-CoV-2; which personal protective equipment (PPE) to wear, its effectiveness and how to wear it.

The following have been defined as exclusion criteria (Appendix II, section 4):

**Criterion A:** articles focused on diagnosis (for instance, diagnosis of respiratory changes)

**Criterion B:** creation of a Clinical Database

**Criterion C:** vaccination

**Criterion D:** virus' immune mechanisms/pathogenicity/immunity

**Criterion E:** other pathologies

**Criterion F:** triage of opting for hospital treatment/treatment at home

**Criterion G:** genetic studies/genome

**Criterion H:** pharmacology

**Criterion I:** articles written in languages other than Portuguese, English, Spanish and French

**Criterion J:** others that are not included in the previous ones

## Selection and Assessment of Evidence

The research, the triage by title, the abstract and the full analysis of the text of the article were conducted/written by two proofreaders (AV; CM). Any disagreements have been solved through direct confrontation in the form of oral discussion. In situations where an agreement could not be reached, resort was had to a third proofreader (AM).

We have made a summary critical assessment of the literature, but we have not formally assessed the risk of the presence of biases resorting to specific scales.

However, we have always briefly and simply indicated the kind of information underlying the clinical recommendations produced by us.

## Ranking of Evidence and Class of Recommendation


The objective of the Clinical Guidelines lies on formulating recommendations based on the evidence which has clinical applicability. The levels of evidence (Appendix II, section 5) rank the different kinds of articles depending on the intrinsic methodology, while the classes of recommendation (Appendix II, section 5) rank the force of the action. In this context, it is expectable that strong recommendations present less frequent variations. The review of scientific evidence allows to gage the ratio risk/benefit of a given action<sup>[3]</sup>.

## Research Results

After duplicates have been removed, the research of the databases resulted in a total of 163 152 references, of which 162 617 did not match the inclusion criteria, being consequently ineligible, and were classified as excluded. We obtained the full articles of the 640 potentially eligible references, selected through systematic and manual research.

After the assessment, 253 studies were excluded for the reasons described in Appendix I, section 4.

Figure 2 - Result of the Systematic Research



# The Pre-Intervention Stages

## The Schedule

### The management of the appointments

The dental health care personnel main concern is the protection of the patient as well as of all those involved in the management and operation of the clinic in a safe way, keeping the levels of effectiveness of treatment and meeting the patient's demands<sup>[76]</sup>. (*Level III, Class IIa*).

---

#### 1.1. How can you manage the schedule?

**Answer:** Every appointment must have a scheduled time (now more extended, due to the pandemic) suited for every procedure, which must be indicated by the dentist. In case a significant delay is identified, the following patient must be contacted in order to avoid his/her permanence in the waiting room. The appointments must be scheduled with an interval long enough to minimize possible contacts between patients in the waiting room. It must be taken into consideration that the procedures to be carried out in between consultations will take longer so that they can be duly performed (long enough for airing, disinfecting the surfaces, payment logistics and scheduling the following appointment)<sup>[77-89]</sup>. (*Level IV, Class IIa*)

---

#### 1.2. Should the patient be alerted to be on time for the appointment?

**Answer:** Yes, the patient must be right on time for the appointment in order to avoid the gathering of patients in the waiting room. Should the patient arrive in advance, he/she can wait for the time of the appointment near the clinic. In case the patient is late, it is up to the dentist (or the duly trained administrative staff) to decide whether the appointment will be postponed/cancelled<sup>[90,91]</sup>. (*Level III, Class IIa*)

---

1.3. Should the patient be alerted to avoid bringing personal ornaments to the appointment?

**Answer:** Yes, the patient must be alerted not to wear any personal ornaments like watches, earrings, bracelets, necklaces, rings<sup>[90,92]</sup>. One of the main routes of contagion are fomites, in other words, any item or inanimate surface capable of absorbing, retaining and transporting infectious organisms. SARS-CoV-2 has a survival time of up to 24 h on paper, 48 h on steel and 72 h on plastic, therefore the restriction of items unnecessary for the appointment represents the elimination of a transmission path of the virus<sup>[12, 93]</sup>. *(Level III, Class IIa)*

---

1.4. Should the patient be alerted to avoid paying the consultation in cash?

**Answer:** Yes, the patient must be alerted to pay using, preferably, the POS terminal or any other electronic means<sup>[90,95]</sup>. *(Level IV, Class IIa)*

---

1.5. What are you supposed to do if a patient asks for a teleconsultation?

**Answer:** The option of providing a teleconsultation is the sole responsibility of the doctor and the clinic. Given the specificity of the necessary treatments in most specialties of dentistry requiring the presence and contact with the patient, the decision must be made case-by-case. The digital consultation may be established by electronic mail or video through several digital platforms, through applications for cellphones, among others, always respecting the patient's privacy as if it were a face-to-face consultation<sup>[95-98]</sup>. *(Level III, Class II)*



## ***Key Points***

- SARS-CoV-2 positive patient can attend the consultation after authorization from the Local Health Authority, only in urgent and urgent cases. ***(Level IV, Class IIa)***
- The appointment time must be adhered to, in order to avoid crowding patients in the waiting room. ***(Level III, Class IIa)***
- The use of personal ornaments by the patient must be restricted. ***(Level III, Class IIa)***
- The payment of the consultation must be made using, preferably, a POS terminal or electronic means. ***(Level IV, Class IIa)***
- The option of teleconsultation is the doctor's and clinic's sole responsibility, being assessed case-by-case. ***(Level III, Class II)***

## Questionnaire to be completed

---

2.1. Which questions must be asked while making an appointment by telephone?

**Answer:** The following questions must be asked:

- Do you have (or have you had) fever over the last 14 days? (temperature  $\geq 38$  °C)
- Do you have (or have you had) cough or trouble breathing over the last 14 days?
- Do you have/experience (or have you had/experienced) a general discomfort, physical fatigue, diarrhea or headaches over the last 14 days?
- Do you complain from (or have you experienced) a total or partial loss of smell or taste over the last 14 days?
- Does any of your family members or close contacts suffer from (or has any of them referred) any of those symptoms over the last 14 days?
- Have you been in contact with anyone suspect or having tested positive for COVID-19?
- Have you already been diagnosed with COVID-19? If the answer is affirmative: Are you still in isolation period (14 days)?

If any of the answers is affirmative, the doctor must be informed about the health condition and weigh up the urgency of the patient's treatment<sup>[12,17,54,90,99-109]</sup>. (*Level IIb, Class I*)

---

2.2. If the patient answers in the affirmative to any of the questions included in the questionnaire referred to in 2.1, but refers absence of fever, how is scheduling supposed to be handled?

**Answer:** Some patients who have tested positive for SARS-CoV-2 may present cough and other symptoms, but no fever or the temperature rise being at the lower threshold (<38 °C), particularly in the first days. If the treatment is not urgent, the appointment must be postponed by 2 weeks and the telephone triage must be repeated while confirming the appointment. If it is an urgent situation, you should deal with it as if it were a case of infection with SARS-CoV-2<sup>[54,109]</sup>. (*Level III, Class I*)

---

2.3. While scheduling an appointment for a patient having a professional profile of risk, as is the case of healthcare workers, or hospital environment, day care center or retirement homes employees, should there be any particular care?

**Answer:** Every mentioned patient, in spite of belonging to a risk group, will be submitted to the same protocol of epidemiologic triage as any other patient and treated according to the standards of individual protection in force for every dental procedure<sup>[110]</sup>. (*Level IV, Class IIb*)

---

2.4. The patient refers having been infected with SARS-CoV-2 but he/she have already tested negative since. How is one expected to proceed?

**Answer:** If the patient has not had a fever for at least 3 days (without using antipyretics), has significantly improved cough, fatigue, and shortness of breath, and has been in isolation at home for at least 10 days in asymptomatic, mild or moderate cases, or 20 days in the most severe cases, you do not need to present a negative laboratory test for SARS-CoV-2 and you can go to the consultation<sup>[111-113]</sup>.

For professionals at high risk of exposure such as health professionals or caregivers of close proximity to vulnerable people, the previous criteria should add the need for a negative test for SARS-CoV-2 with a test (rRT-PCR for SARS-CoV-2) If the laboratory test result for SARS-CoV-2 is positive, the isolation is prolonged until 20 days have elapsed since the onset of symptoms, at which point the isolation end is determined, without the need for additional testing. laboratory testing. (*Level IV, Class IIa*)

---

2.5. How should one tackle the situation of a patient having a clinical history of COVID-19 and who is in home isolation?

**Answer:** AA patient undergoing a period of isolation due to COVID-19 in need of urgent dental treatments can only leave the isolation after authorization by the Local Health Delegate. This clearance is the only one allowing the patient to visit the clinic or dental practice and to be treated according to the standards of individual protection in force for every dental procedure<sup>[113-115]</sup>. In the imperative need to observe a suspected or

confirmed case of COVID-19, in urgent situations, consider scheduling the appointment, at the end of the morning or in the afternoon, at specific times, so that there is no sharing of the waiting room, always consulting the patient with the use of PPE for high-risk procedures<sup>[95]</sup>. (*Level IV, Class IIa*)

---

2.6. A patient goes spontaneously to the clinic to schedule an appointment without having previously called. What is the receptionist supposed to do?

**Answer:** It must be explained that going to the clinic without a previous contact must be avoided. After carrying out the epidemiologic triage, the need of an urgent consultation should be assessed<sup>[90]</sup>. (*Level IV, Class IIa*)

---

2.7. Should the accompanying person be advised not to come, whenever possible, and/or avoid his/her permanence in any place of the clinic?

**Answer:** Yes, the presence of the accompanying person is only indicated when the patient is underage, needs help to move around or depends on others<sup>[71,91]</sup>. (*Level IV, Class IIa*)

---

2.8. If the patient comes accompanied, is it necessary for the accompanying person to go through the triage for COVID-19?

**Answer:** Yes, since this person may be a suspect case<sup>[17,90]</sup>. (*Level III, Class I*)

## ***Key Points***

- In case of the patient's suspected infection with SARS-CoV-2, the dentist must be informed, as well as weigh up the urgency of the treatment of the patient in question. ***(Level IIb, Class I)***
- If the patient no longer has a fever for at least 3 days (without the use of antipyretics) have significant improvement in cough, fatigue, and shortness of breath, and was in home insulation at least 10 days in asymptomatic cases, mild or moderate or 20 days in the most severe cases, you do not need to submit a negative laboratory test for SARS-CoV-2 and you can go to the consultation. ***(Level IV, Class IIa)***
- In patients positive to SARS-CoV-2 in home isolation, even without fever, every nonurgent dental treatment must be postponed by two weeks, being reassessed afterwards. ***(Level III, Class I)***
- In patients positive to SARS-CoV-2 in home isolation, the treatment is only ensured in urgent situations and subject to an authorization by the local Health Delegate. ***(Level IV, Class IIa)***
- The patient should avoid going to the clinic without previously contacting the reception. ***(Level IV, Class IIa)***
- The presence of accompanying person is only indicated for patients underage or requiring special needs, namely elderly people, pregnant women, people presenting visible physical or mental limitations. ***(Level IV, Class IIa)***
- The accompanying person must be asked, all the same, the questions related to the screening of infection with SARS-CoV-2. ***(Level III, Class I)***
- Every individual (patients/accompanying people) presenting an acute respiratory condition of cough, fever or dyspnea/respiratory difficulty is considered suspect of infection with SARS-CioV-2. ***(Level III, Class I)***

# Reception, Waiting Room and Sanitary Facilities

## Organization of space and furniture

---

3.1. How can you make means of disinfection and protection available to those entering the clinic?

**Answer:** Install alcohol-based antiseptic solution (ABAS) dispensers near the entrance of the clinic and require its use (through an infographic). Make shoe covers, bouffant cap (optional) and surgical masks available to those who are going to remain inside the clinic<sup>[12,17,116]</sup>. (*Level III, Class I*)

---

3.2. Which special measures must be applied in this customer service area?

**Answer:** Allocate a pen to each employee and another one for the exclusive use of the patients (which must be disinfected after every single use, using either a wipe or disinfectant spray). In case of coming into contact with items belonging to the patient or with potentially contaminated surfaces (cards, bills, coins, cell phone, among others), you should perform the antiseptics of the hands<sup>[90,91,117,118]</sup>. (*Level IV, Class IIa*)

---

3.3. How should you organize the waiting room?

**Answer:** The waiting room must be organized in such a way that the available seats are two meters apart from one another in order to guarantee social distancing. If every occupant wears a mask, the distance may be shorter, making sure that there is no physical contact between them<sup>[115]</sup>. The patient must wait seated until his/her turn comes. In case the patients prefer so, or when the waiting room cannot accommodate any more users, they must be advised to wait inside their vehicles or nearby (ensuring social distancing), being contacted by phone when it is their turn<sup>[119,120]</sup>. (*Level III, Class I*)

---

3.4. How many people can remain in the waiting room? What is the advised capacity?

**Answer:** One must make sure that the place where users are to wait only accommodates 1/3 of its usual capacity<sup>[116,120]</sup>. One must expedite procedures so that the least possible number of people seat in the waiting room and during the shortest time possible<sup>[116,119]</sup>. *(Level IV, Class IIb)*

---

3.5. Which furniture can there be in the waiting room?

**Answer:** In the waiting room there must only be the furniture which is strictly necessary to accommodate the patients, one that is easily washable and disinfectable, and a waste container to store the disposables worn by the patients. It is recommended to remove every decorative furniture such as paintings, sofas, support tables, flower boxes, among others<sup>[116,120]</sup>. *(Level III, Class I)*

---

3.6. Should magazines or newspapers be available?

**Answer:** Preferably, every newspaper, magazine and toy must be removed<sup>[120]</sup>. *(Level III, Class I)*

---

3.7. Should the TV and air conditioning remote controls be available?

**Answer:** No, these must be out of public's reach<sup>[118,121]</sup>. *(Level IV, Class IIa)*

---

3.8. Should the water dispenser or the coffee machine be available?

**Answer:** No, they must be removed since they may be a source of contagion<sup>[90,121]</sup>. *(Level IV, Class IIa)*

---

3.9. Which changes must be made in the sanitary facilities?

**Answer:** The towels and hand driers must be replaced by paper towels dispensers. Every non-essential element must be removed<sup>[90]</sup>. (*Level IV, Class IIa*)

---

3.10. Which means of decontamination must be available at the sanitary facilities?

**Answer:** Liquid soap and ABAS must be available<sup>[88,120]</sup>. (*Level IV, Class IIa*)



## ***Key Points***

- Make alcohol-based antiseptic solution available near the entrance and make surgical masks and shoe covers available to the individuals who are going to remain inside the clinic. ***(Level III, Class I)***
- Each employee has his/her own pen; there is a pen for the exclusive use of the patients, which must be disinfected after every single use. ***(Level IV, Class IIa)***
- Contact with objects belonging to the patient: sanitize the hands with antiseptic solution. ***(Level IV, Class IIa)***
- Do not share the telephone without it being previously disinfected. ***(Level IV, Class IIa)***
- Only the furniture which is strictly necessary must remain in the waiting room. ***(Level III, Class I)***
- The waiting room must allow for social distancing and must not exceed 1/3 of its maximum capacity. ***(Level III, Class I)***
- The TV or air conditioning remote controls must be out of public's reach. ***(Level IV, Class IIa)***
- The water dispensers and coffee machines must be removed since they are a potential source of contagion. ***(Level IV, Class IIa)***
-

## Physical Barriers and Signage

---

4.1. Should there be physical barriers between the reception and the user?

**Answer:** One must make sure that customer service is provided through physical barriers which limit the proximity between employees and users (for instance, the installation of an acrylic barrier which might limit exposure)<sup>[88,90,116]</sup>. One must make sure that customer service is provided while keeping an appropriate distance of, at least, 1 m by means of signaling to this effect - namely through signage and markings on the floor (Level IV, Class IIa)

---

4.2. Which signposting can indicate the organization of the waiting spaces or the interdiction of crossing without permission?

**Answer:** Signage and/or markings on the floor delimiting the safety area around the reception desk of, at least, 1 m must be used. Visible notices at the entrance and on the counter raising awareness about the distancing<sup>[116,122]</sup>. (*Level III, Class IIa*)

---

4.3 Which informative billboards on COVID-19 must be posted and where?

**Answer:** Some infographics explaining the hand washing<sup>[89,116,122]</sup> must be posted near the ABAS dispensers, in the waiting room, at the reception and in the sanitary facilities. Infographics explaining the respiratory etiquette<sup>[89,90,116,120]</sup> must be posted at the entrance of the facilities, in the waiting room, in the sanitary facilities and at the reception. At the entrance of the facilities, there must also be available infographics explaining how to put on and remove a surgical mask. (*Level IV, Class IIa*)

## ***Key Points***

- The safety distances and expected conduct of patients and staff must be properly posted at the clinic's entrance and on the counter. ***(Level III, Class IIa)***
- The customer service must be provided while keeping a distance of, at least, 1 m, alongside with the implementation and application of physical barriers. ***(Level IV, Class IIa)***
- Information about respiratory etiquette, hand sanitization and donning/removal of the mask must be posted. ***(Level IV, Class IIa)***

## Useful Information

---

### 5.1. What is the rationale behind hand washing?

**Answer:** Hand hygiene has been considered the most effective, simplest and cheapest single measure to reduce the risk of cross infection of microorganisms<sup>[89]</sup>. The virus may persist on several types of surface from hours to days, depending on the type of material and the environmental conditions (temperature and humidity), which implies the importance of a thorough disinfection of surfaces in a dental clinic/practice and dissemination of routine practices of hand hygiene and asepsis, both on the part of the health professionals and of the administrative staff and users<sup>[17,122]</sup>. It is fundamental to establish a protocol of adherence to these hygiene standards because they are recurrently neglected by the health professionals, as reports a systematic review referring adherence rates lower than 20%<sup>[123]</sup>. Another systematic review has determined the effectiveness of physical interventions in containing the contagion of respiratory viruses<sup>[71]</sup>. 32 studies were included regarding the topic of hand washing (12 RCT and 16 of observational nature), in several environments like health units, schools, day care centers, military conscription centers, fraternities/sororities and households. The highest methodological quality RCTs have determined a high effectiveness of hand washing in the prevention of contagion of respiratory viruses, especially in children. We can infer from the meta-analysis derived from the studies of checking cases that the implementation of simple and cheap hygienic measures is effective in the containment of SARS-CoV. The review concludes that frequent hand washing, with or without the adjuvant use of antiseptics, is effective in the reduction of viral transmission and that it must, preferably, be combined with other interventions of containment, such as the use of protective equipment and the triage of suspect cases. The public sanitary measures seem to be highly effective, especially if they are part of a structured program which includes simultaneous information and education. In the future, it will be of the essence to fill in some gaps displayed by the evidence through broad pragmatic trials, in order to assess the best combinations of sanitary and hygienic measures both in the community and in a private and hospital clinical setting, regarding several respiratory viral agents, focusing on SARS-CoV-2.

Hand hygiene has been considered the most important single measure to prevent the transmission of microorganisms. However, in order to optimize its effectiveness, this measure must be included in a general protocol of asepsis and spread through an

educational program of promotion of enabling habits and practices, both through training actions destined for health professionals, diffusion through information leaflets (or others considered suitable) and motivational reinforcement for the users, as well as through the implementation of physical devices integrated in the premises of dental clinics. *(Level Ia, Class I)*

---

## 5.2. Which products must be used to sanitize the hands?

**Answer:** It is recommended the use of the technique described by the WHO, following their recommendations for the 5 moments of hand hygiene. The leaflet and/or billboard explaining the hand washing must be posted near the areas provided for that purpose, namely sanitary facilities and dental office<sup>[116,122,125]</sup>. The antisepsis of the hands is a prerequisite for the dental practice; however, besides the health professionals at the clinic, it must eventually be extended to the administrative staff and users. The practice of the proper hand hygiene must be reinforced and carried out in an attentive and conscious way. The technique described by the WHO implies 6 chronologically chain-linked steps for the hand washing (40 to 60 s) or antiseptic friction of the hands (20-30 s): washing the palm of the hands – back with fingers interwoven – palm with fingers interwoven – fingers – thumbs – pulp of fingers. The health professionals in a clinic should adhere to the WHO's recommendations called "My 5 moments of hand hygiene": before touching a patient, before the cleaning or aseptic procedures, after risk or exposure to body fluids, after touching a patient and after touching around the patient<sup>[126]</sup>. *(Level Ib, Class I)*  
(Appendix III, section 1)

---

## 5.3. Which products must be used to sanitize the hands?

**Answer:** Hand hygiene with ABAS is universally used as the most effective, simplest, safest and cheapest procedure to fight the transmission of COVID-19. When hands are dirty or visibly contaminated with protein matter or blood, you must use, alternatively, water and soap. In the WHO's guidelines on hand hygiene in health care<sup>[126]</sup>, they develop an extensive narrative review about the effectiveness and efficiency of the several preparations for this purpose, through several experimental studies (in vitro and in vivo assays and clinical studies). It is established that ABAS is the preparation of choice for a routine antisepsis of the hands, with the highest level of evidence (Ia). Through the

denaturation of proteins, the alcohol inactivates the virus with envelope (lipophilic), including the coronaviruses, and it is effective with a concentration of ethanol of, at least, 60%. In a systematic review about the effectiveness of ABAS, it was concluded that it removes microorganisms in a more effective way, in less time and with a lower irritant capacity than washing the hands with water and soap or other antiseptic agents<sup>[127]</sup>. ABAS' formulations containing 80% (v/v) of ethanol or 75% (v/v) of isopropanol are virucidal for SARS-CoV and MERS-Cov<sup>[126]</sup>. In a clinical context, it is fundamental to comply with the recommended washing time and volume of product in order to achieve a real effectiveness. The product's ideal volume to be applied is still not totally defined and may vary depending on the formulations. However, as a general rule, if you experience a sensation of dry hands after washing them during 10-15 s, it is likely that you have applied an insufficient quantity of product. Nonetheless, alcohols are not good cleaning agents and their use is not recommended when hands are dirty or visibly contaminated with organic matter or blood. Under the circumstances, you must use water and soap, whose effectiveness is well established with a high level of evidence<sup>[89,126,128,129]</sup>.

As a future recommendation for research, we must implement new adaptations to the experimental methodology of the antimicrobials for the asepsis of the hands. Besides the in vivo laboratory assays of effectiveness, it is key to conduct effectiveness testing that can be extrapolated to real-life conditions of application in a clinical setting. (*Level Ia and Ib, Class I*)

---

#### 5.4. Which are the methods that keep hands' skin healthy?

**Answer:** In spite of the high clinical safety of ABAS' formulations, their frequent use for the antiseptics of the hands may cause skin dehydration. There are several strategies intended to minimize skin irritation or dermatitis: selecting products which are not as irritant, avoiding noxious practices and using skin care moisturizing products. It is estimated that the hands' occupational dermatitis may affect 30% of the health professionals. However, the shift towards a routine use of ABAS' formulations improves these professionals' skin health if some care is applied<sup>[117,130]</sup>. Several clinical studies have shown that the use of moisturizers or other conditioning agents, such as ABAS's excipients, is associated to a lower skin irritation compared to simple or antimicrobial soap. The addition of 1-3% of glycerol may minimize this side effect. You must also avoid noxious practices, like washing regularly your hands with water and soap immediately before or after using

ABAS, as well as put on gloves while your hands are still wet after washing them, regardless of the product that has been used. Health professionals at risk of developing contact dermatitis might benefit from the complementary use of emollient or moisturizing creams<sup>[126]</sup>. (*Level Ib, Class I*)

---

5.5. Should the patients be allowed to brush their teeth, prostheses or removable braces before stepping inside the consultation office?

**Answer:** That must not be allowed, in order to avoid cross contamination, being recommended the existence of some information concerning this prohibition in the sanitary facilities<sup>[120]</sup>. (*Level IV, Class IIa*)

## ***Key Points***

- Hand hygiene using alcohol-based antiseptic solutions (ABAS) is used worldwide as the most effective, simplest, safest and cheapest procedure at fighting the transmission of COVID-19. When hands are dirty or visibly contaminated, you must use water and soap. ***(Level Ia and Ib, Class I)***
- We recommend the use of the technique described by the WHO, following their recommendations for the 5 moments of hand hygiene. ***(Level Ib, Class I)***
- Patients must not brush their teeth nor sanitize prostheses or removable braces in the clinic's sanitary facilities. ***(Level IV, Class IIa)***



## Cleaning the Waiting Room and the Sanitary Facilities

---

### 6.1. How often should you clean?

**Answer:** The facilities must be cleaned and disinfected, at least, once a day, using appropriate agents, in every customer service area, waiting room and corridors leading to the consultation offices. The spaces where children might be must be cleaned more often<sup>[131,132]</sup>. (Level IV, Class IIb)

### 6.2. How should you clean surfaces?

**Answer:** The cleaning must always be wet - do not use dry vacuum cleaners in public areas, unless they are equipped with a tank which collects the dirt in the water; this reservoir must be emptied and washed after every single use<sup>[91,121,131]</sup>. (Level IV, Class IIb)

### 6.3. How should you do this cleaning?

**Answer:** It must always be done from top to bottom and from the cleanest areas to the dirtiest ones<sup>[121,131]</sup>: (Level IV, Class IIb)

1. Walls (where indicated);
2. Equipment existing in the areas;
3. Surfaces above the floor (working benches, tables, chairs, handrails and others);
4. Washbasins;
5. Floor.

### 6.4. Which cleaning/disinfection products must be used?

**Answer:** The cleaning products must contain disinfectants certified for use in a hospital/ clinical setting and comply with EN 14476 for virucidal products, and they must be applied according to the manufacturer's instructions.

As an option for those less critical surfaces, bleach-based products (sodium hypochlorite) having, at least, 0,1% of free chlorine may be used; in practice, it equals 20,5 ml of common bleach (5%) for every liter of water, which must not be prepared until they are

about to be used. It is important to leave the bleach for 10 min, rinsing it afterwards<sup>[118, 131]</sup>. (Level IV, Class IIb)

---

#### 6.5. What are the most critical surfaces?

**Answer:** The most critical surfaces are keyboards, mouse, POS terminal, door handles, the reception counter, hand rails and disinfectant dispensers<sup>[118]</sup>. (Level IV, Class IIb)

---

#### 6.6. How often should the most critical areas or surfaces be cleaned?

**Answer:** The cleaning frequency of the surfaces which are frequently touched must be, at least, once an hour or after every single use<sup>[133]</sup>. (Level IV, Class IIa)

---

#### 6.7. What are the materials you must use while cleaning the most critical areas?

**Answer:** The cleaning cloths must be, preferably, of single use and disposable, or wipes moistened with disinfectant and supplied by their own dispenser.

Use a wipe for every surface and then throw it away; do not dry the surface out after using the disinfectant wipe because you need that one to be wet for some minutes until it dries in the air in order to be effective<sup>[121,131]</sup>. (Level IV, Class IIa)

---

#### 6.8. How often must you clean the sanitary facilities?

**Answer:** Draw up a cleaning and sanitizing plan for the facilities, which must be posted in a visible place. This must be done, at least, 3 times a day, or whenever it proves necessary<sup>[131,132]</sup>. (Level IV, Class IIa)

---

#### 6.9. How should you clean the sanitary facilities?

**Answer:** While cleaning the sanitary facilities, the sequence below must be followed<sup>[121, 131]</sup>: (*Level IV, Class IIa*)

1. Clean the walls (where indicated);
2. Start the cleaning with the basins (first the taps and only then the basin) and the surfaces around these;
3. Clean the toilets focusing on the internal cleaning, leaving the product for 5 min, scrub with a brush and activate the flush with the brush still inside, so this one might be cleaned as well. In addition, disinfect properly the flush button;
4. Clean the floor;
5. At the end, disinfect the taps and the doorknobs of the restrooms once again.

---

#### 6.10. Which are the most critical surfaces in the sanitary facilities?

**Answer:** In the sanitary facilities, the most frequently touched surfaces are the dispensers (both of hand paper towels and of soap or disinfectant solution), taps, door handles, basins and toilet<sup>[132,134]</sup>. (*Level IV, Class IIa*)

---

#### 6.11. Which products must be used to clean the sanitary facilities?

**Answer:** For the critical surfaces, certified medical-hospital products (EN14476) must be used; for the remaining surfaces, the most widely used detergents are the common ones or the ones for domestic use:

- 70 % ethyl alcohol solution;
- Sodium hypochlorite containing, at least, 5% of free chlorine in its original form diluted in water.

The solution of diluted bleach must have a concentration of sodium hypochlorite of, at least, 0.1%. In practice, it equals 20.5 ml of common bleach for every liter of water, and it must not be prepared until it is about to be used. One must leave the sodium hypochlorite solution for 10 min, rinsing afterwards<sup>[121,131,134]</sup>. (*Level IV, Class IIa*)

---

6.12. What are the materials that you must use to clean the sanitary facilities?

**Answer:** Preferably disposable cloths or, as an option, define different colors for the cleaning of distinct areas, namely the basin and the outer side of the toilet.

The inner side of the toilet must be scrubbed with its own brush and disinfectant-based detergent.

The floor must be washed, preferably, with a product containing, in its composition, detergent and disinfectant since it is easier to apply and has a higher disinfecting power<sup>[131]</sup>. (*Level IV, Class IIa*)

## ***Key Points***

- Critical surfaces must be disinfected, at least, once an hour or after every single use. The areas where children might be must be disinfected several times a day. ***(Level IV, Class IIa)***
- The cleaning of critical surfaces must be accomplished using specific products (EN14476), alcohol solution at 70% or sodium hypochlorite solution-based products at 0,1%. ***(Level IV, Class IIa)***
- The cleaning of the common areas, as well as of the sanitary facilities, must be accomplished using sodium hypochlorite solution at 0,1% (or equivalent), and starting from the least dirty to the dirtiest ones. ***(Level IV, Class IIb)***

## People flow

---

7.1. How can the receptionist keep a social distancing, minimizing the fear that this measure might arouse on the patient?

**Answer:** The receptionist must seek, while keeping customer service cordial, to be as quick as possible, in order to minimize the patient dwell time in the clinic's common areas. Avoid physical contact with the patient and make sure that there is a physical barrier in place keeping receptionist and patient apart - protective acrylic glass (even if partial) with a restricted area (a window) for handing in documents/POS terminal/money. In case there is more than one patient waiting to be served, something which must be avoided, a safety distance of 2 m must be kept between patients. Everyone must wear a mask and avoid physical contact. Request the patient not to touch the counter, door knobs and support areas<sup>[14,116,135]</sup>. *(Level IV, Class I)*

---

7.2. What should you do after finishing serving a patient/accompanying person and before serving the following one?

**Answer:** One must sequentially apply surface disinfectant on the counter in between two patients; perform the antisepsis of the hands using ABAS; make sure that the reception area is aired out at all times<sup>[136]</sup>. *(Level IV, Class IIa)*

---

7.3. Should there be separate entrances for users and professionals?

**Answer:** It has not been established. Where possible, one should choose to make the admission to the practice avoiding that staff and users, as well as waiting users and those on their way out cross paths<sup>[17,103,137]</sup>. *(Level IV, Class IIb)*

---

7.4. What is the minimum mandatory recommendation of personal protective equipment at the dental clinic?

**Answer:** The transmission of SARS-CoV-2 occurs by inhalation-ingestion-direct contact of the mucous membranes with the saliva droplets. Since this virus can survive on different surfaces for several days, the objects and the hands are possible sources of

contamination. The protective equipment inside the clinic must be determined according to the area where the professional works and the activity/procedure the dentist is going to perform, knowing that the basics includes uniform or disposable gown, dedicated shoes or shoe covers and surgical mask<sup>[12,17,92,101,103,110,138-143]</sup>. (*Level IIb, Class I*)

## ***Key Points***

- After a patient has been served, some surface disinfectant must be applied at the reception (on the counter) in between two patients and you must make sure that the reception area is properly aired out. ***(Level IV, Class IIa)***
- The minimum PPE inside the clinic must be determined according to the area where the professional works and the activity/procedure that the dentist is going to perform, the uniform or disposable gown, dedicated shoes or shoe covers and surgical mask being the mandatory minimum inside the clinic for everyone. ***(Level IIb, Class I)***



# Access to the Clinic

## Dentists and Staff

---

### 8.1. Am I able to go to work?

**Answer:** If you have a clinical condition compatible with infection with SARS-CoV-2, you should not go to work<sup>[54,100,119]</sup>. You must contact the SNS (NHS' Portuguese counterpart) 24 support line - 808 24 24 24. *(Level III, Class I)*

---

### 8.2. I have a relative (who is part of my household) who is sick with COVID-19. Can I go to work?

**Answer:** No. If you cohabit or have cohabited with someone infected, you must not report to work during the 14 days following your last contact with the one infected; this situation is considered a close contact with a high risk of exposure<sup>[54,91,100,135,144]</sup>. *(Level III, Class I)*

---

### 8.3. I am pregnant or more than 65 years old, I suffer from related systemic pathologies of higher risk. Can I go to work?

**Answer:** Yes, but you must take into consideration the risks associated to the practice of the profession. It is suggested that the professionals who fit into the categories indicated below must not be assigned to the provision of care involving a higher degree of risk<sup>[119,146]</sup> (presence of chronic medical conditions, including immunosuppression conditions; pregnancy). *(Level III, Class II)*

---

### 8.4. How is it possible to control if the staff are infected?

**Answer:** The measurement and recording of the body temperature must be carried out twice a day. Any other symptom which might be compatible with infection with SARS-CoV-2 must be reported immediately to the clinical director. Should there be a suspicion, the evolution of the case must be monitored by the clinical director<sup>[17,91, 100,119]</sup>. *(Level III, Class IIa)*

## ***Key Points***

- If you have a clinical condition of symptomatology compatible with infection with SARS-CoV-2, you must not go to work. ***(Level III, Class I)***
- If you cohabit (or have cohabited) with someone infected with SARS-CoV-2, you are considered a close contact having a high risk of exposure and should, therefore, remain 14 days in prophylactic isolation, after the last contact. ***(Level III, Class I)***
- It is suggested that the professionals who fit into the categories indicated below must not be assigned to the provision of care involving a higher degree of risk (presence of chronic medical conditions, including immunosuppression conditions; pregnancy). ***(Level III, Class II)***

## Patients and Accompanying People

---

### 9.1. What shall be checked as soon as the patient enters the clinic?

**Answer:** The patient's temperature must be measured, ideally with a contactless thermometer. Next, administer the epidemiologic questionnaire in order to identify the patients at risk of transmitting an infection by SARS-CoV-2<sup>[12,17,54,91,101,110,119,147]</sup>. (*Level IIb, Class I*)

---

### 9.2. The patient is feverish or reports having had fever and cough or sore throat and muscular pains over the last 24h. What should you do?

**Answer:** If it is not an urgent treatment: ask the patient to return home and to contact *Saúde 24* (telephone health advice line). The appointment must be postponed by 2 weeks, always respecting the period of quarantine, and the phone triage must be repeated while confirming the next appointment. In case of emergency, you must consider this patient as a risk patient, who must wait his/her appointment outside or who skips directly to the office, after putting on the appropriate PPE, without remaining in the waiting room<sup>[100,119]</sup>. (*Level III, Class I*)

---

### 9.3. The patient is not feverish. What should you do next?

**Answer:** A surgical mask, a bouffant cap (optional) and shoe covers must be supplied to every patient, whether he/she displays respiratory symptoms or fever or not, upon his/he entrance in the clinic. The way how to put on and remove the mask must be taught/checked. Start by putting the shoe covers on. Before putting the mask on, the patient must be required/taught how to sanitize his/her hands with ABAS<sup>[49,54,107,120,157,147-149]</sup>. (*Level IIb, Class I*)

---

### 9.4. Can the patient come accompanied?

**Answer:** The patient must be advised to come alone, as far as possible, the exception being those patients underage or requiring special needs, such as elderly people,

pregnant women and people displaying visible physical or mental limitations.<sup>[119,135]</sup>. (Level IV, Class IIa)

---

#### 9.5. Should the accompanying person wear a mask too?

**Answer:** Yes, the accompanying person must sanitize his/her hands with ABAS, put a surgical mask and shoe covers on<sup>[17,49,120,124,140]</sup>. He/She will have to be submitted to the same triage and to the same procedures of protection than the patient<sup>[119]</sup>. (Level IIb, Class IIa)

---

#### 9.6. In case the accompanying person walks into the clinical office, where should he/she remain and how should you proceed?

**Answer:** The accompanying person must not be present at the clinical office. In exceptional situations, this person may remain seated 2 m away from the dental equipment, with a surgical mask and shoe covers on<sup>[136]</sup>. (Level IV, Class IIa)

---

#### 9.7. In case the accompanying person does not enter the clinical office, where should he/she wait?

**Answer:** The accompanying person must, ideally, wait outside the clinic or in the waiting room, in case its capacity so allows<sup>[119,136]</sup>. (Level IV, Class IIa)

---

#### 9.8. What should you do with the patient's belongings (and the accompanying person's as well, if he/she happens to walk into the consultation office) before putting on the appropriate PPE?

**Answer:** The patient's belongings must be put inside a clean plastic bag, which must be available at the entrance, and they must be carried around by the patient during his/her stay at the clinic. These belongings will remain inside the bag until the patient leaves the facilities. (Level IV, Class IIb)

## **Key Points**

- Upon arrival to the clinic, some procedures involving the patient and the accompanying person must be carried out (if applicable):
  - ◉ Patients must be told to sanitize their hands, to put a surgical mask and the shoe covers on. **(Level IIb, Class I)**
  - ◉ Temperature check and administration of the COVID-19 questionnaire. **(Level IIb, Class I)**
  - ◉ In case the patient has fever and has presented suspect symptoms over the last 24 h, he/she must return home, if the appointment has not an urgent nature, and contact Saúde 24. **(Level III, Class I)**
- The accompanying person must not walk into the clinical office. **(Level IV, Class IIa)**
- The patient's belongings must be kept in a clean bag made available at the reception. **(Level IV, Class IIa)**

## Maintenance Technicians

---

### 10.1. How should maintenance technicians be welcome?

**Answer:** The technician must avoid direct contact with the clinic's employees and respect the signing delimiting the safety distance. The measurement of body temperature must be performed using a contactless thermometer. The circulation of technicians must be restricted to those areas which their intervention strictly requires. Upon accessing the premises, they must be told to disinfect their hands with ABAS, and put the surgical mask and the shoe covers on<sup>[91,100,116]</sup>. *(Level III, Class I)*

## Delivery men

---

### 11.1. How should delivery men be welcome?

**Answer:** These must not go any further than the clinic's entrance and avoid direct contact with the clinic's employees, as well as respect the signaling delimiting the safety distance. The parcels must be dropped off and remain at the entrance, being stored afterwards by a clinic employee after their disinfection with spray or a wipe containing alcohol at 70%. The employee must sanitize his/her hands with ABAS<sup>[93,116]</sup>. *(Level IV, Class IIa)*

## Waste Collectors

---

### 12.1. How should waste collection employees be welcome?

**Answer:** The responsible for the collection must avoid direct contact with the clinic employees and respect the signaling delimiting the safety distance. He/She must restrict circulation to those areas which his/her intervention strictly requires. The professionals in charge of the transport of hospital waste should wear the appropriate PPE and follow the rules determined by the respective Services of Health and Safety at Work/Occupational Health. Outside the packages/packaging containers for transport of waste, particularly in the lock area/lid and in the handle area, you must apply a disinfectant solution of sodium hypochlorite containing, at least, 1% of chlorine<sup>[150]</sup>. *(Level IIb, Class I)*

## ***Key Points***

- Maintenance technicians:
  - ◉ Avoid direct contact with the clinic staff; respect the safety distances; restriction of circulation to the intervention areas; briefing for the hand hygiene, put a surgical mask and shoe covers on. ***(Level III, Class I)***
- Parcel delivery men:
  - ◉ Do not cross the entrance of the clinic; avoid direct contact with the clinic staff; respect the safety distances; the parcels must be disinfected. ***(Level IV, Class IIa)***
- Waste collection employees:
  - ◉ Avoid direct contact with the clinic staff; respect the safety distances; restriction of circulation to the intervention areas; disinfection of the lock area and container handles ***(Level IIb, Class I)***

# Means of Protection

## Reception/administrative staff

---

### 13.1. What is the appropriate protective equipment?

**Answer:** The reception staff must:

- Wear clothes for exclusive use at the clinic;
- Washable dedicated shoes (for exclusive use at the clinic) or disposable shoe covers;
- Bouffant cap (optional, which must cover the entire hair);
- Eye protection or face shield in case social distancing is not guaranteed at all times;
- Surgical mask, which must be replaced if it is wet or, at most, after 6 h of use<sup>[103,140]</sup>. (*Level IV, Class I*)

Hands must be sanitized before putting the PPE on. In parallel, the eye protection or face shield must be disinfected<sup>[90,116,137,138,151]</sup>. (*Level IIb, Class IIa*)



## Cleaning Staff

---

14.1. What is the appropriate protective equipment?

**Answer:** Clothes for exclusive use at the clinic, including waterproof shoes, bouffant cap, surgical mask, apron and thick disposable gloves<sup>[103]</sup>. (*Level IV, Class I*)

## ***Key Points***

- Receptionists and administrative staff must wear the clinic's own clothing, dedicated shoes or shoe covers, bouffant cap, eye or facial protection and surgical mask. ***(Level IV, Class I)***
- The cleaning staff must wear the clinic's own clothing, dedicated shoes, surgical mask, bouffant cap, apron and thick disposable gloves. ***(Level IV, Class I)***
- Other people coming from outside must wear a surgical mask and shoe covers. ***(Level IV, Class I)***

# The Stages of Peri-Intervention or Peri-Intervention Stages?

## Definition of Risk of the Procedure

---

### 15.1. Should risk areas be identified or marked at the clinic?

**Answer:** The limitation of transmission in a practice/clinic requires a set of preventive and control measures assigned of hierarchically. Setting up physical barriers and specific circuits/flows for patients and professionals allows to minimize contact and exposure. The movement must follow the principle of “three areas and two hallways” - contaminated, potentially contaminated and clean areas - with a buffer area in between the contaminated and the potentially contaminated areas.

There is no need to identify risk areas; however, internally, the definition of areas presenting a risk of differentiated contagion must be present among all the staff members. We can conceptually define the following areas<sup>[17,88,99,121,136]</sup>:*(Level IV, Class IIb)*

- **Green or low risk area:** safe and rest area;
- **Yellow or medium risk area:** area designed for the passage and sterilization of materials;
- **Red or high-risk area:** treatment area.

---

### 15.2. What are the differentiated recommendations for the different risk areas?

**Answer:** The control of exposure to the source of contamination by installing physical barriers or ventilation, and decontamination of the environment allows to reduce the risk of contamination. In different risk areas, only those professionals designated to that veryvery area are allowed, and according to an activity plan and the procedures to be performed, reducing the number of professionals in that area and their entrances/exits. The full and appropriate use of the PPE by the professionals, including protective respiratory equipment, protects physically the professionals themselves and the patients from contagion. The use of breathable materials does not lead to further contamination

and allows for a more comfortable use, which leads to a greater compliance with the usage requirements. The direct instructions, active training and video screening showing the correct donning and doffing are important to improve the effectiveness of the use of the PPE<sup>[17,88,103,135,152-156]</sup>. (*Level III, Class I*)

- **Green area:** basic equipment (uniform or disposable gown, surgical mask and/or social distancing; shoe covers);
- **Yellow area:** degree 1 protection PPE;
- **Red area:** degree 2 and 3 protection PPE.

(read how to put on PPE - [Appendix III, section 2](#))

---

15.3. How should the general procedures in a clinic be ranked in terms of risk?

**Answer:** At the dental clinics there are several levels of equipment to be worn according to the risk to which one is exposed. Considering every patient as a potentially infected individual, the risk of contagion must be considered real and you must not underrate it. The risk is defined by the proximity of exposure as a result of the kind of tasks one performs and by the recurrence of the exposure. We can consider the following levels of risk of the procedure:

- Low risk: sporadic contact with people, considering that social distance is kept and that there is no contact with potentially infected people;
- Moderate risk: constant interaction with people, having, however, the possibility of keeping safety distancing, adopting a respiratory etiquette and wearing masks;
- High risk: frequent and close contact with patients (the distance from them being less than 1 m), but without performing aerosol-generating procedures;
- Very high risk: frequent and close contact with patients (the distance from them being less than 1 m), but with aerosol generation <sup>[88,154,171,157]</sup>.

At the clinic, the administrative and secretariat staff may present a moderate or low risk, however, assistants, hygienists and doctors present a high or very high risk of exposure <sup>[88,154,158,159,171]</sup>. (*Level III, Class I*)

---

#### 15.4. How should the different clinical procedures be ranked in terms of risk?

**Answer:** The risk of the procedure is defined by how closely to the patient it is performed and the need to adopt aerosol-generating measures.

Every procedure involving the generation of aerosols must be ranked as having a very high risk of exposure<sup>[82,85,91,103,119,137,152,154,160-165]</sup>. (*Level III, Class I*)

The clinical procedures may be ranked according to the following levels of exposure:

- Moderate risk - procedures which might be carried out at a minimum distance of 1 m, namely clinical history, extraoral X-rays and extraoral photographs. A PPE which provides for degree 1 protection must be worn.
- High risk - procedures carried out in close proximity with the patient (the distance being less than 1 m), namely intra or perioral procedures. A degree 2 protection PPE must be worn.
- Very high risk - procedures involving the generation of aerosols. A degree 3 protection PPE must be worn.

## **Key Points**

- We can define 3 areas: green area or safe and rest area (low risk); one yellow area or area designed for the passage and material sterilization (intermediate risk); one red area or treatment area (high risk). **(Level IV, Class IIb)**
- Green area/low risk: uniform or disposable gown, surgical mask and/or social distancing and shoe covers. **(Level III, Class I)**
- Yellow area/moderate risk: surgical mask (type IIR), waterproof disposable gown, bouffant cap, dedicated shoes; the use of gloves and goggles (mandatory in procedures of material disinfection and sterilization) may be considered. **(Level III, Class I)**
- Red area/high or very high risk: FFP2 or FFP3 respirator, waterproof disposable gown, surgical cap or cap with a neck protection, goggles and/or face shield, 1 or 2 pairs of gloves, waterproof disposable boot covers (alternatively hood-equipped coverall, preferably made of breathable fabric, and shoe covers). **(Level III, Class I)**
- At the clinic, the administrative and secretariat staff may present a low or moderate risk, depending on the facilities; however, assistants, hygienists and doctors present a high or very high risk of exposure. **(Level IV, Class I)**

# Preparing the office and individual protection

## The space and the equipment

---

### 16.1. What is the stability of the virus in the atmosphere?

**Answer:** In laboratory experimental conditions, the virus has a half-life and a viability of approximately: 1 h and 3 h in aerosols; 7 h and 72 h on plastic; 6 h and 48 h on stainless steel; 4 h and 24 h on cardboard; 45 min and 4 h on copper<sup>[93]</sup>. Moreover, the virus may remain viable on clothes for up to 1 day or on the outer lining of surgical masks for up to 7 days. It also presents a varying stability depending on the ambient temperature: up to 14 days at 4 °C, 7 days at 22 °C, 1 day at 37 °C and up to 10 min at 56 °C<sup>[190]</sup>. The evidence in natural contexts shows the virus' prolonged environmental persistence in spaces occupied by symptomatic and asymptomatic patients for up to 17 days, as well as a potential of nosocomial infections<sup>[168-170]</sup>. The SARS-CoV-2 may remain in the air and on some surfaces, like, for instance, clothes, plastic, stainless steel, cardboard and copper, for hours or even days; consequently, the contact with the surfaces and equipment confined to the consultation office must take place while wearing the appropriate PPE, namely respiratory PPE and disposable gloves, knowing that the latter must not come into contact with the face or any other part of the body. (*Level III, Class I*)

---

### 16.2. What is the reach of contaminated aerosols released in normal breathing in a closed environment?

**Answer:** The classical model on the transmission paths of infectious respiratory diseases lies on a predictive model developed in 1930, which served to explain the transmission of tuberculosis and serves as a basis to the recommendations issued by the main international organizations on COVID-19. The current scientific consensus establishes that the transmission via respiratory secretions emitted by an infected individual happens, mostly, through contact and droplets of bigger dimension (>5-10 µm), without significant capacity for propagation within a distance of up to 2 m. Based on the actual knowledge, derived from laboratory experimental simulations, and from data of quantification of aerosols and dynamics of contagion in hospital units, these premises may be simplified and not reflect the reality regarding the potential of transmission of SARS-CoV-2<sup>[87]</sup>. New

models have shown that cough, sneezes or exhalations are not only formed by mucosalivary droplets with semiballistic trajectories of limited range, but mostly by a multiphase turbulent gaseous cloud which imprisons clusters of droplets, including those of smaller dimension (<5-10 µm), with a potential of aerosolization<sup>[170]</sup>. This cloud develops in a wet and warm microenvironment, protecting the droplets from a precocious evaporation, and enables their spread up to 7-8 m<sup>[171]</sup>. Even though there are no studies which directly assess the potential of infectiousness of the aerosols released by patients infected with SARS-CoV-2, the characteristics of the dynamics of formation of the gaseous cloud of exhaled aerosols and their role in the infectious transmission must be considered.

During the inter-human airborne transmission, the cloud of airborne bioaerosols, generated when someone coughs, sneezes or exhales, may reach distances of 7-8 m, which exceeds the considered safe distance (2 m). (*Level III, Class I*)

---

### 16.3. Which distance may reach the aerosols generated during the operative procedures in Dentistry?

**Answer:** Several factors, including humidity, temperature, dimension of the particles, ventilation, instrumental and operatory postures may influence the potential of dissemination and infection of aerosols contaminated by biological agents. Regarding non-surgical dental procedures, the (sonic and ultrasonic) scaling is the one which has the highest incidence of aerosol spreading, followed by high rotation instruments, polishing by sandblasting and air/water syringe. Those aerosol-generating procedures in the conventional environment of a dental office may generate contamination to distances over 1.5 m<sup>[82]</sup>. The most contaminated areas are number 1 and 2 (0.5 and 1 m around the patient's head, respectively), with an emphasis on the patient's chest (main contamination area), doctor's and assistant's face area, the doctor's right arm and the assistant's left arm, lower neck region, the lamp and its handles, the cutting instruments' support area, the rotary cutting instruments themselves, ultrasounds, and air and water syringe, the doctor's and the assistant's support trays (in case they exist), the terminals and the area where surgical aspirators fit and the spittoon area.

Whenever a well-positioned high flow suction system is associated, the contamination beyond a 60 cm radius is residual<sup>[172]</sup>. The peak of the aerosol decreases to baseline



levels within 10 to 30 min with a quick dispersion or deposition of particles after the dental procedures<sup>[173]</sup>. However, the airborne aerosols may potentially reach any area of the office, meaning that there are no “green” areas which might be guaranteed contamination-free. (*Level Ib, Class I*)

---

#### 16.4. How should dental equipment and the surrounding surfaces be protected?

**Answer:** Besides the dentist/hygienist and the assistant, the targets which are the most directly contaminated by the aerosols in the consultation office are: the lamp and its respective handles, the spittoon and adjacent areas, the movable support tray, the doctor’s operating instruments and their support area, the assistant’s support tray (when it is available), the suction terminals and their respective bracket. Besides the external surface, the rotary or ultrasound cutting operating instruments, and the air and water syringe may also be contaminated internally<sup>[54,174-177]</sup>. However, the aerosols generated while working in the mouth may potentially contaminate any exposed place or infrastructure existing in the dental office. (*Level Ib, Class I*)

Therefore, we recommend you to:

- Protect the head support, arms, chair, lamp, handles, the doctor’s and the assistant’s support trays, the spittoon’s adjacent area, the surgical aspirators’ terminal, the cutting operating instruments, and the air and water syringe with plastic or tin foil barriers;
- Use disposable air and water syringe terminals;
- Use removable and autoclavable lamp handles (when they are available);
- Remove the patient’s chair’s arm and hand supports (where possible);
- You should only keep on the workbenches the material and equipment required for the procedure in question;
- Only the cutting operating instruments, preparation and the like strictly necessary to the defined procedure must be mounted in the work equipment;
- Protect the movable equipment that sits on your workbench, like LED-curing light, endodontic motors, scanners, photo cameras and others using plastic barriers or tin foil;
- Protect the radiographic films and the sensors of digital imaging with a plastic casing, being then coated with a fingerstall, aiming to reduce the salivary

stimulation triggered by the nuisance caused by the grooves in the casing, as well as to reduce the probability of its rupture;

- Possess a system of inner air renewal so as not to allow or reduce the permanence and “deposition” of aerosols on the several surfaces of the offices.

---

### 16.5. How should the disinfection of the several components and infrastructures of the consultation office be carried out?

**Answer:** Human coronaviruses are inactivated by processes of disinfection which act by oxidation with alcohol-based products, hydrogen peroxide and sodium hypochlorite, or by processes of sterilization by heat, like autoclaving<sup>[54,176,178]</sup>. (*Level Ib, Class I*)

- Every area, equipment, instrument and work material in the dental office must be protected with adapted mechanical barriers, as far as possible, and disinfected following specific protocols. The movable material which meets the conditions for that must be sterilized by autoclaving;
- Concerning human coronaviruses, the disinfection of the inner surfaces, and of non-disposable or non-autoclavable materials and equipment, can be done using ethanol-based products at 70%, hydrogen peroxide at 0.5-1% or sodium hypochlorite at 0.1-0.5%, with a minimum exposure time of 1 min. Other physical and chemical agents with virucidal potential can also be used (ultraviolet radiation, glutaraldehyde);
- Promote a flow of air renewal (natural and forced) in the office with a minimum renewal rate of 6 times per hour (namely during aerosol-generating interventions);
- Promote a good surgical suction (8 mm nozzle or slightly superior, close to the source of emission of the aerosol and capacity of aspirating 2.8 m<sup>3</sup> of air per minute, at least). The use of adaptations with larger nozzles connected to the surgical suction system and placed at the entrance of the oral cavity requires a compensation (increase) of the suction flow in order to keep the effectiveness;
- Once the procedure has been completed, you should disinfect the surgical suction systems using ammoniacal or enzymatic disinfectants mixed with water.

---

## 16.6. What is the most suitable ventilation system in a clinical office?

**Answer:** The dental office's ambient air may become easily changed in chemical, physical and microbiological parameters harmful to human beings. Besides the microbiological contamination, the dental office's inner air has parameters which are important for the assessment of the health conditions and well-being, such as: temperature, relative humidity, levels of carbon dioxide and of volatile organic compounds. The heating, ventilation and air conditioning (HVAC) systems, however necessary, may turn into an outbreak of dissemination of contaminants of the ambient air. Microbiological agents which make their way into the ventilation systems and are redistributed by them may spread outbreaks of potential infection through an entire building.

The flow of forced ventilation may influence the risk of residual contamination by aerosols in the ambient air. However, besides the flow, the rate of renewal by "clean" air, the filtration systems and the ducts' physical location seem to be very important factors in the effectiveness of the HVAC systems.

Whenever there is a high concentration of aerosols, one single air renewal may remove approximately 63% of the aerosols. After 5 renewals, it may remain only 1% of the aerosols. That is to say, at a renewal rate of 10 to 12 per hour, 20 min will be enough for an almost total decontamination. With about 6 renewals per hour, it will take an hour. However, if the exhaust system has already been switched on before the generation of aerosols starts, their saturation will never be total and even at lower renewal rates, it may prove possible to keep fairly low environmental values<sup>[77-89]</sup>. (*Level Ib, Class I*)

- The dental office must allow for some natural ventilation. Keep a window open where possible and appropriate;
- The natural ventilation must be complemented by a HVAC system which must take some basic assumptions into account:
  - The rate of minimum air renewal must be 6 times as high as the office's total volume per hour, or about 24 m<sup>3</sup>/hour/person. Whenever there is a high emission of potentially contaminated aerosols, the ideal would be 12 total renewals per hour;
  - It seems consensual that, as far as aerosols are concerned, the installation of the internal exhaust ducts must be carried out at a superior level (near the ceiling);
  - Possess suitable and well-positioned filtering systems;

- Take into consideration the location of the external inlet and exhaust ducts so that they are not too close to one another (a minimum of 5 m, according to the Ordinance 353-A, 2013<sup>[197]</sup>);
- The simple combination of mechanical ventilation (with upper exhaust) with an open window may, by itself, generate a beneficial air flow;
- The most effective ventilation method of all is the following one: the insufflation of cold air at a low speed near the floor and extraction near the ceiling, as part of a displacement-like ventilation strategy, it allows for a unidirectional flow and thermal stratification<sup>[197]</sup>;
- You must make sure that the HVAC systems receive proper technical maintenance on a regular basis.

---

#### 16.7. Is the use of additional systems of purification of biological contaminants in the air of the clinical office justified?

**Answer:** As far as the individual protection in environments with a potential microbiological contamination of the inner air is concerned, the measures of mitigation may be implemented at 5 levels: 1 - control of the source of emission (patient); 2 - localized and immediate extraction at the source (suction); 3 - dilution of the pollution load in the internal environment (ventilation systems); 4 - purification of the ambient air; 5 - use of PPE.

There is no robust evidence as to the need of decontaminating the offices' ambient air by forced purification as a way to prevent the transmission of diseases. There are several available systems, starting with ultraviolet radiation devices (fixed and movable), ozone devices, local aspirators with several types of filters and air treatment (HEPA, coal, UVC). The use of UVC (100-280 nm) in the offices may help decontaminate the air since it has a fungicidal, virucidal and bactericidal potential through the destruction of DNA and denaturation of proteins. In spite of the diversity of supply in this regard, there are no studies where these systems are compared as far as their effectiveness and advantage over one another are concerned, in the several conditions (variables) of the dental medicine offices, nor the side effects and hazards inherent in their use. <sup>[178,180,181]</sup>(*Level IV, Class IIb*)

- Due to the absence of benchmark comparative studies duly applied within the scope of dentistry and to the dynamics of market offer at this point, it seems

pertinent to advise some caution when it comes to select and apply at once a system of purification of the ambient air. Until further information is available, you must promote the correct use of HVAC systems, combined with natural ventilation (where possible) and all the hygiene standards outlined for the remaining four levels of intervention.

---

#### 16.8. What must the acclimatization conditions at the clinical office be?

**Answer:** The temperature, relative humidity and air renewal may affect not only human health globally, but also the existence and survival of some chemical, physical and microbiological agents inside the dental office, including the levels of carbon dioxide, of volatile organic compounds and of bacteria, viruses and fungi. At the ambient temperature of 21°C, an increase of relative humidity from 30% to 50% more than doubles the survival time of coronaviruses which might be present in the aerosols. However, globally, a relative humidity ranging between 30 and 60% is accepted as good. An increase of ambient temperature to 30-40 °C may reduce the survival time of some coronaviruses. Inversely, the drop of temperature to values of 4 to 6 °C may increase the survival of the coronaviruses<sup>[[80,81,87,178,180,181]</sup>. (*Level Ib, Class I*)

- You must promote an inner environment ranging between 21-23 °C (Winter) and 23-25 °C (Summer). The distribution of the temperature in the office must be as uniform as possible, without too fast displacements (flows) of air;
- The relative humidity must be kept at between 30 and 50%. Keeping humidity at the most inferior level (around 30%) may prove beneficial in the reduction of some microorganisms' survival, but may also cause some respiratory discomfort.

---

#### 16.9. Is it advisable to use means of microbiological decontamination in the water supply systems and water reservoirs of the equipment?

**Answer:** The water supply systems and water reservoirs of the dental equipment may contain and convey several kinds of microorganisms such as bacteria, fungi, viruses or protozoa. The microorganisms which are present in the water may also contribute to the microbiological load of aerosols and the exposure to them may cause several serious respiratory, eye and dermal pathologies, among others<sup>[54,85,175,182-184]</sup>. (*Level IIa Class I*)

- Installing systems of physical and/or chemical treatment (ultraviolet, *calvenium*, hypochlorite, povidone-iodine, chlorhexidine) in the devices of water supply and water reservoir of the dental equipment may significantly reduce their microbiological load;
- The use of non-return valves and constant positive pressure may also reduce the microbiological load of the water made available at the cutting terminals (turbine, counter-angle and micromotor);
- Activate the irrigating rotary instruments for 30 s in between patients. Due to the prolonged nocturnal stagnation, the cleaning time in the beginning of the workday must be longer (at least 1 min). After longer inactivation periods, like weekends and holidays, the time to purge the water from the instruments must also be longer;
- Check and replace regularly the filters of the supply water.

---

16.10. What is the effectiveness of suction systems in the reduction of the generation of aerosols at the source of contamination?

**Answer:** Depending on their dimensions, positioning and flow of suction, the suction systems may very significantly reduce the amount of aerosols and droplets expelled to the office's environment<sup>[54,82,185,187]</sup>: (*Level IIa, Class I*)

- Whenever aerosol-generating procedures are used, we recommend the permanent use of an effective and well-positioned surgical suction.
- Even in procedures which are not aerosol-generating, the surgical suction may always be used if you are not using a rubber dam.

---

16.11. Does the use of antimicrobials in the irrigation system of the ultrasonic devices reduce the risk of contamination during aerosolization?

**Answer:** The use of antimicrobial irrigating solutions associated to ultrasonic instrumentation must be understood in the actual context concerning two perspectives: effectiveness on the periodontal clinical parameters and effectiveness in the reduction of the viral load of the generated aerosol.

As far as the first one is concerned, a recent systematic review and meta-analysis have determined the effect of the ultrasonic instrumentation combined with antimicrobials on

the clinical parameters of periodontal inflammation<sup>[187]</sup>. The global results have not detected an added clinical benefit with the use of antimicrobials. A sub-analysis of the data has reported a favorable weak effect for the povidone-iodine (concentrations of 0.5-10%), through a gain of clinical insertion of 0.23 mm, while for chlorhexidine and the essential oils, the considered effect was nil. This is why we do not recommend the use of antimicrobial coadjuvants in the irrigation of ultrasonic instruments from a periodontal clinical efficiency standpoint.

As far as the second item is concerned, two RCT have assessed this intervention in the reduction of contaminated aerosols regarding the bacterial load. Both studies have determined a significant reduction with chlorhexidine compared to distilled water, and one of them highlighted, on top of that, a significant reduction with the use of povidone-iodine compared to distilled water and a superiority of chlorhexidine over povidone-iodine<sup>[189]</sup>. Due to the lingering doubts about the ineffectiveness of chlorhexidine in SARS-Cov-2, povidone-iodine might become an alternative in order to reduce the viral load of the cloud of aerosols. However, there is no available evidence about the impact of the antimicrobials, like irrigating solutions, in the ultrasonic instruments on viruses. (*Level IV, Class I*)

- The available evidence is robust enough to not advise the use of antimicrobial solutions in the irrigation system of ultrasound devices from the clinical standpoint of periodontal benefit. However, from the microbiological safety standpoint, there might be some benefit, still to be proved, in the use of povidone-iodine in order to reduce the aerosols' viral load. This procedure should be considered based on a cost-benefit analysis case-by-case.

## **Key Points**

- SARS-CoV-2 may remain in the air and on some surfaces, like, for instance, clothes, plastic, stainless steel, cardboard and copper, for hours or even days; consequently, the contact with the surfaces and equipment confined to the office must be done with the due PPE on, namely disposable gloves, knowing that these must not come into contact with the face or any other part of the body. **(Level III, Class I)**
- The most contaminated areas are number 1 and 2 (0,5 m and 1 m around the patient's head, respectively). **(Level Ib, Class I)**
- You must promote an inner environment ranging between 21-23 °C (Winter) and 23-25 °C (Summer). / An internal atmosphere ranging between 21-23 °C (Winter) and 23-25 °C (Summer) must be promoted. The distribution of the temperature in the office must be as uniform as possible, without too fast displacements (flows) of air. **(Level Ib, Class I)**
- The relative humidity must be kept at between 30 and 50%. Keeping humidity at the most inferior level (around 30%) may prove beneficial in the reduction of some microorganisms' survival, but may also cause some respiratory discomfort. **(Level Ib, Class I)**
- Whenever aerosol-generating procedures are used, we recommend the permanent use of an effective and well-positioned surgical suction. **(Level IIa Class I)**



# Personal Protective Equipment (PPE)

## PPE donning

---

17.1. Must the PPE to be worn be defined by the risk involved in the procedure?

**Answer:** The risk of the procedure to be performed defines the type of PPE to be worn. However, when it comes to select the PPE, you must consider factors other than its function, like its fitting, how comfortable it is to put on and wear, the possibility of decontamination, being disposable and its cost. The professional must select the combination of elements of the PPE that best fit, given that he/she might have to wear them for long periods and, therefore, they must be as comfortable as possible. As a general rule, the type of PPE must be selected according to the following levels of risk of the procedure<sup>[88,91,152-155,160-162,165,190-194]</sup>. (*Level Ib, Class I*)

- Low risk - basic equipment (uniform or disposable gown, dedicated shoes or shoe covers and surgical mask);
- Moderate risk - degree 1 protection PPE;
- High risk - degree 2 protection PPE;
- Very high risk - degree 3 protection PPE.

(read donning of the PPE - [Appendix III, section 2](#))

---

17.2. What is the appropriate PPE according to the levels of protection?

**Answer:** The selection of the equipment is conditioned by the level of risk and the degree of protection inherent in the procedure to be carried out. Additionally, every equipment must be certified for use in the EU. It is necessary to undergo some training for its donning and, mainly, for its removal, since many of the cases of contamination of the health professionals occur during the removal of the contaminated equipment. (*Level III, Class I*) The basic equipment must be worn upon entering the clinic after hand hygiene. It must be kept on at all times and only replaced by a better PPE, according to one of the three degrees of strategic occupational protection to be applied at the dental clinics<sup>[12,91,98,152,154,160-162,165,190,195-197]</sup>: (*Level IIa, Class I*)

- **Degree 1**
  - Surgical mask (type IIR);
  - Disposal waterproof gown;
  - Bouffant cap;
  - Dedicated shoes;
  - Observation gloves (latex or nitrile).
  
- **Degree 2**
  - FFP2 respirator or equivalent;
  - Disposal waterproof gown;
  - Surgical cap (bouffant caps with a neck protection are optional);
  - Eye protection (goggles with lateral protection and/or face shield);
  - Disposable gloves, preferably with a long sleeve;
  - Disposable boot covers.
  
- **Degree 3**
  - FFP2 or FFP3 respirator;
  - Disposable waterproof gown or coverall;
  - Surgical cap and cap with a neck protection (alternatively, coverall);
  - Eye protection with goggles and face shield;
  - Two sets of disposable gloves;
  - Disposable waterproof boot covers.

---

17.3. Which type of masks are there? And what is the indication for its clinical use?

**Answer:** Face masks are medical devices destined to minimize the transmission of infectious agents between professionals and patients. These masks must comply with EN 14683:2019. They are classified in different types (Type I, Type II and Type IIR) according to their capacity of bacterial filtration and resistance to fluids and splatters. Type IIR mask displays a bacterial filtration effectiveness >98% and resistance to fluids >160 KPa, being called surgical mask, while the Anglo-Saxons call it *Fluid Resistant Surgical Mask* (FRSM). Self-filtering masks or respirators are destined to personal protection against one or more risks. They are encompassed by the PPE and comply with EN 149:2001+A1:2019, for <sup>85</sup>

standards of effectiveness of filtration and total leak into the interior of the mask, and are classified as: FFP1 (*Filter Face Piece 1*) of low effectiveness, leak into the interior 22% and infiltration into the material 20%; FFP2 (N95 of US – NIOSH certification) of medium effectiveness, leak into the interior 8% and infiltration into the material 6%; FFP3 (N99 of US – NIOSH certification) of high effectiveness, leak into the interior 2% and infiltration into the material 1%. Surgical masks must be worn essentially to protect third parties, whereas the valveless respirators allow a bidirectional protection and these must be the ones to be worn by the health professionals. In a context of epidemiologic prevention, the use of respirators equipped with valves is not recommended<sup>[222]</sup>. The respirators are individual protection devices and are more recommended for use by the health professionals to provide a higher protection in contexts with a higher infective load<sup>[[88,91,122,123,137,152,154,155,177,178,181,219]</sup>. (*Level III, Class II*)

---

#### 17.4. What should you do before putting the PPE on?

**Answer:** Before putting a PPE on<sup>[92,93]</sup>: (*Level IV, Class II*)

1. Tie your hair;
2. Remove watch and jewelry to avoid contamination;
3. Drink water before wearing a protective equipment to avoid dehydration;
4. We suggest that you go to the bathroom in order to avoid interruptions;
5. Men must avoid to wear beard, since it affects the mask's marginal sealing;
6. Perform hand washing and hand disinfection.

---

#### 17.5. How are you supposed to put the PPE on?

**Answer:** There are different ways of putting the PPE on, but every single one of them aiming to prevent the secondary transmission of diseases. The process of donning must be carried out near the area where it is supposed to be worn, it improves the safety of the whole process when it is carried out with the active assistance of a team member. Face masks, surgical masks or respirators must be put on and adjusted with your hands properly washed and after putting on the surgical cap. Every mask must provide for the best sealing possible. For that, it must be well tight and adjusted over the nasal bridge and facial contour in order to prevent the entrance of contaminated air through the side of

the mask or respirators. An adaptation test must be conducted at the end. You must conduct a qualitative assessment of the adaptation by softly inhaling in order to detect any strong odor. On the other hand, you must check the sealing, first by producing negative pressure, inhaling to create vacuum inside, which is supposed to cause a slight flattening of the respirator's surface if the sealing is effective; immediately thereafter you must conduct the test of positive pressure, blowing moderately to cause the bulging of the respirator's surface in case of sealing<sup>[152,154,160,162]</sup>.

As you enter the clinic, you must put on the first protective "layer", starting with the removal of the ornaments and the hand washing. Next, you put the bouffant cap and the mask on. Then you must assess the procedure to be performed and put the appropriate PPE on: (Level IV, Class I)

- Wash your hands;
- Adapt the respirator and make the adjustment test;
- Put the disposable waterproof gown or the waterproof coverall on;
- Put the foot protection (shoe or boot covers) on;
- Disinfect the hands using ABAS;
- Put on gloves – make sure they are well fitted-over the gown/coverall;
- Put the goggles on, where required;
- Put a surgical cap/bouffant cap or hood of the coverall on;
- Put a face shield protection on, where required;
- Disinfect the inner gloves using ABAS and put the second pair of gloves on, where required degree 3 protection;
- Check the adaptation of every element. (Level IIa, Class I)

NOTE: The elements of the PPE that you need depend on the level of risk and the adequate degree of protection. Donning a full PPE obeys to a systemization, the dentist being ideally helped by an assistant having previously washed his/her own hands; in order to create surgical conditions of sterilization the assistant must perform the hand disinfection in between two steps<sup>[91,152,155,159,160,162,196,199]</sup>.

---

## 17.6. What care should be taken over the use of PPE?

**Answer:** You should never touch your face to adjust the mask, the respirator or the goggles without having previously made a careful disinfection of the inner gloves.

The use of a second mask or face shield, in addition to the goggles, is advised since it helps keeping the FFP2 mask and the goggles dry, increasing, however, the respiratory and visual discomfort.

The use of a second pair of gloves reduces the tactile sensibility and makes it harder to perform the procedures; however, it reduces the risk of contamination while removing the degree 3 protection PPE<sup>[201]</sup>. (*Level III, Class I*)

---

17.7. What are you supposed to do if the first pair of gloves (inner pair in degree 3 protection PPE) touches a potentially contaminated area?

**Answer:** You must disinfect the gloves using ABAS<sup>[92,106,117]</sup>. (*Level IV, Class I*)

## ***Key Points***

- The selection of the PPE must be made according to the risk of exposure and the degree of protection necessary to the procedure to be performed. ***(Level IIb, Class I)***
- A very careful and ordained donning and doffing of the PPE is required in order to guarantee its protective effectiveness and to avoid that it turns into a focus of cross contamination, respectively. ***(Level IIa, Class I)***
- Surgical masks must be worn essentially to protect third parties. ***(Level III, Class I)***
- Valveless respirators allow a bidirectional protection and these must be the ones to be worn by the health professionals. ***(Level III, Class I)***
- In a context of epidemiologic prevention, the use of respirators equipped with valves is not recommended. ***(Level III, Class I)***

## Treatment

---

### 18.1. How can you reduce microbiological burden in intraoperative stage?

**Answer:** In the current pandemic context, it has been recommended the implementation of mouthrinses with antiseptic solutions prior to the execution of intraoral surgical procedures, in an attempt to reduce the viral load of SARS-CoV-2. The American Dental Association (ADA) and the US Center for Disease Control and Prevention (CDC) have issued recommendations in this regard, despite the low level of evidence, advising the use of oral antimicrobials through mouthrinses with hydrogen peroxide (PXH), povidone-iodine (IPVD), chlorhexidine gluconate (CHX), essential oils or cetylpyridinium chloride (CPC).

To date, there has been no clinical study to support the virucidal effects of any oral solution as a preoperative protocol. The evidence was based only on *in silico* and *in vitro* studies, which established the principle of proof of the rationale of action of some of these antimicrobials, through the disruption of various metabolic pathways, oxidation or rupture of the SARS-CoV-2 envelope.

More recently, several literature reviews have emerged<sup>[2001-204]</sup> that reported the first clinical studies that define the validation of these agents against SARS-CoV-2. An uncontrolled prospective study detected only a transient unsustainable decrease until 2 h in the viral load after mouthrinses with CHX (0.12%) in the saliva of COVID patients<sup>[205]</sup>. A series of 4 uncontrolled cases detected an effective reduction in the 2 patients with the highest viral load up to 3 hours after mouthwash with IPVD (1%, 1min)<sup>[117]</sup>. Another prospective, uncontrolled study with 10 cases, reported that mouthrinses with PXH (1%, 30 sec.) did not reduce the intraoral viral load<sup>[207]</sup>. Finally, a randomized controlled trial with 16 patients evaluated the influence of IPVD (0.5%), CHX (0.2%) and CPC (0.075%). A decrease in viral load was detected in a sustainable way until 6 h, after CPC and IPVD mouthrinses<sup>[208]</sup>.

These studies have significant limitations, derived from the applied methodology, sample size or validation biases, compromising reliable extrapolations. However, in the absence of more robust evidence, the following can be advised in the light of the most current data:

- **the recommendation hydrogen peroxide mouthrinses is questionable and should not continue to be supported, until further evidence is available.** (*Level III, Class IIb*)

- rinse with solutions of povidone-iodine (0.5%) or cetylpyridinium chloride (0.075%) for 30 seconds, before any oral procedure that generates aerosols and/or droplets. (*Level Ib, Class IIa*)

- pre-operative mouthrinses with antimicrobial solutions should be an integral part of the microbial load control protocol in the dental office, namely, in a rational articulation with surgical aspiration and effective ventilation systems to minimize the infectiousness of aerosolization.

---

18.2. Should a perioral decontamination be performed? How should it be performed?

**Answer:** On the patient's face, nose surrounding areas (including the upper lip area) and the inner corner of the eyeball concentrate a great variety and amount of microbiological agents. Contact with oral, nasal and ocular areas is a route of contamination and transmission of viruses and bacteria<sup>[54,209-211]</sup>. (*Level Ib, Class I*)

- Performing the decontamination of perioral and perinasal areas with ABAS, hydrogen peroxide, povidone-iodine or a solution of chlorhexidine and alcohol.

---

18.3. What is the importance of using a rubber dam?

**Answer:** Despite some controversy related to the fact that using a rubber dam may not reduce the aerosol amount, it certainly reduces the production of bio-aerosols and droplets contaminated with blood, saliva and several microbiological agents during the handling of rotary cutting instruments, ultrasonic instruments and saliva and air/water syringe. If the water and air which feed cutting operating instruments are not contaminated with micro-organisms, tooth surfaces to be treated remains as the only potential contamination route of the aerosol which is generated. Wearing a rubber dam reduces the patient's oral breathing, and privileges nasal breathing<sup>[54,82,85,212-214]</sup>. (*Level Ib, Class I*)

- If it is possible and suitable, use a rubber dam in every procedure;



- Cleaning and decontaminating the teeth which are exposed after placing the rubber dam with sodium hypochlorite, povidone-iodine or hydrogen peroxide;
- Special care removing the rubber dam at the end of the consultation, once due to its elastic properties, contaminated droplets which have adhered to its internal surface which was in contact with the mucous membranes, fluids and teeth may be projected. Using surgical suction during removal prevents aerosol generation and it may also prevent the patient from getting choked with his/her own saliva.

---

#### 18.4. Should priority be given to the use of single-dose materials?

**Answer:** All the consumable materials should be dispensed as strictly necessary, in order to minimize the exposure to aerosols. Caps should always be placed when products are not being used. Wherever possible, packagings should be protected by a physical barrier (e.g.: plastic wrap). There should be a circumscribed and duly respected place on the working table to place these materials. The materials used with one patient should not go back to their storage place before being disinfected after removing the packaging physical barriers. The disinfection of all the vials and other packaging of reusable consumables used during a consultation should be performed with a compressed soaked in sodium hypochlorite or a disinfectant wipe<sup>[215]</sup>. (*Level IV, Class IIa*)

### **Key Points**

- Patients with acute respiratory conditions should only be treated in extremely urgent situations. (*Level Ib, Class I*)
- Before the consultation, mouthrinsing and gargling with 1-%% hydrogen peroxide or 02-3.0% povidone iodine solution during 60 seconds. (*Level Ib, Class I*)
- Using a rubber dam whenever it is possible and suitable. (*Level Ib, Class I*)
- Consumable materials should have a physical protection to be removed after the end of the consultation, and they should be decontaminated with a suitable disinfectant wipe, alcohol, hydrogen peroxide or a sodium hypochlorite solution after being stored. (*Level IV, Class IIa*)

## Rotary Instruments

---

### 19.1. What is the impact of operating instruments on aerosol generation?

**Answer:** Microbiological contamination derived from aerosols can provide from the patient (oral cavity and respiratory track), the water, the air, the equipment, or even from the instruments and work material. Rotary instruments, ultrasonic devices, air and particle jets and the air/water syringe can generate a great amount of droplets and aerosols containing micro-organisms and fluids, such as saliva, nasopharyngeal secretions or blood. High-flow surgical aspirators can, when correctly applied, significantly reduce the amount of droplets and aerosols released to the office's internal environment. Using ultrasound and rotary cutting systems without water can also generate contaminated aerosols, even though they are not easily visible to the naked eye. Aerosols and aerosolized droplet nuclei can remain in the air for 30 minutes or more after the end of the procedures.

Procedures using ultrasonic or piezoelectric devices are the ones which generate a greater aerosol amount.

Procedures using compressed air propulsion instruments, namely dental preparation with a turbine, but also with a counter-angle, polishing with particle jet, air/water syringe or dental preparation with abrasion, also generate great amount of aerosols.

Thus, we recommend to<sup>[54,82,85,154,175,216]</sup> (*Level Ib, Class I*)

- Use the turbine as little as possible, replacing it with multiplying or 1:1 counter-angles (with adjusted water and air flow);
- Using a rubber dam (whenever it is possible and suitable) and high-flow surgical aspirators at their maximum exhaustion capacity (8 mm nozzle or above and minimum suction capacity of 2.8 m<sup>3</sup> air/minute,
- Practice, whenever possible, handed work;
- If applicable, we recommend that caries lesions removal should be performed using mechanical or chemo-mechanical techniques;
- Reduce active microbiological load in patients' oral cavity by means of suitable elixir before operative procedure (and also during, if applicable);
- Clean, disinfect, lubricate and sterilize cutting and cleaning equipment (turbine, counter-angles, hand-piece, scalers, air and particle jets and similar jets);
- Efficient air renewal in the office;

- Physical and/or chemical decontamination treatment of the water used in dental equipment, in order to reduce the microbiological load which will subsequently integrate bioaerosols;
- Use effective PPE regarding the risk of inhalation and the contact of aerosols and droplets with skin and mucous membranes;
- Inside the office, even after finishing the consultation, professional must keep their mask and goggles and they should only touch the surfaces which were protected and/or were already disinfected.

---

## 19.2. Which specifications should cutting/preparation operating instruments have? Are non-return systems better?

**Answer:** When using rotary cutting operating equipment or ultrasonic surgical equipment, microbiological agents can penetrate into the tubing and cause an inverse contamination of the equipment's air and water systems. (*Level Ib, Class I*)

- After a consultation and before starting the next one, the residues inside the tubes of cutting instruments and similar should be eliminated, running them during 20 to 30 seconds, in order to purge the air/water feeding system.
- We also recommend the use of operative equipment with non-return valves and constant positive pressure in order to reduce the microbiological load of the water available in the terminals and the risk of contamination<sup>[54,85,175,183,184,217]</sup>.

## ***Key Points***

- Use the turbine as little as possible, replacing it with multiplier or 1:1 counter-angles (with adjusted water and air flow). ***(Level III, Class IIa)***
- Physical and/or chemical decontamination treatment of the water used in dental equipment, in order to reduce the microbiological load which will subsequently integrate bioaerosols. ***(Level III, Class IIb)***
- Use turbines and counter-angles with non-return systems. ***(Level Ib, Class I)***

## After-Intervention Stages

### End of the Consultation and Disinfection of the Medical Office

Disinfectant Solutions in [Appendix III, section 3](#)

---

20.1. I have finished the operative procedure. What comes next?

**Answer:** We recommend you to perform subsequently as follows:

- Lift the patient and instruct him/her to mouthrinse and gargle with the antiseptic;
- Ask the patient to empty his glass and place the napkin in it after;
- Perform the antiseptis of the perioral area using a compress soaked in an alcohol solution or 1% hydrogen peroxide or a povidone-iodine solution (dermal solution at 10%, pay attention to pregnant women, children and iodine intolerant patients);
- Instruct the patient to put on the mask and stay sit until he/she is told to leave the office. Except for the perioral areas and the area strictly necessary to perform the procedures, the dentist should never place his/her hands above his/her waist when he/she is wearing PPE, nor touch oneself or the patient<sup>[140]</sup>. (*Level IV, Class IIa*)

---

20.2. Which are the critical surfaces?

**Answer:** Inside the office, the critical surfaces identified are drawer handles, equipment (tips and air/water syringe), imaging sensors, spittoons, lamps, working table (hanging tray, accessory cart and counter), keyboards, mice, door handles, taps, detergent and disinfectant manual dispensers and consumables packing<sup>[93,118,128]</sup>. (*Level III, Class I*)

---

20.3. How can you minimize contamination and perform the cleaning of critical areas?

**Answer:** In these areas, disposable waterproof barriers, such as plastic wrap, aluminum foil, plastic sleeves or adhesive covers should be used if possible. After the consultation, these barriers should be removed and discarded, and surfaces should be cleaned and disinfected<sup>[91,128,131]</sup>. *(Level III, Class I)*

---

20.4. Should protection barriers in screen surfaces, keyboards and mice be removed between patients?

**Answer:** Yes, all barrier protections should be removed after each consultation, even if they had not been used. In the end of the consultation, surfaces must be disinfected<sup>[177]</sup>. *(Level IV, Class II)*

---

20.5. What care should I take with the casing of the digital imaging sensor and the intraoral x-ray films?

**Answer:** After being used, the protection casing of the digital x-ray sensor, which is disposable, should be discarded in the group III waste container. After being removed from their positioner conventional or phosphor x-ray films must be disinfected (immersed in a solution or cleaned with a wipe) before being placed in the developer or in the reading equipment. In the transition to the developer, films should not be handled with contaminated gloves. Never touch an x-ray film with gloves contaminated with the patient's saliva. If this happens, that film should be disinfected with a compress soaked in 70% alcohol, rubbing both sides of the film for 30 seconds<sup>[178]</sup>. *(Level IV, Class IIa)*

---

20.6. How should X-ray positioners be disinfected?

**Answer:** After the consultation, positioners should be washed with water and detergent, and after that they should be placed in ultrasonic container. Sterilizing procedures should observe the manufacturer's recommendations<sup>[174,177,215,219-221]</sup>. *(Level IV, Class IIa)*

---

20.7. How should digital imaging sensors, intraoral cameras and scanners be protected and disinfected?

**Answer:** Regarding digital imaging sensors, to ensure the prevention of cross contamination, It is recommend the use of a plastic protection sheath combined with a silicone fingerstall (for example, the finger of a glove), which will increase the insulation guaranteed by the commercialized sheaths. Protection sheaths should cover the camera's handle and/or part of the sensor's handle in order to ensure the protection from the contact with saliva. The disinfection of these devices should be performed with virucidal disinfectant wipes and removable parts should be properly sterilized<sup>[76,126,174,177,215]</sup>. (*Level IV, Class IIa*)

---

20.8. Should the purge of rotary instruments be performed before being removed to be sterilized?

**Answer:** After the consultation, rotary instruments should be left running for 20-30 seconds, including the instruments with anti-suck-back ability. The use of rotary instruments with no anti-suck-back ability is discouraged. After removing the physical barrier, they should be disconnected from their coupling and placed on the examination tray<sup>[54,128]</sup>. (*Level IV, Class IIa*)

---

20.9. Is chemical external disinfection of turbines, counter-angles and scalers enough against SARS-CoV2 or should they be sterilized between patients?

**Answer:** No. Cleaning and external disinfection with suitable disinfectants does not reduce the risk of cross infection in instruments which are not equipped with a anti-suck-back system. In addition to the external disinfection of these devices, it is recommended that their sterilization be made between patients<sup>[54,177,222-224]</sup>. (*Level IIIb, Class IIa*)

---

20.10. What is the sequence for physical barriers and rotary instruments removal?

**Answer:** Barrier removal should be performed from the least contaminated area towards the most contaminated area. Regarding physical barriers, firstly you should remove the ones on the handles (lamp, equipment tray, X-ray tubes) and finally the ones on the aspirators, the air/water syringe and rotary instruments; these should be removed to be disinfected in the sterilization area. After the removal of physical barriers and instruments, surfaces must be disinfected with a virucidal wipe<sup>[128]</sup>. (*Level IV, Class IIa*)

---

20.11. How should physical barriers to protect the equipment be eliminated?

**Answer:** Physical barriers must be placed in the group III waste container<sup>[177]</sup>. (*Level IV, Class II*)

---

20.12. Regarding surface disinfection (equipment and workbenches), which product is more efficient and for how long should contact time be? be the contact time?

**Answer:** Surface disinfection can be performed using a disinfectant detergent to achieve a faster procedure, i.e. a product containing compatible detergent and disinfectant (2 in 1). Generally, commercially available products (such as virucidals), duly certified by the relevant authorities should be used. They can have different presentations: liquid, gel, foam or spray. Alternatively, the use of alcohol-based solutions (70%, at least 1 min), hydrogen peroxide (0.5-1% at least 1 min), sodium hypochlorite (0.1-0.5%, at least 10 min) has also been proven as being effective. Metal surfaces which are not compatible with sodium hypochlorite due to the risk of corrosion, should be disinfected with alcohol-based solutions or another suitable product for, at least, 1 min<sup>[65,66,128,131,225-229]</sup>. (*Level III, Class I*)

---

20.13. Is UV radiation an effective sterilization method?

**Answer:** UV radiation is effective when combined with surface disinfectants with virucidal properties<sup>[227,230]</sup>. (*Level Ia, Class IIa*)



---

20.14. Regarding wrapping and packaging of the consumables used, which is the most suitable disinfection product and what is its contact time?

**Answer:** Consumable packaging used during the consultation should be disinfected with a certified virucidal disinfectant or, alternatively, with 0.1% sodium hypochlorite or 70% alcohol, which should be left on the surface for at least 1 min<sup>[112,178]</sup>. (*Level III, Class I*)

---

20.15. Is it necessary to make the suitable aeration of the clinical office before starting the following consultation?

**Answer:** The aeration of the clinical office is recommended by most authors. However, the time necessary to this aeration depends on the room's volume and on the available ventilation features, among other factors, and it is not fully clarified. Some authors refer natural aeration for, at least, 10 minutes or the implementation of an HVAC system<sup>[77-86,88,89,227]</sup>. (*Level IV, Class IIa*)

---

20.16. How often should the clinical office's floor be cleaned and disinfected?

**Answer:** It should be disinfected, at least, 3 times a day, using suitable cleaning agents and, additionally, whenever contamination with saliva, blood or other type of contaminant has noticeably occurred<sup>[91,128,131]</sup>. (*Level III, Class I*)

---

20.17. Regarding the clinical office's floor, which product should be used and for how long should it be left in contact to achieve a suitable disinfection?

**Answer:** Disinfection must be performed using a washing detergent and adding a certified virucidal disinfectant, 0.2% sodium hypochlorite or 0.5% hydrogen peroxide, and it must be left to dry<sup>[108,121,178,189]</sup>. Floor cleaning should always be performed using a wet method, and avoid the use of vacuum cleaners<sup>[108,178]</sup>. (*Level IV, Class I*)

---

## 20.18. How should the clinical office cleaning be performed?

**Answer:** Cleaning should always be performed downwards (from top to bottom) and from the cleanest areas to the dirtiest ones<sup>[128,131]</sup>: *(Level IV, Class I)*

1. Walls;
2. Equipment present in the area;
3. Surfaces above the floor (working benches, tables, chairs, others);
4. Sinks;
5. Floor.

---

## 20.19. Which cleaning/disinfection products should be used?

**Answer:** Cleaning products must contain certified disinfectants to be used in hospital/ clinic settings and comply with EN 14476, about virucidal products. They must be applied according to the manufacturer's instructions.

As an option, for less critical surfaces, sodium-hypochlorite-based (0.1-0.5%) can be used. The solution must be prepared only at the time it will be used (active for 24 h). It is important to leave the product in contact during 10 min and then rinse it<sup>[112,131]</sup>. *(Level IV, Class I)*

---

## 20.20. Which types of presentation should cleaning products have?

**Answer:** Cleaning products can be used as sprays, disposable cleaning cloths or wet wipes in an individual dispenser. Regarding disposable products, a different one should be used for each surface to be cleaned, and it should be left to dry, guaranteeing that the disinfectant is left for the recommended time<sup>[112,131]</sup>. *(Level IV, Class I)*

---

## 20.21. What is the most suitable disinfectant to be used in the form of wipes?

**Answer:** Soaked wipes require a short time to take effect (up to 1 min); the most recommended disinfectant is 62-71% ethanol, with 1 min of exposure<sup>[178]</sup>. *(Level III, Class I)*

---

20.22. How should surfaces be cleaned in the event they are contaminated with blood or other organic products?

**Answer:** Should that happen, the spillage should be completely absorbed with absorbent paper, wearing gloves and goggles; after that, a 5% sodium hypochlorite solution (bleach) or a suitable disinfectant should be applied and left for, at least, 10 min; finally, the dirty area should be washed with water and a common detergent, rinsed with water and left to dry in air<sup>[112,131]</sup>. (*Level IV, Degree I*)

## **Key Points**

- Performing antiseptics of perioral area with 1 % hydrogen peroxide or a 10 % povidone-iodine solution, and leave the clinical office wearing a mask. **(Level IV, Class IIa)**
- Protection barriers placed on clinical contact surfaces should always be removed after the end of the consultation. **(Level IV, Class II)**
- Barriers should be removed from the least potentially contaminated place to the most contaminated one, and subsequently surfaces should be disinfected with an alcohol-based solution. **(Level IV, Class IIa)**
- X-ray sensors and intraoral films protections, as well as physical barriers removed should be placed in a group III waste container. **(Level IV, Class IIa)**
- Positioners should be washed and disinfected according to the manufacturer's recommendations. **(Level IV, Class IIa)**
- Digital imaging sensors, intraoral cameras and scanners require a double protection with a plastic barrier and a silicone fingerstall. **(Level IV, Class IIa)**
- After each usage, rotatory systems should be activated for 20 to 30 s, and their sterilization between consultations is recommended. **(Level IIb, Class IIa)**
- The clinical office surfaces should be disinfected with 2 in 1 solutions (detergent and disinfectant). Regarding metal surfaces, alcohol-based solutions should be chosen. **(Level III, Class I)**
- Consumable packaging should be disinfected with a virucidal product or, alternatively, a 0,1 % sodium hypochlorite or 70% alcohol solution. **(Level III, Class I)**
- The products used to clean the office should have a certified virucidal action. Cleaning should be performed downwards, from the cleanest areas to the dirtiest ones. **(Level IV, Class I)**
- The office's floor should be cleaned, at least, 3 times a day. **(Level III, Class I)**

## Materials and devices to be sent to the prosthetics laboratory

---

### 21.1. What is the protocol to disinfect dental impressions?

**Answer:** Firstly, impressions should be washed with water for 10s and then they should be disinfected with a specific product according to the material involved: elastomers (silicones and polyesters) should remain immersed for, at least 5 min, in ammonia and aldehydes-based disinfectants, 10min in 5% sodium hypochlorite or 10 min in 3% hydrogen peroxide; alginates should be sprayed with a 1% sodium hypochlorite-based disinfectant and placed in a closed plastic bag for 10 min<sup>[231,232]</sup>. (*Level IIIa Class I*)

### 21.2. What is the protocol to disinfect dental stone models?

**Answer:** Regarding dental models handled during the consultation and contaminated with saliva, they should be immersed in a solution of slurry water with 0.5% sodium hypochlorite or 2% glutaraldehyde for 5 min, without the risk of dimensional change<sup>[233-235]</sup>. (*Level IIIb, Class IIa*)

### 21.3. What is the protocol for try-in devices (copings, registration waxes, preliminary crowns/bridges and removable prosthesis) disinfection?

**Answer:** Regarding try-in devices, they can be disinfected with alcohol at 70% for 5 min<sup>[178]</sup>. (*Level IIb, Class I*)

### 21.4. What is the protocol to disinfect pre-existing prosthesis?

**Answer:** Regarding acrylic prosthesis, they should be immersed in an antiseptic solution containing sodium hypochlorite at 1% for 10 min. Prosthesis with metal parts should not be immersed in sodium hypochlorite; alternatively, hydrogen peroxide at 1% should be used during 10 min<sup>[236]</sup>. (*Level IIa, Class IIa*)

## ***Key Points***

- Prints should be washed with running water and subsequently they should be placed into dental impression disinfectant:
  - Elastomers: ammonia and aldehydes-based disinfectants for 5 min; or 5% sodium hypochlorite for 10 min or 3% hydrogen peroxide for 10 min.;
  - Alginates: they should be sprayed with 1% sodium hypochlorite and kept in a closed bag for 10 min. ***(Level IIa Class I)***
- Regarding plaster models handled during the consultation and contaminated with saliva, they should be immersed in a solution of slurry water with 0.5% sodium hypochlorite or 2% glutaraldehyde for 5 min. ***(Level IIIb, Class IIa)***
- Try-in testing devices should be decontaminated in an alcohol-based solution at 70% for 5 min. ***(Level IIb, Class I)***
- Pre-existing acrylic prosthetics should be immersed in a 1% sodium hypochlorite solution for 10 min; if they have metal parts, use 1% hydrogen peroxide instead for 10 min. ***(Level IIa, Class IIa)***

## Removal of PPE

---

22.1. What is the sequence to remove PPE in each one of the levels (1, 2 and 3)?

**Answer:** There are different ways of removing PPE safely and without contaminating clothes, skin, eyes or oral and nasal mucosae with potentially infectious droplets. All the PPE should be removed before getting out of the office, except the respirator, if applicable. The removal sequence can vary and depends on the patient's risk level or on the procedure to be performed. Goggles and facial protection should be removed from the back, by lifting head band or ear pieces. Remove the face shield, goggles and place them in a specific place on the workbench for subsequent disinfection, if they are reusable; otherwise, discard / dispose of remove them directly in a waste container<sup>[54,92,98,106,128,139]</sup>. (*Level IV, Class I*)

### **Level 1:**

- Remove the water-resistant gown from the outside, pulling away from shoulders, and rolling it gradually and gently in order to generate the less volume possible and remove hands simultaneously;
- Hands antisepsis with an ABAS;
- Remove cap and surgical mask and wash your hands again.

### **Level 2:**

- Remove the water-resistant gown from the outside, pulling away from shoulders, as in the previous level;
- Remove disposable gloves together with the gown;
- Hand antisepsis with an ABAS;
- Remove eye protection (face shield and goggles) from the back to the front and surgical cap from front to back;
- Remove shoe covers;
- Hand antisepsis with an ABAS;

- Remove FFP2 respirator or equivalent (from the back to the front, pulling elastic bands, without touching the outer side of the mask);
- Hand antisepsis with an ABAS;

**Level 3:**

- Remove the outer pair of gloves;
- Disinfect inner pair of gloves with ABAS;
- Remove eye protection and surgical cap /hood of the surgical suit (if you are wearing a surgical cap untie it first):
- If you are wearing a face shield: remove the face shield and surgical cap/bouffant cap/hood with a single gesture, from the back to the front, touching only the outside;
- If you are wearing goggles: remove goggle and, if they are reusable, place them in a suitable container; then, remove surgical cap /hood from the back to the front, touching only the outside;
- Remove the gown (unfasten side ties and pull gown away from shoulders touching only the outside, with an abrupt movement) and simultaneously remove inner gloves; or remove surgical suit and then gloves;
- Remove shoe covers;
- Perform hand antisepsis with ABAS;
- Remove respirator FFP2 or FF3 respirator (from the back to the front, pulling elastic bands, without touching the outer side of the mask);
- Hand antisepsis with an ABAS;

Remark: if your hands touch accidentally a contaminated surfaces during removal, immediately disinfect with ABAS.

(read on how to put on PPE - [Appendix III, section 2](#))

---

22.2. Where should disposable PPE be placed after being removed? In what type of container?



**Answer:** Disposable PPE should be discarded directly into a group III waste container<sup>[237]</sup>.  
(Level IV, Class I)

---

### 22.3. What is the protocol to remove gloves?

**Answer:** Except for the perioral areas and the area strictly necessary to perform the procedures, the dentist should never place his/her hands above his/her waist when he/she is wearing PPE, nor touch oneself or the patient, once the gloves outer side is contaminated.

Glove removal protocol<sup>[92,106,139,200,237]</sup>: (Level IV, Class I)

- Using a gloved hand, grasp the palm area of the other gloved hand and peel off the first one;
- Hold removed glove in gloved hand;
- Slide fingers of ungloved hand under remaining glove and peel off second glove over first glove;
- Discard gloves in a contaminated waste container;
- Disinfect remaining gloves with ABAS;
- After removing the first pair of gloves, hands should be immediately washed using a suitable detergent and antiseptic.

---

### 22.4. When and how should the dentist remove the surgical mask?

**Answer:** Once the patient has left the office, the dentist can remove the surgical mask or respirator, according to the following steps<sup>[92,237]</sup>: (Level IV, Class I)

The outside of the mask/respirator is contaminated - do not touch it!

Grasp bottom ties or elastics of the surgical mask/respirator and remove without touching the outer area;

Place the surgical mask/respirator and the gloves in the group III waste container.

---

## 22.5. When should the assistant remove PPE?

**Answer:** PPE should only be removed when the office has been completely decontaminated<sup>[137]</sup>. In routine check-ups, after disinfection and removal of all the disposable means of protection and barriers, the assistant should guarantee the office's aeration, and then put on a new pair of gloves to place again all the means of protection and physical barriers in the office, devices and other consumables, to be ready for the following consultation. *(Level IV, Class I)*

---

## 22.6. Can FFP2 (N95) and FFP3 respirators be reused?

**Answer:** FFP2 and FFP3 respirators can be used for 5 to 6 hours. When used with different patients, they should be covered with a surgical mask, and the latest should be exchanged in the end of each procedure. FFP2 or FFP3 respirators masks should be exchanged every time they get humid or visibly contaminated. It is important to avoid touching the mask's external surface after putting it on<sup>[238,239]</sup>. *(Level IV, Class II)*

---

## 22.7. In which place should PPE be removed? In the clinical office or somewhere else?

**Answer:** PPE should be removed before leaving the clinical office, except for the respirator. It is important that you never leave the office or touch surfaces (handlers, tables, support chairs, working benches, drawers, walls, all the surrounding material) with outer or inner pair of gloves which you are wearing<sup>[237]</sup>. *(Level IV, Class I)*

---

## 22.8. Where should the assistant place PPE after being removed?

**Answer:** The reusable protection suit should be placed in a suitable place inside a closed bag for later processing. Disposable contaminated material should be placed in group III waste containers. Face shields and goggles should be placed in a suitable place for subsequent disinfection<sup>[137]</sup>. *(Level IV, Class IIa)*

---

22.9. How should the face shield and the goggles be washed and disinfected?

**Answer:** In this context, we recommend the following sequence:

- (1) Washing the inner surface and then the external surface with a detergent;
- (2) Disinfecting the external surface with a suitable disinfectant;
- (3) Washing it with running water or spraying with alcohol;
- (4) Drying it with absorbent paper or leave it exposed in the air<sup>[240]</sup>.

Remark: this procedure should be performed wearing gloves. (Level IV, Class I)

---

22.10. Can uniforms be worn again the following day without being washed?

**Answer:** Uniforms should not be worn again the following day without being washed<sup>[241]</sup>.  
(Level IV, Class IIa)

---

22.11. What kind of treatment should be given to non-critical fabric materials: uniforms?

**Answer:** Even though this kind of clothing represents a very reduced risk of SARS-CoV-2 infection and spread, during this pandemic stage it should be washed more often and handled with gloves, or you should always wash your hands after handling them. Wash uniforms according to the manufacturer's instructions. Select the hottest water possible program, and dry them completely in a dryer machine and/or iron them<sup>[241]</sup>. (Level IV, Class IIa)

*At home:*

---

22.12. After a day at work how should you behave when you get home?

**Answer:** Shoes should be left at the door, take your clothes off and have a shower immediately<sup>[111]</sup>. (Level IV, Class IIb)

## ***Key Points***

- The removal level 1, 2 and 3 PPE should follow a sequential order. ***(Level IV, Class I)***
- After being removed, the PPE should be placed in the group III waste residues. ***(Level IV, Class I)***
- While you are wearing the PPE, your hands should not touch below your waist, and gloves should be removed according to a particular sequence. ***(Level IV, Class I)***
- The dentist can remove the surgical mask after getting out the office; the assistant can remove the surgical mask after decontaminating the office. ***(Level IV, Class I)***
- FFP2 and FFP3 respirators may be kept for up to 6 hours, once they are protected by a surgical mask exchanged between patients. ***(Level IV, Class II)***
- Face shield and goggles should be washed and disinfected, firstly in the inner side and then the outer side, with water, soap and ABAS. This procedure should be performed wearing gloves. ***(Level IV, Class I)***
- Uniforms should be washed at the end of each day. (Level IV, Class IIa)After a working day, shoes should be taken off and be left at the door, take your clothes off and have a shower immediately. ***(Level IV, Class IIb)***

## Patient care after treatment

---

23.1. Does the use of a surgical mask by the patient inside the clinic reduce the risk of cross contamination with SARS-CoV-2?

**Answer:** As soon as the procedure is finished and the patient has mouth rinsed, the individual protection mask should immediately be put on<sup>[116,140]</sup>. (*Level Ib, Class I*)

---

23.2. Should patients' movement inside the clinical office be limited after the operative procedures?

**Answer:** The patient should remain seated while registration procedures are performed and the next appointment is made<sup>[17]</sup>. (*Level IV, Class IIa*)

---

23.3. Should additional information be provided to the patient in digital form instead of paper form?

**Answer:** You should ensure that a message is sent to the patient informing about the date and time of the next appointment; make also sure a notification will be sent the day after the consultation. Regarding medical prescriptions, preference should be given to electronic media (e.g. text messages, e-mail)<sup>[17]</sup>. (*Level IV, Class IIa*)

---

23.4. How should hand antisepsis be performed before leaving the office, regarding the dentist, the assistant, the patient/accompanying person?

**Answer:** In the clinical office, there should be an ABAS dispenser to perform hand antisepsis before leaving the office. This action should be repeated at the reception, in the event a surface is touched or if sanitary facilities are used<sup>[17, 116]</sup>. (*Level IV, Class IIa*)

## ***Key Points***

- The patient must remain seated until the next appointment is well-marked, being informed of this by message. ***(Level IV, Class IIa)***
- The patient must leave the consultation with the mask provided at the entrance. ***(Level Ib, Class I)***
- The patient must perform hand antisepsis with ABAS before leaving the office and doing it again at the reception, if necessary. ***(Level IV, Class IIa)***

## Patient's return to the Reception

---

24.1. Should the patient keep social distance at the reception and means of protection after consultation?

**Answer:** Safety distance should be kept regarding the counter and according to the signaling and physical contact with surfaces and with the receptionist should be avoided<sup>[54]</sup>. (*Level IV, Class IIa*)

---

24.2. Should the patient keep the mask on in the reception area after the consultation?

**Answer:** The patient should always be wearing a protection mask in all the clinic's environments<sup>[140]</sup>. (*Level IV, Class I*)

---

24.3. Which important additional information should be provided to the patient by the receptionist, in the context of COVID-19 infection, in the post-consultation stage?

**Answer:** The receptionist should:

Inform the patient that he/she should contact the clinic/office within 15 days after the consultation if he/she develops symptoms compatible with COVID-19;

Inform the patient in the event any medical staff or non-medical staff (assistants/receptionist) develop symptoms compatible with COVID-19 within 15 days after the consultation;<sup>[112]</sup>. (*Level IV, Class IIa*)

---

24.4. Which information should be confirmed to the patient by the receptionist before leaving the clinic?

**Answer:** The receptionist should confirm the following information with the patient: consultation registration, validating if the message scheduling the following appointment has been received, and if applicable, reception of medical prescription. Providing antiseptic gel or solution, so that the patient can sanitize his/her hands before leaving;

remind the patient to remove the mask when and only he/she arrives home or, at least, outside the clinic premises; say goodbye to the patient warmly and positively<sup>[136]</sup>. (*Level IV, Class IIa*)



## **Key Points**

- Guaranteeing the observance of social distance at the reception. **(Level IV, Class IIa)**
- The patient should always wear a mask at the clinic. **(Level IV, Class I)**
- In the event the patient develops COVID-19 symptoms within 15 days after the consultation, he/she should contact the clinic; similarly, if any professional at the clinic develops any symptoms, the patient should be informed. **(Level IV, Class IIa)**
- Before the patient leaves the clinic, clinical registration should be confirmed, as well as the appointment of the following consultation and the validation of the confirmation of the following appointment. **(Level IV, Class IIa)**
- Before leaving the clinic, the patient should sanitize his/her hands and he/she should also be reminded not to take off the mask until he/she gets home. **(Level IV, Class IIa)**

## Payment methods

---

25.1. Which payment methods and receipt issuing methods should be recommended?

**Answer:** Payment and receipt issuing should, preferably, be made using electronic means (dematerialized paperless), namely using contactless cards or *MBway*, *iPAY*, among others. In the event that it is not possible, the patient should swipe his/her own card for strip or magnetic chip reading. In this case, the patient should enter the code only pressing the keys and without holding the POS terminal; keep the payment terminal machine duly protected with plastic wrap and place another film over the keys, whenever a patient needs to enter a personal code. Once the patient has used and entered the code, the plastic wrap should immediately be removed or a 70% alcohol disinfectant should be applied<sup>[242]</sup>. Using a payment card makes cleaning and disinfection easier comparatively with paper money, once there is not a suitable disinfection method for this kind of money, despite the fact that the risk of infection due to the handling of this kind of money seems to be very reduced in the light of the existing evidence. (*Level IV, Class IIa*)

## ***Key Points***

- Payments made preferably using electronic means - contactless, MBway, iPAY and others. If that is not possible, the patient should handle his/her own card. ***(Level IV, Class IIa)***
- After using the POS terminal, the protection should be replaced or a 70% alcohol solution should be applied. ***(Level IV, Class IIa)***

## Sterilizing

### Material Cleaning, Disinfection and Sterilization

#### Standards

---

26.1. Does contamination with SARS-CoV-2 imply a change of the procedures used to clean, decontaminate and sterilize materials and instruments?

**Answer:** It does not imply a change of good practice, however the recommended procedures should be reminded. Non-rotary instruments should be pre-processed (decontaminated) with a suitable disinfectant or with a 5% sodium hypochlorite solution (pay attention to metals), for 30 minutes; washed with water, placed in the ultrasonic container with an enzymatic disinfectant for the recommended time and at the recommended temperature; after getting dry, they should be placed in a sterilization sleeve (in kits or individually) and placed in the autoclave. Rotary instruments should be submitted to a pre-treatment with a 70% alcohol solution, a detergent washing and, after getting dry, they should be lubricated, placed in the sterilization sleeve and in the autoclave. After the autoclave and before being used again, rotary instruments can be lubricated again, manually or in dedicated equipment. Instruments which get in contact with the patient's oral mucous membranes and fluids (e.g., dental mirror, tweezers, restorative material applicators) should be considered as semi-critical and should be sterilized using thermal methods (steam and pressure - autoclave; chemical vapor - formaldehyde; dry heat); semi-critical instruments sensitive to heat should not be reused, they should be discarded instead. Surgical instruments and other instruments which penetrate soft tissues or bones (forceps, periodontal cures, bone drills, etc.) should be considered as critical and, if they are reusable, they should be sterilized in the autoclave<sup>[86,143,174,177,215,243]</sup>. (*Level IIIb, Class IIa*)

---

26.2. Which kind of bag is indicated to pack group III waste?

**Answer:** Waste resulting from each consultation should be packed in a suitable bag sealed with ties, sprayed with sodium hypochlorite and placed in a storing container to be collected by the certified company<sup>[54]</sup>. (*Level IV, Class I*)

---

26.3. Should the placement of group III waste in the corresponding container be performed in the clinical office or in the sterilizing room?

**Answer:** It should be performed as close as possible to the place where the waste is produced, i.e. inside the office<sup>[54]</sup>. (*Level IV, Class I*)

---

26.4. Does contamination with SARS-CoV-2 imply a change of the procedures used to handle and process waste?

**Answer:** This virus contamination degree implies protocolized and multiplied care to minimize the risk of cross infection<sup>[137]</sup>. (*Level IV, Class IIa*)

---

26.5. Should the material/instruments handled by the dentist be considered as semi-critical or critical?

**Answer:** All the material/instruments handled by the dentist should be considered as critical. It should be sterilized, as long as its characteristics allow it. For biosafety purposes, non-critical material should be processed as semi-critical material when it is handled by the dentist<sup>[244]</sup>. (*Level IV, Class I*)

Basic principles to follow:

Whatever can be sterilized should not be disinfected;

Always give preference to physical sterilizing methods, namely by heat (autoclave).

---

26.6. Should the material/instruments handled by the dental assistant be considered as semi-critical or critical?

**Answer:** All the material/instruments handled by the dental assistant and which came into contact with the patient's mucous membranes, saliva or other fluids should be considered as critical. If it is not disposable, it should be sterilized, as long as its characteristics allow it. When that is not possible, it must be disinfected<sup>[244]</sup>. (*Level IV, Class I*)

---

26.7. Is it possible to disinfect FFP2 (N95) and FFP3 respirators?

**Answer:** FFP2 respirators can be disinfected with hydrogen peroxide vapors and UV light; however, these procedures are only justified in times of lack of equipment, once there is not enough evidence in literature to validate these procedures<sup>[245]</sup>. (*Level IIb, Class IIb*)

## ***Key Points***

- All the material handled by the doctor and the assistant is considered as critical and should be sterilized in autoclave, whenever manufacturer's instructions allow it. ***(Level IV, Class I)***
- Non-rotary instruments should be decontaminated (Level IIb, Class IIb) with a suitable solution or with a 5% sodium hypochlorite solution, washed with water, placed in the ultrasonic container (enzymatic detergent), dried and placed in a sterilization sleeve and autoclave, according to manufacturer's instructions. ***(Level IIb, Class IIa)***
- Rotary instruments: 70% alcohol solution, detergent washing, dried, placed in sterilization sleeve and autoclave. ***(Level IV, Class IIa)***
- Material which gets in contact with the oral cavity: we recommend sterilization in the autoclave. ***(Level IV, Class IIa)***
- Group III waste should be placed in a suitable bag, tied and sprayed with hypochlorite. Its disposal in the container should be performed in the clinical office. ***(Level IV, Class I)***

## Accidental Exposure

### How to act in the event of Accidental Exposure

---

#### 27.1. Is the oral cavity a tropism location for SARS-CoV-2?

**Answer:** There is an urgent need for information about the specific local tropism of this virus in the human body, to allow us to understand its replication, immune response and infectiousness. Most coronaviruses bind to cell receptors through a spike protein. SARS-CoV-2 uses ACE2, expressed in different organs, as a key receptor<sup>[146]</sup>. The affinity for this receptor seems to be greater with this virus than with other coronaviruses<sup>[176]</sup>. The predominant expression of ACE2 in the lower respiratory tract is believed to have determined the natural history of SARS-CoV-2 as a respiratory infection of the lower tract. However, there is evidence of tropism for oropharyngeal cells, with an active replication during the first 5 days after the onset of symptoms and with a high viral load in saliva in the posterior oropharynx<sup>[176, 177]</sup>. In a preclinical model with non-human primates, salivary gland ducts were the early target cells of SARS-CoV and, therefore, a source of viral load in saliva<sup>[178]</sup>. Posterior oropharynx is a meeting point for secretions from the posterior nasopharynx, the tracheobronchial tree and salivary glands. Preliminary data show that there is a high expression of ACE2 receptors in the epithelial cells of oral mucosa, which was higher in tongue, indicating that the oral cavity may be a potentially higher risk route of SARS-CoV-2 infection<sup>[250]</sup>.

Several places in the oral mucosa were identified, expressing the main key receptor which establishes the connection to SARS-CoV-2 and mediates intracellular invasion. Even though there is no evidence of the direct action of this tropism as a preferential niche for viral colonization, preliminary data establish rational proof of this mechanism, which indicates that the oral cavity may be a highly susceptible place for SARS-CoV-2 infection.

*(Level IV, Class I)*

---

#### 27.2. In the event of an accidental exposure to biological contaminants from a COVID-19 patient, which procedure should be adopted?



**Answer:** Regarding the hands (without protection or with gloves which have lost their protection barrier, ex accidental perforation), you should perform their immediate antiseptics with ABAS. The skin is the most external barrier of the human body. Health professionals working with COVID-19 patients are prone to damages of their skin and mucosa barriers. To minimize the break of skin and mucosa barriers, health professionals should observe the standards related to the use of PPE and avoid an excessive exposure. Regarding mucous membranes, namely the ocular conjunctiva, an immediate washing with water or a saline solution should be performed. In the case of a suspected exposure when removing the goggles, the periocular area should immediately be cleaned with a cotton ball soaked in ethanol at 70%. Bear in mind that the excess of ethanol can cause conjunctiva and cornea irritation. Additionally, we recommend the implementation of measures related to skin and mucosa care during work<sup>[212,251]</sup>. (read [question 6.4.](#), section 6) (*Level IV, Class I*)

## ***Key Points***

- In the event of accidental cutaneous exposure to biological contaminants from a COVID-19 patient, you should immediately perform antisepsis with ABAS. ***(Level IV, Class I)***
- In the event of accidental exposure of the ocular conjunctiva, you should remove your goggles, wash with water and a saline solution and decontaminate the periocular area with a 70% ethanol solution. ***(Level IV, Class I)***
- We recommend prophylactic measures to preserve skin and mucosa integrity during working hours. ***(Level IV, Class I)***

## Conclusion

(Appendix IV to read the recommendations in the form of a schedule, as well as check-lists to support you during adaptation phase)

## Future Perspectives

The group intends to answer all the questions concerning the different intervention stages, pre-, peri- and post-. Nonetheless, at this stage, we considered it was more prudent to establish an urgency hierarchy of response to the questions which intrigue health professionals and all their team the most. The critical evaluation of the evidence according to standardized systems shall also be a long-term aim, once the evidence on this topic is growing exponentially, with methods increasingly structured.

Presently, there are surveys in the field of seasonal variation of SARS-CoV-2 transmissibility, once it has the potential to modulate the spread of the virus. The spread of the virus in tropical climates, such as Singapore, is well known, and, however, winter is not a necessary condition to the spread of SARS-CoV-2. Furthermore, these surveys expose issues which may be preponderant to establish future attitudes and prevention measures<sup>[181]</sup>:

Reductions in the prevalence may be related to the seasonal variation of transmissibility, without neglecting well-succeeded interventions;

Subpopulation dynamics is very uneven in terms of counting methods of positive cases;

Seasonal variation can be a window of opportunity for a better preparation of health systems to fight a new outbreak;

SARS-CoV-2 may become an endemic seasonal CoV, similar to *Influenza* virus.

The future envisages the adaptation and adoption of new measures, such as *Big Data* and telemedicine.

*Big Data* is gaining international dimension in the perception of the necessary measures to monitor and detect prematurely new outbreaks of infectious diseases. It is a public health vehicle to detect, report and control emerging outbreaks. It is based on data sets and computational approaches which allow to reveal trends, patterns and predictive

probabilities. It is based on the Vs triad: volume (amount of data collected); velocity (rate at which data are collected and spread); variety (multiplicity of sources used to compile data)<sup>[182]</sup>.

Telemedicine allows to treat patients in their own homes with a great virtual medical support. It was a solution designed with the aim of facilitating medical consultations in disaster periods and public health emergencies<sup>[183]</sup>. It is necessary to understand its applicability in dentistry; nonetheless, in pandemic times caused by SARS-CoV-2 it has been an essential tool in patients' triage process.

SARS-CoV-2 will be a milestone in the history of humankind and it made health professionals aware of the importance, even grater nowadays, of continuous monitoring and early detection. The greatest challenge is indeed the time for the preparation and response to situations such as the one we live in<sup>[253]</sup>!

## Updating

Taking into account the experience of just 4 months with COVID-19, these Clinical Guidelines shall be object of continuous updating, and they can also be kept, revised or completely redrafted.

## Acknowledge

The authors would like to acknowledge NSK, Tokyo, Japan for supporting the translation of this Clinical Guidelines.



# Appendix I

Advisory Commission

## | *Advisory Board*

### *Doctor Ana Lúcia de Pereira Neves Messias*

Guest Teaching Assistant at the Faculty of Medicine of the University of Coimbra

### *Professor Professor Aníbal Diógenes*

Associate Professor at the University of Texas, *Health Center of San Antonio*, United States of America

### *Doctor and Professor António Vaz Carneiro*

Full Professor of Medicine at the Faculty of Medicine of the University of Lisbon and Director of Cochrane Portugal

### *Doctor and Professor Carlos Robalo Cordeiro*

Director of the Faculty of Medicine of the University of Coimbra

### *Doctor and Professor Gil Alcoforado*

Full Professor

### *Doctor and Professor Ion Zabalegui*

Guest Professor at the University *Complutense* of Madrid, Spain

### *Doctor José Manuel da Costa Rodrigues de Oliveira*

Dentist

### *Doctor and Professor José Frias Bulhosa*

Representative of the Portuguese Dentist National Association of the National Health Service

### *Doctor Luís António Silva Ferreira*

Dentist

### *Doctor and Professor Luís Pires Lopes*

Head of the Faculty of Dental Medicine of the University of Lisbon

### *Doctor and Manuel Gameiro*

Full Professor at the Mechanical Engineering Department of the University of Coimbra

### *Doctor and Professor Maria Helena Raposo Fernandes*

Full Professor at the Faculty of Dental Medicine of the University of Oporto

*Doctor and Professor Mário Bernardo*

Coordinator of the Integrated Master Degree in Dental Medicine at the Faculty of Dental Medicine at the University of Lisbon

*Doctor and Professor Mário Bernardo*

Full Professor at the University Complutense of Madrid, Spain

*Doctor and Professor Marlene Barros*

Head of the Faculty of Dental Medicine of the Catholic University of Portugal

*Doctor Paulo Manuel de Oliveira Nunes*

Dentist

*Doctor Nuno Ventura*

Dentist

*Doctor Ricardo Jorge Martins*

Dentist

*Doctor and Professor Sandra Gavinha*

Head of the Faculty of Health Sciences of Fernando Pessoa University



## Appendix II

### Materials and Methods

## Section 1 | Questions

<b>PRE-</b>
1.1. How can you manage the schedule?
1.2. Should the patient be alerted to be on time for the appointment?
1.3. Should the patient be alerted to avoid bringing personal ornaments to the appointment?
1.4. Should the patient be alerted to avoid paying the consultation in cash?
1.5. What are you supposed to do if a patient asks for a teleconsultation?
2.1. Which questions must be asked while making an appointment by telephone?
2.2. If the patient answers in the affirmative to any of the questions included in the questionnaire referred to in 2.1, but refers absence of fever, how is scheduling supposed to be handled?
2.3. While scheduling an appointment for a patient having a professional profile of risk, as is the case of health professionals, or hospital environment, day care center or retirement homes employees, should there be any particular care?
2.4. The patient refers having been infected with SARS-CoV-2 but they have already tested negative since. How is one expected to proceed?
2.5. How should one tackle the situation of a patient having a clinical history of COVID-19 and who is in home isolation?
2.6. A patient goes spontaneously to the clinic to schedule an appointment without having previously called. What is the receptionist supposed to do?
2.7. Should the accompanying person be advised not to come, whenever possible, and/or avoid their permanence in any place of the clinic?
2.8. If the patient comes accompanied, is it necessary for the accompanying person to go through the triage for COVID-19?
3.1. How can you make means of disinfection and protection available to those entering the clinic?
3.2. Which special measures must be applied in this customer service area?
3.3. How should you organize the waiting room?
3.4. How many people can remain in the waiting room? What is the advised capacity?
3.5. Which furniture can there be in the waiting room?
3.6. Should magazines or newspapers be available?
3.7. Should the TV and air conditioning remote controls be available?
3.8. Should the water dispenser or the coffee machine be available?
3.9. Which changes must be made in the sanitary facilities?
3.10. Which means of decontamination must be available at the sanitary facilities?

4.1. Should there be physical barriers between the reception and the user?
4.2. Which signposting can indicate the organization of the waiting spaces or the interdiction of crossing without permission?
4.3. Which informative billboards on COVID-19 must be posted and where?
5.1. What is the rationale behind hand washing?
5.2. How and when should you wash your hands?
5.3. Which products must be used to sanitize the hands?
5.4. Which are the methods that keep hands' skin healthy?
5.5. Should the patient be allowed to brush their teeth, prostheses or removable braces before stepping inside the consultation office?
6.1. How often should you clean?
6.2. How should you clean surfaces?
6.3. How should you do this cleaning?
6.4. Which cleaning/disinfection products must be used?
6.5. What are the most critical surfaces?
6.6. How often should the most critical areas or surfaces be cleaned?
6.7. What are the materials you must use while cleaning the most critical areas?
6.8. How often must you clean the sanitary facilities?
6.9. How should you clean the sanitary facilities?
6.10. Which are the most critical surfaces in the sanitary facilities?
6.11. Which products must be used to clean the sanitary facilities?
6.12. What are the materials that you must use to clean the sanitary facilities?
7.1. How can the receptionist keep a social distancing, minimizing the fear that this measure might arouse on the patient?
7.2. What should you do after finishing serving a patient/accompanying person and before serving the following one?
7.3. Should there be separate entrances for users and professionals?
7.4. What is the minimum mandatory recommendation of personal protective equipment at the dental clinic?
8.1. Am I able to go to work?
8.2. I have a relative (who is part of my household) who is sick with COVID-19. Can I go to work?
8.3. I am pregnant or more than 65 years old, I suffer from related systemic pathologies of higher risk. Can I go to work?
8.4. How is it possible to control if the staff are infected?
9.1. What should be checked as soon as the patient enters the clinic?

9.2. The patient is feverish or reports having had fever and cough or sore throat and muscular pains over the last 24h. What should you do?
9.3. The patient is not feverish. What should you do next?
9.4. Can the patient come accompanied?
9.5. Should the accompanying person wear a mask too?
9.6. In case the accompanying person walks into the clinical office, where should they remain and how should you proceed?
9.7. In case the accompanying person does not enter the clinical office, where should they wait?
9.8. What should you do with the patient's belongings (and the accompanying person's as well, if they happen to walk into the consultation office) before putting on the appropriate PPE?
10.1. How should maintenance technicians be welcome?
11.1. How should delivery men be welcome?
12.1. How should waste collection employees be welcome?
13.1. What is the appropriate protective equipment?
14.1. What is the appropriate protective equipment?

## **PERI-**

15.1. Should risk areas be identified at the clinic?

15.2. What are the differentiated recommendations for the different risk areas?

15.3. How should the general procedures in a clinic be ranked in terms of risk?

15.4. How should the different clinical procedures be ranked in terms of risk?

16.1. What is the stability of the virus in the atmosphere?

16.2. What is the reach of contaminated aerosols released in normal breathing in a closed environment?

16.3. Which distance may reach the aerosols generated during the operative procedures in Dentistry?

16.4. How should dental equipment and the surrounding surfaces be protected?

16.5. How should the disinfection of the several components and infrastructures of the consultation office be carried out?

16.6. What is the most suitable ventilation system in a clinical office?

16.7. Is the use of additional systems of purification of biological contaminants in the air of the clinical office justified?

16.8. What must the acclimatization conditions at the clinical office be?

16.9. Is it advisable to use means of microbiological decontamination in the water supply systems and water reservoirs of the equipment?

16.10. What is the effectiveness of suction systems in the reduction of the generation of aerosols at the source of contamination?

16.11. Does the use of antimicrobials in the irrigation system of the ultrasonic devices reduce the risk of contamination during aerosolization?

17.1. Must the PPE to be worn be defined by the risk involved in the procedure?

17.2. What is the appropriate PPE according to the levels of protection?

17.3. Which type of masks are there? And what is the indication for its clinical use?

17.4. What should you do before putting the PPE on?

17.5. How are you supposed to put the PPE on?

17.6. What care should be taken over the use of PPE?

17.7. What are you supposed to do if the first pair of gloves (inner pair in degree 3 protection PPE) touches a potentially contaminated area?

18.1. How can you reduce microbiological burden in intraoperative stage?

18.2. Should a perioral decontamination be performed? How should it be performed?

18.3. What is the importance of using a rubber dam?

18.4. Should priority be given to the use of single-dose materials?

19.1. What is the impact of operative instruments on aerosol generation?

19.2. Which specifications should cutting/preparation operating instruments have? Are non-return systems better?

## **POST-**

20.1. I have finished the operative procedure. What comes next?

20.2. Which are the critical surfaces?

20.3. How can you minimize contamination and perform the cleaning of critical areas?

20.4. Should protection barriers in screen surfaces, keyboards and mice be removed between patients?

20.5. What care should I take with the casing of the digital imaging sensor and the intraoral x-ray films?

20.6. How should X-ray positioners be disinfected?

20.7. How should digital imaging sensors, intraoral cameras and scanners be protected and disinfected?

20.8. Should the purge of rotary instruments be performed before being removed to be sterilized?

20.9. Is chemical external disinfection of turbines, counter-angles and scalers enough against SARS-CoV2 or should they be sterilized between patients?

20.10. What is the sequence for physical barriers and rotary instruments removal?

20.11. How should physical barriers to protect the equipment be eliminated?

20.12. Regarding surface disinfection (equipment and workbenches), which product is more efficient and for how long should contact time be?

20.13. Is UV radiation an effective sterilization method?

20.14. Regarding wrapping and packaging of the consumables used, which is the most suitable disinfection product and what is its contact time?

20.15. Is it necessary to make the suitable aeration of the clinical office before starting the following consultation?

20.16. How often should the clinical office's floor be cleaned and disinfected?

20.17. Regarding the clinical office's floor, which product should be used and for how long should it be left in contact to achieve a suitable disinfection?

20.18. How should the clinical office cleaning be performed?

20.19. Which cleaning/disinfection products should be used?

20.20. Which types of presentation should cleaning products have?

20.21. What is the most suitable disinfectant to be used in the form of wipes?

20.22. How should surfaces be cleaned in the event they are contaminated with blood or other organic products?

21.1. What is the protocol to disinfect dental impressions?

21.2. What is the protocol to disinfect dental stone models?

21.3. What is the protocol for try-in devices (copings, registration waxes, preliminary crowns/bridges and removable prosthesis) disinfection?
21.4. What is the protocol to disinfect pre-existing prosthesis?
22.1. What is the sequence to remove PPE in each one of the levels (1, 2 and 3)?
22.2. Where should disposable PPE be placed after being removed? In what type of container?
22.3. What is the protocol to remove gloves?
22.4. When and how should the dentist remove the surgical mask?
22.5. When should the assistant remove PPE?
22.6. Can FFP2 (N95) and FFP3 respirators be reused?
22.7. In which place should PPE be removed? In the clinical office or somewhere else?
22.8. Where should the assistant place PPE after being removed?
22.9. How should the face shield and the goggles be washed and disinfected?
22.10. Can uniforms be worn again the following day without being washed?
22.11. What kind of treatment should be given to non-critical fabric materials: uniforms?
22.12. After a day at work how should you behave when you get home?
23.1. Does the use of a surgical mask by the patient inside the clinic reduce the risk of cross contamination with SARS-CoV-2?
23.2. Should patients' movement inside the clinical office be limited after the operating procedures?
23.3. Should additional information be provided to the patient in digital form instead of paper form?
23.4. How should hand antisepsis be performed before leaving the office, regarding the dentist, the assistant, the patient/accompanying person?
24.1. Should the patient keep social distance at the reception and means of protection after consultation?
24.2. Should the patient keep the mask on in the reception area after the consultation?
24.3. Which important additional information should be provided to the patient by the receptionist, in the context of COVID-19 infection, in the post-consultation stage?
24.4. Which information should be confirmed to the patient by the receptionist before leaving the clinic?
25.1. Which payment methods and receipt issuing methods should be recommended?
26.1. Does contamination with SARS-CoV-2 imply a change of the procedures used to clean, decontaminate and sterilize materials and instruments?
26.2. Which kind of bag is indicated to pack group III waste?
26.3. Should the placement of group III waste in the corresponding container be performed in the clinical office or in the sterilizing room?



26.4. Does contamination with SARS-CoV-2 imply a change of the procedures used to handle and process waste?

26.5. Should the material/instruments handled by the dentist be considered as semi-critical or critical?

26.6. Should the material/instruments handled by the dental assistant be considered as semi-critical or critical?

26.7. Is it possible to disinfect FFP2 (N95) and FFP3 respirators?

27.1. Is the oral cavity a tropism location for SARS-CoV-2?

27.2. In the event of an accidental exposure to biological contaminants from a COVID-19 patient, which procedure should be adopted?

## Section 2 | Bibliographic Research

Table 1 - Algorithm established by the proofreader - Stage 1 (AV; CM)

1	SARS.ti,ab,kw
2	MERS.ti,ab,kw
3	SARS-Co-2.ti,ab,kw
4	COVID-19 or COVID-19 or coronavirus*.ti,ab,kw
5	OR/3,4
6	Influenza.ti,ab,kw
7	Dentistry/
8	AND/1,7
9	AND/2,7
10	AND/5,7
11	AND/6,7
12	Aerosol.ti,ab,kw
13	Air.ti,ab,kw
14	OR/12, 13
15	AND/1, 14
16	Ventilator.ti,ab,kw
17	Oral hygiene.ti,ab,kw
18	OR/16, 17
19	AND/7, 18

Table 2 - Algorithm adapted from NICE - Stage 2<sup>[1]</sup>

1	exp coronavirus/
2	((corona* or corono*) adj1 (virus* or viral* or virinae*)).ti,ab,kw.
3	((coronavirus* OR coronovirus* OR coronavirinae* OR Coronavirus* OR Coronovirus* OR Wuhan* OR Hubei* OR <b>huaian</b> OR "2019-nCoV" OR 2019nCoV OR nCoV2019 OR "nCoV-2019" OR "COVID-19" OR COVID19 OR "CORVID-19" OR CORVID19 OR "WN-CoV" OR <b>Inco</b> OR "HCoV-19" OR HCoV19 OR CoV OR "2019 novel*" OR ncov OR "n-cov" OR "SARS-CoV-2" OR "SARSCoV-2" OR "SARSCoV2" OR "SARS-CoV2" OR SARSCov19 OR "SARS-Cov19" OR "SARSCov-19" OR "SARS-Cov-19" OR <b>ncolor</b> OR Ncorona* OR Ncorono* OR NcovWuhan* OR NcovHubei* OR NcovChina* OR NcovChinese*) .ti, ab, kw.) all
4	((respiratory* adj2 (symptom* or disease* or illness* or condition*)) or "seafood market*" or "food market*") adj10 (Wuhan* or Hubei* or China* or Chinese* or Huanan*).ti,ab,kw.
5	((outbreak* or wildlife* or pandemic* or epidemic*) adj1 (China* or Chinese* or Huanan*).ti,ab,kw.
6	"severe acute respiratory syndrome*".ti,ab,kw.
7	or/1-6
8	limit 7 to yr="2019 -Current"

### Section 3 | Articles Included

Reference	Author, Year	Participants	Intervention	Comparison	Outcome	Remarks
1	Guerra F. 2020	Pacientes infectados ou não pelo vírus SARS-CoV-2	Any type of dental procedure, with or without aerosol production, in a pandemic situation due to SARS-CoV-2	Any type of dental procedure, with or without aerosol production, before the SARS-CoV-2 pandemic	Dental Medicine practice and practice guidelines after general confinement	-
2	NICE. 2014	-	-	-	NICE Guidelines elaboration manual	-
3	Rosenfeld RM. 2009	-	-	-	Manual de desenvolvimento de uma Guideline	-
4	NICE. 2020	-	-	-	Guidelines for the preparation of quickl NOC	-
5	El Zowalaty ME, Jarhult. 2020	-	-	-	SARS-CoV-2 as a potential pandemic	-
6	Xu J, Chen Y, et al. 2020	-	-	-	Coronavirus: a global pandemic?	-
7	Sohraibi C, Alsafi Z, et al. 2020	-	-	-	Coronavirus - the 21st century pandemic	-
8	Amodio E, Vitale F, et al. 2020	-	-	-	SARS-CoV-2 -an emerging pandemic	-
9	Chan KW, Wong VT,	-	-	-	SARS-CoV-2: epidemiology, clinical aspects, preventive and therapeutic measures	-
10	Coronaviridae Study Group of the International Committee on Taxonomy 2019	-	-	-	Classification of the new coronavirus	-
11	Guan W, Ni Z, et al. 2020	-	-	-	Clinical features of the SARS-CoV-2 virus, diagnosis	Screening process for patients / companions
12	Spagnuolo G, de Vito D, et al. 2020	-	-	-	Care to be taken before scheduling an emergency appointment	-
13	Ungaro RC, Sullivan T, et al. 2020	-	-	-	Prevention measures related to the new coronavirus	-
14	Borges Nascimento IJ 2020	-	-	-	SARS-CoV-2 - inter-human transmission	-
15	WHO 2020	-	-	-	Screening of patients. EPI's according to the patient's risk, 5 key moments of hand washing, empirical precautions, aerosol formation precautions, surface cleaning	-
16	Xia N, Wang G, et al. 2020	-	-	-	Serological tests	-
17	Meng L, Hua F, Bian Z. 2020	-	-	-	Infection control in consultation with MD + recommendations	-

18	Sahni V, 2020	-	-	-	Possible relationship between COVID-19 and Periodontal Disease - cytokine expression	Letter to Editor
19	Takahashi Y. 2020	-	-	-	Possible worsening of symptoms by COVID-19 due to the presence of periodontal bacteria; Contribution of oral hygiene care to prevention	-
20	Manzalawi R. 2020	SARS-CoV-2 positive patients	-	-	Gingival hemorrhage as a sign of SARS-CoV-2 infection	Case report
21	Di Spirito F. 2020	-	-	-	Oral manifestations associated with SARS-CoV-2 infection and the importance of early diagnosis of oral lesions by specialists in oral medicine and pathology	Letter to Editor
22	Badran Z. 2020	-	-	-	Periodontal pockets as possible reservoirs for the SARS-CoV-2 virus	-
23	Xiang Z. 2020	-	-	-	Oral locations as potential reservoirs for SARS-CoV-2 and implications for changes in host microbiota	-
24	Amorim dos Santos J. 2020	SARS-CoV-2 positive patients	-	-	Prevalence of oral lesions / changes in positive SARS-CoV-2 patients	-
25	Bemquerer LM. 2021	-	-	-	Correlation between oral dermatological lesions in positive SARS-CoV-2 patients	Letter to Editor
26	Corchuelo J. 2020	SARS-CoV-2 positive patients (asymptomatic)	-	-	Presence of oral lesions in asymptomatic positive SARS-CoV-2 patients	Case report
27	Brandão TB. 2021	SARS-CoV-2 positive patients	-	-	Presence of oral lesions in positive SARS-CoV-2 patients; possible consideration of the oral cavity as a target for infection with this virus	Case series
28	Riad A. 2020	SARS-CoV-2 positive patients	-	-	SARS-CoV-2 positive patients with mucositis: possible relationship?	Case series
29	Pérez-Sayáns M. 2021	-	-	-	COVID-19 Tongue; pathognomonic of COVID-19?	Letter to Editor
30	Hathway RW. 2021	-	-	-	COVID-19 Tongue	Letter to Editor
31	Santosh ABR. 2020	-	-	-	Possible transmission of COVID-19 disease through saliva in asymptomatic patients	Letter to Editor
32	Sarode GS. 2020	SARS-CoV-2 positive patients	-	-	Efficacy of salivary and nasopharyngeal samples in the detection of SARS-CoV-2	-
33	Medeiros da Silva RC. 2020	-	-	-	Salivary samples as a means of diagnosing SARS-CoV-2 infection	-
34	Fernandes LL. 2020	SARS-CoV-2 positive patients	Salivary Samples	-	Salivary samples as a means of diagnosing SARS-CoV-2 infection	-
35	Michailidou E. 2020	-	-	-	Salivary samples as a means of diagnosing SARS-CoV-2 infection, salivary diagnostic methods	-
36	Kashiwagi K. 2021	-	-	-	Salivary samples as a means of diagnosing SARS-CoV-2 infection	Case report
37	Hamid H. 2020	-	-	-	Salivary samples as a means of diagnosing SARS-CoV-2 infection	-

38	Chen L. 2020	-	-	-	Salivary samples as a means of diagnosing SARS-CoV-2 infection and classification of oral symptoms	-
39	Hung K. 2020		Salivary Samples	Nasopharyngeal samples	Salivary samples as a means of diagnosing SARS-CoV-2 infection	-
40	Azzi L. 2021	-	Salivary Samples	Nasopharyngeal samples	Salivary samples as a means of diagnosing SARS-CoV-2 infection; virus detection techniques in samples	-
41	Fakheran O. 2020	-	-	-	Salivary samples as a means of diagnosing SARS-CoV-2 infection	-
42	Altawalah H. 2020	Patients suspected of having SARS-CoV-2 infection	Salivary Samples	Nasopharyngeal samples	SARS-CoV-2 detection via RT-PCR	Cross-sectional study
43	Czumbel LM. 2020	SARS-CoV-2 positive patients	Salivary Samples	Nasopharyngeal samples	Consistency and reliability of viral RNA detection in salivary samples	Metanalysis
44	Gupta S. 2021	SARS-CoV-2 positive patients	Sample of crevicular fluid	Nasopharyngeal and salivary samples	Detection of viral RNA	-
45	Droegemeier K. 2020	-	-	-	Stabilization on SARS-CoV 2 surfaces, incubation period	-
46	Bai Y, Yao L, et al. 2020	-	-	-	Transmission of SARS-CoV-2 in asymptomatic patients	-
47	Wang Q, Yu C. 2020	-	-	-	Endemic areas of SARS-CoV-2	-
48	Zou L, Ruan F, et al, 2020	-	-	-	Viral load of the upper respiratory tract by SARS-CoV-2	-
49	National Research Council 2020.	-	-	-	Bioaerosols: potential for permanence on Sars-Cov-2 surfaces	-
50	CDC 2020	-	-	-	SARS-CoV-2 infection prevention protocol	-
51	Huang C, Hang Y, et al, 2019	-	-	-	Clinical aspects of patients infected with SARS-CoV-2	-
52	Santarpia J, Rivera D, et al. 2020	-	-	-	Transmission of SARS-CoV-2 in saliva	-
53	Li Y, Leung GM, et al. 2020	-	-	-	Air duration control: space ventilation, negative pressure, etc.	-
54	Peng X, Xu X, et al. 2020	-	-	-	Mechanisms of transmission of the Covi-19 virus in dentistry, as well as pre, peri and postoperative prevention measures	-
55	Zhu F. 2020	Adult patients who have not contracted COVID-19 disease	Ad5 vectored COVID-19 vaccine ( $5 \times 10^{10}$ , $1 \times 10^{11}$ , and $1 \cdot 5 \times 10^{11}$ viral particles)	-	Adverse effects 7 days after vaccine administration	Dose-escalation, open-label, non-randomised, first-in-human trial
56	Zhao J. 2020	-	-	-	Updates vaccination COVID-19	-
57	Awadasseid A. 2021	-	-	-	Updates vaccination COVID-19	-
58	Kim KD. 2020	-	-	-	Updates vaccination COVID-19	-
59	DGS. 2020	-	-	-	Norma 021/2020: Vaccination Campaign against COVID-19 COMIRNATY® Vaccine	-

60	DGS. 2021	-	-	-	Norma 001/2021: Vaccination Campaign against COVID-19 COVID-19 Vaccine MODERNA®	-
61	DGS. 2021	-	-	-	Norma 003/2021:Vaccination Campaign against COVID-19 COVID-19 Vaccine AstraZeneca®	-
62	Centers for Disease Control and Prevention. 2021	-	-	-	Updates vaccination COVID-19	-
63	Bernal JL. 2021	-	-	-	Vaccination with BNT162b2 mRNA and ChAdOx1 adenovirus vector	-
64	Brandolin BA. 2020	-	-	-	Impact of COVID-19 in the dentistry sector	-
65	Banakar M. 2020	-	-	-	SARS-CoV-2 infection prevention protocol	-
66	Bizzoca ME. 2020	-	-	-	Changes in Dentistry practice during the pandemic by COVID-19	-
67	Kathree BA. 2020	-	-	-	Impact of COVID-19 in the dentistry sector	-
68	Coulthard P. 2020	-	-	-	Impact of COVID-19 in the dentistry sector	-
69	Wijesooriya NR. 2020	-	-	-	Changes and adaptations in the practice of Dentistry after the emergence of COVID-19: education, teleconsultation	-
70	Crawford E. 2020	-	-	-	Efficacy and usefulness of telemedicine	-
71	Ghai S. 2020	-	-	-	Telemedicine	-
72	Telles-Araujo GT. 2020	-	-	-	Telemedicina como meio de suporte à pandemia por COVID-19	-
73	Tapia ROC. 2020	SARS-CoV-2 positive patients	-	-	Description of oral lesions	Case series
74	Petruzzi M. 2016	-	-	-	Telemedicine	-
75	Saccomanno S. 2020	-	-	-	Orthodontics and Telemedicine	-
76	CDC. 2005	-	-	-	Mycobacterium transmission prevention guidelines	-
77	Administração Central do Sistema de Saúde IP. 2008	-	-	-	Technical Specifications for HVAC Installations.	ET 06/2008 V.2014
78	CDC. 2003	-	-	-	Appendix B - air quality and cross infection control	-
79	New York Stat Department of Health. 2014	-	-	-	Guidelines for building hospitals and clinics.	-
80	US Department of Veterans Affairs. Office of Construction & Facilities Management. 2014	-	-	-	Guide to dental clinics	-

81	Harrel SK, Molinari J. 2004	-	-	-	Aerosols in Dentistry: implications for infection control	-
82	Helmis CG, Tzoutzas J, et al. 2007	-	-	-	Air quality in the dentistry office	-
83	Li Y, Leung GM, et al. 2007	-	-	-	The role of ventilation in the airborne transmission of infectious agents in the built environment - a systematic multidisciplinary review	-
84	Raghunath N, Meenakshi S. 2016	-	-	-	Aerosols in Clinical Practice	-
85	Qian H, Zheng X. 2018	-	-	-	Ventilation control for air transmission of human exhaled bio-aerosols in buildings	-
86	Silva, MG. 2020	-	-	-	Modes of transmission of the SARS-CoV-2	-
87	OSHA. 2020	-	-	-	Guidelines for preparing different sectors of post-pandemic work	Risk definition of the procedures, of the different classes
88	WHO. 2020	-	-	-	Guidelines for monitoring patients suspected of SARS-CoV-2 infection	-
89	Bioseguridad para la Atención Odontológica COVID 19 2020	-	-	-	Measures aimed at prevention: Telemedicine, Instructions for consultation, admission to the clinic, reception, waiting room, bathroom	-
90	Zhang W, Jiang X. 2020	-	-	-	Prevention and control of corona virus in dental clinics - control of medical teams, screening of patients	-
91	Huh S. 2020	-	-	-	PPE's for the treatment of COVID positive patients	-
92	van Doremalen N, Bushmaker T, et al. 2020	-	-	-	Aerosol stability in sars-cov 1 and 2	-
93	DGS. 2020	-	-	-	Orientação 022/2020: COVID-19:Procedures in Clinics, Offices or Oral Health Services of Primary Health Care, Social and Private Sector.	-
94	Lurie N, Carr BG. 2020	-	-	-	Telemedicine in disaster response and support	-
95	Reeves JJ, Hollandsworth HM. 2020	-	-	-	Describe the implementation of an important technological support system to optimize clinical management during the COVID-19 pandemic.	-
96	Lv N. 2020	-	-	-	Monitoring of emergency consultations during COVID-19	-
97	León S. 2020	-	-	-	Inequalities in access to health care for the elderly	-
98	DGS. 2020	-	-	-	SARS-CoV-2 Infection Prevention and Control (COVID-19): PPE	Norma 007/2020 29/03/2020
99	Smales FC, Samaranyake LP. 2003.	-	-	-	Maintenance of MD consultations in sars-cov1 (2003)	-



100	Li RW, Leung KW, et al. 2004	-	-	-	Recommendations in the management of sars-cov-1 patients (possible cases, unprotected contacts, contacts with sars-cov 1 patients in less than 10 days or asymptomatic patients; cross-infection control measures	Coughing and sneezing can produce aerosols from 0.001 µm to 10,000 µm
101	Samaranayake LP, Peiris M. 2004	-	-	-	Implications of sars-cov-1 in dentistry: identifying patient, hand hygiene, EPI's	Mouthwash with chx between 0.12 and 2% before consultation
102	Huang Z, Zhao S, et al. 2020	-	-	-	Establish management strategies and protocol for the performance of procedures in the radiology department during the COVID-19 outbreak	Strategies for staff and patients
103	Lai T, Tang E, et al. 2020	-	-	-	Proposed model to be adopted in the ophthalmology consultation to prevent covid cross infection (pre, peri)	Includes questionnaire to be made to the patient before the consultation
104	Maragakis, L. 2020	-	-	-	Questionnaire that helps the patient to understand whether he or she may be infected with SARS-CoV-2 and whether or not to schedule an appointment.	-
105	Repici A, Maselli R, et al. 2020	-	-	-	Use of PPE's according to the patient's risk; patient's clothing; OS PPE's: depending on the risk; how to dress and remove. Negative pressure room treatments	High and low endoscopy. Possible extrapolation to dentistry?
106	Zhang Y, Sun Z, et al. 2020	-	-	-	Procedures adopted in a hospital in China to control patients infected with COVID-19	The questionnaire at the end seems to me relevant for the classification of individuals at risk - I leave it up to the experts to consider the relevance of the evidence for inclusion in the recommendations.
107	DGS. 2020	-	-	-	Norma 020/2020: COVID-19: COVID-19 Case Definition.	-
108	DGS. 2020	-	-	-	Norma 004/2020: COVID-19: Patient Approach with Suspected or Confirmed COVID-19.	-
109	DGS. 2020	-	-	-	COVID-19: Mitigation Phase. Procedures for Residential Structures for the Elderly (ERPI), Integrated Continuous Care Units (UCCI) of the National Network of Continuing Care (RNCCI) and other responses dedicated to the elderly; institutions for the care of children and young people at risk.	Orientação 009/2020 de 11/03/2020. Updated to 07/04/2020.
110	ADA 2020	-	-	-	Minimize COVID-19 transmission risk	-
111	ECDC. 2020	-	-	-	Criteria for the identification of COVID-19	-
112	DGS. 2020	-	-	-	SARS-CoV-2 infection (COVID-19) - Social detachment and isolation	Orientação 010/2020 de 16/03/2020.
113	OMD 2020	-	-	-	Care of patients with COVID-19 or high risk of contagion	-
114	OMD 2020	-	-	-	Preventive care and Dentistry consultations	-
115	DGS. 2020	-	-	-	SARS-CoV-2 infection (COVID-19): Measures to prevent transmission in public service establishments	Orientação 011/2020 17/03/2020

116	Boyce JM, Pittet D, et al. 2002	-	-	-	Guidelines for hand hygiene in healthcare institutions	-
117	Guangming Y, Hualiang L, et al. 2020	-	-	-	Air contamination by the SARS-CoV-2 virus	Calling attention to the importance of PPE
118	CDC 2020	-	-	-	Guidelines for infection control in hospital settings	-
119	Consejo Dentistas. Organizacion Colegial de Dentistas de España 2020	-	-	-	Guidelines of the College of Dentists for action in the pandemic COVID 19	-
120	CDC. 2019	-	-	-	Guidelines for cleaning healthcare facilities	-
121	Lotfinejad N, Peters A, et al. 2020	-	-	-	Recommendations regarding the importance of hand washing in a pandemic period due to COVID-19	-
122	Allegranzi B, Bagheri NS, et al. 2011	220 artigos para revisão final	-	-	Developing countries - prevalence / incidence of infections (including general infections associated with health care, main sites of infection and microbiological causes)	-
123	Jefferson T, Del Mar CB, et al. 2011	66 artigos	-	-	Physical intervention measures to reduce or stop the spread of viruses associated with respiratory diseases	SARS-CoV-2 is a virus associated with Acute Respiratory Syndrome
124	Basile C, Combe C, et al. 2020	-	-	-	Guidelines for transmission control and monitoring practices for patients infected with COVID-19 in a dialysis center	-
125	WHO. 2009	-	-	-	Educational guidelines for the hand hygiene process. Awareness of its importance as a preventive measure	-
126	Picheansathian W. 2004	-	-	-	Effectiveness of hand washing solutions	-
127	CDC. 2003	-	-	-	Infection control in dental institutions	Adaptation to SARS-CoV-2
128	Sebastiani FR, Dym H, et al. 2017	-	-	-	Guidelines for infection control practices in the MD practice	Improve the effectiveness and impact of public health interventions.
129	Kampf G, Löffler H. 2003	-	-	-	Dermatological changes in hands exposed to SABA	-
130	DGS.2020	-	-	-	SARS-CoV-2 infection (COVID-19): Cleaning and disinfecting surfaces in public or similar establishments	Orientação 014/2020 21/02/2020
131	CDC. 2003	-	-	-	Guidelines for infection control in hospital settings	-
132	Kundrapu S, Sunkesula V, et al. 2012	215 rooms for patients with ICD and MRSA	Efficacy of disinfecting frequent touch surfaces in PS contamination	-	Daily disinfection of the surfaces of frequent touch reduces the acquisition of these pathogens in the hands of PS after contact with these surfaces	-
133	European Centre for Disease Prevention and Control 2020	-	-	-	Disinfection / Cleaning of healthcare facilities	-

134	Consejo Dentistas. Organizacion Colegial de Dentistas de España 2020	-	-	-	Guidelines of the College of Dentists for action in the pandemic COVID 19	-
135	Consejo Dentistas. Organizacion Colegial de Dentistas de España 2020	-	-	-	Recommendation of good practices for emergency dentistry appointments	-
136	Handbook of COVID19 Prevention and Treatment. 2020	-	-	-	Prevention of SARS-CoV-2: Isolation of areas, team management, individual protection, hospital protocols and digital support	-
137	Chughtai AA, Seale H, et al. 2020	-	-	-	Regulation for the use of masks by health professionals in hospitals	-
138	Giwa AL, Desai A, et al. 2020	-	-	-	Epidemiological, preventive, immunological characteristics of the COVID-19 pandemic	"Use of Personal Protective Equipment"
139	Leung NHL, Chu DKW, et al. 2020	-	-	-	Surgical masks for the transmission of SARS-CoV-2 and Influenza	-
140	Murthy S, Gomersall CD, et al. 2020	-	-	-	PS PPE for treatments with and without aerosols (at the level of COVID + patient treatment)	-
141	Ng K, Poon BH, et al. 2020	41 Health professionals exposed to COVID + patient aerosols	Exposure of at least 10 min at a distance of less than 2m	Use of surgical mask or N95 + Isolation with symptom monitoring	PS did not develop the disease. Negative covid test	Case report
142	Wang Q, Yu C. 2020	-	-	-	Indications for different masks	-
143	Chinese Center for Disease Control and Preventions	-	-	-	Key points for the prevention of health professionals against SARS-CoV-2	-
144	CDC 2019	-	-	-	Risk of exposure of doctors	-
145	Rimmer A. 2020	-	-	-	PPE in covid patients +	-
146	Rombolà G, Hedemperger M, et al. 2020	-	-	-	Practical indications for the prevention and management / handling of SARS-Cov-2, experienced-based	Extrapolation to COVID-19?
147	Krithikadatta J, Nawal RR, et al. 2020	-	-	-	Guideline of recommendations for the prevention of the dentist in the consultation of endodontics. Prevention of cross-infection in the community	-
148	Cowling BJ, Aiello A. 2020	-	-	-	Epidemiology, preventive measures in the community to combat the transmission / spread of COVID-19	-
149	DGS. 2020	-	-	-	Epidemiology, preventive measures in the community to combat the transmission / spread of COVID-19	Orientação n.º 012/2020 de 19/03/2020

150	WHO 2020	-	-	-	Guide for Health Professionals	-
151	-	-	-	-	COVID-19: Infection, prevention	-
152	ECDC. 2020	-	-	-	Infection, preparation and prevention measures for COVID-19 disease in healthcare settings	-
153	Association of Occupational health Professionals in Healthcare (AOHP) 2020	-	-	-	Infection through bioaerosols	-
154	Verbeek JH, Rajamaki B, et al. 2019	1950 patients in a total of 17 studies	21 interventions related to full body PPE, dressing and removing PPE; Compliance of health professionals	PPE How to dress and remove without self-contamination by healthcare professionals, evaluate OS training methods to improve compliance	Breathable PPE with low quality evidence, 2 pairs of gloves for removing PPE, face-to-face training have better rates in preventing self-contamination	Extrapolation to dentistry ?
155	Long Y, Hu T, et al. 2020	9171 patients from 6 RCTs	Effectiveness of N95 respirators	Effectiveness of surgical masks	N95 respirators are not associated with a lower risk of influenza	Study with Influenza
156	CDC. 2020	-	-	-	Risk monitoring guidelines in professional environments	-
157	CAL/OSHA	-	-	-	Aerosol-borne diseases	-
158	Pan Y. 2020	-	-	-	Vias de transmissão do vírus e medidas de contenção da disseminação do mesmo	-
159	ECDC. 2020	-	-	-	Guidelines for PPE placement and removal criteria when treating patients suspected of SARS-CoV-2 infection	-
160	ECDC. 2020	-	-	-	PPE needed to care for suspected or confirmed SARS-CoV-2 infection patients	-
161	ECDC. 2020	-	-	-	Safe use of PPE when treating infectious patients	-
162	Tang JW, Li Y. 2006	-	-	-	Factors involved in the transmission of aerosol infections and ventilation control	-
163	Asadi S, Wexler As, et al. 2019	-	-	-	Emission of aerosols during phonetics	-
164	Sociedad Española de Patología Digestiva; Asociación Española de Gastroenterología. 2020	-	-	-	General and individual guidelines in endoscopy units during the SARS-Cov-19 pandemic	Extrapolation to dentistry ?
165	Chin AWH, Chu JTS, et al. 2020	-	-	-	SARS-CoV-2 stability on surfaces	-
166	Moriarty LF, Plucinski, et al. 2020	-	-	-	SARS-CoV-2 control: isolation of a ship and its importance in Public Health	-

167	Ong SWX, Tan YK. 2020	3 symptomatic positive COVID 19 patients	Cleaning of space and material	Samples before and after decontamination * tests were not carried out in the 2 times nor were the same sites evaluated	Effectiveness of correct cleaning?!?!	Possible extrapolation to MD cabinet?
168	Jiang Y, Wang H, et al, 2020	-	-	-	Clinical data on hospital environmental hygiene monitoring and medical team protection during the 2019 coronavirus disease outbreak	-
169	Zayas G, Chiang MC, et al. 2012	45 participants	Voluntary cough (Evaluation of the precise and time-dependent distribution of droplet size using a laser diffraction system)	-	Droplets smaller than 1 micron represent 97% of the total number of droplets contained in the cough aerosol. Age, sex, weight, height and body mass have no statistically significant effect on the composition of the aerosol (size and number)	-
170	Bourouiba L. 2020	-	-	-	Cloud and aerosol production: respiratory transmission - implications and how to prevent	-
171	Bentley CD, Burkhart NW, et al. 1994	-	-	-	Study of sprays and aerosols produced in the dental medicine consultation	-
172	Bennett AM, Fulford MR, et al.2000	-	-	-	Exposure risk of dentists: transmission of diseases via aerosols produced in the consultation	-
173	Hokett, s. D., honey, j	-	-	-	Effectiveness of sensor protections	-
174	Swaminathan Y, Thomas J. 2013	-	-	-	Aerosols produced during the consultation are the main source of infection in the office, the main one being the control of infection.	-
175	Otter JA, Donskey C, et al. 2020	-	-	-	Survival of potentially pandemic viruses on surfaces	-
176	Rutala, WA, Weber DJ, et al. 2019	-	-	-	Disinfection and cleaning guidelines	-
177	Kampf G, Löffler H. 2003	-	-	-	Persistence of SARS-CoV-2 on inanimate surfaces	-
178	-	-	-	-	Diário da República, regulamentação do ar	-
179	NORMA ISO 7730/2005	-	-	-	Ergonomics of the thermal environment: Analytical determination and interpretation of thermal comfort through the calculation of PMV and PPD indices. Local thermal comfort criteria	-
180	Ijaz MK, Brunner AH, et al. 1985	-	-	-	Survival characteristics of SARS-CoV-2	-

181	Offner D, Floretti F, et al. 2016	-	-	-	Contamination of the dental team water system	-
182	Ji XY, Fei CN, et al. 2019	-	-	-	Bacterial infection of the dental chair water systems	-
183	Lizzadro J, Mazzotta M, et al. 2020	-	-	-	Cross-infection in the water lines of a dental unit	-
184	Desarda H, Gurav A, et al. 2014	-	-	-	Microbiological study on high-volume extractors in which their effectiveness is only registered when they are modified	-
185	Avasthi A. 2018	-	-	-	High-volume extractors for aerosol reduction in clinics	-
186	Van der Sluijs M, et al. 2016	-	-	-	Periodontal inflammation after non-surgical treatment	-
187	Jawade R. 2016	-	-	-	Effectiveness of different liquids in dental aerosols	-
188	Sethi KS. 2019	-	-	-	CHX vs. Cinnamon extract to reduce viral load in aerosols	-
189	Howard BE. 2020	-	-	-	Risk of aerosol production in the context of COVID-19	-
190	Zimmermann M. 2020	-	-	-	Monitoring of patients undergoing surgery in the context of a pandemic	-
191	Alharbi A. 2020	-	-	-	Guidelines for care in the context of dental medicine in the context of a pandemic	-
192	Ayyed AB. 2020	-	-	-	Guidelines for care in the context of dental medicine in the context of a pandemic	-
193	Falahchai M. 2020	-	-	-	Guidelines for care in the context of dental medicine in the context of a pandemic	-
194	Park, S H. 2020	-	-	-	Personal protective equipment in the context of a pandemic	-
195	Samaranayake LP. 2020	-	-	-	Personal protective equipment in the context of a pandemic	-

196	Jamal M. 2020	-	-	-	Guidelines for care in the context of dental medicine in the context of a pandemic	-
197	Agência Nacional de Vigilância Sanitária				Respiratory Protection Primer against Biological Agents for Health Workers	
198	Gallagher JE. 2020	-	-	-	Personal protective equipment in the context of a pandemic	-
199	Kantor J. 2020	-	-	-	Injuries associated with the use of PPE	-
200	Reis INR. 2021	-	-	-	Efficacy of preoperative mouthwashes in reducing the transmission of SARS-CoV-2	-
201	Castro-Ruiz C. 2020	-	-	-	Efficacy of povidone-iodine in reducing the risk of transmission of SARS-CoV-2	Narrative review
202	Carrouel F. 2021	-	-	-	Efficacy of preoperative mouthwash solutions in reducing salivary viral load	Critical review
203	Ather A. 2021	-	-	-	Efficacy of preoperative mouthwash solutions in reducing salivary viral load	Scoping Review
204	Yoon JG. 2020	-	-	-	Clinical importance of high doses of salivary SARS-CoV-2	-
205	Martínez Lamas L. 2020	-	-	-	Efficacy of povidone-iodine in reducing the risk of transmission of SARS-CoV-2	In vivo
206	Gottsauner MJ. 2020	-	-	-	Effectiveness of hydrogen peroxide in reducing salivary viral load	Prospective clinical pilot study
207	Seneviratne CJ. 2020	-	-	-	Efficacy of preoperative mouthwash solutions in reducing salivary viral load	Randomized control trial
208	Nejatidanesh F, Khosravi Z, et al. 2013	-	-	-	Risk of cross contamination of the face of dentists: the parts that are most exposed are the peri-nasal and peri-ocular areas. Face protection recommendations	-
209	Caldeira D, David C, et al., 2011	-	-	-	RS on antiseptics for the skin in the prevention of contamination in venous punctures.	-
210	Kamel C, McGahan L, et al. 2011	-	-	-	RS antiseptic skin preparation techniques - infection prevention	-
211	Samaranayake LP, Reid J, et al. 1989	-	-	-	Effectiveness of the rubber dam in controlling aerosol contamination	-

212	El-Din AMT, Ghoname NAEH. 1997	-	-	-	Use of the rubber dam in pediatrics	-
213	Al-Amad SH, Awad MA, et al. 2017	-	-	-	Effect of the levee on the control of bacterial transmission	-
214	CDC. 2020	-	-	-	Infection prevention guidelines	-
215	Day CJ, Sandy JR, et al. 2006	-	-	-	Types of particles produced during dental treatments	-
216	Hu, T. 2007	-	-	-	Risk of transmission of Hepatitis B in dental instruments	-
217	CDC. 2003	-	-	-	Cleaning and disinfection of rotating devices	-
218	Hamedani S. 2020	-	-	-	Guidelines for safe imaging protocols in the context of a pandemic by COVID-19	-
219	Doriguëtto PVT. 2020	-	-	-	Guidelines for safe imaging protocols in the context of a pandemic by COVID-19	-
220	Ilhan B. 2020	-	-	-	Guidelines for safe imaging protocols in the context of a pandemic by COVID-19	-
221	Tuttnauer Team 2020	-	-	-	Sterilization of dental materials	-
222	CDC. 2020	-	-	-	Provisional guidelines for infection control and prevention in patients with suspected or confirmed coronavirus disease 2019 (COVID-19) in health services	-
223	Hu, X. 2020	-	-	-	Thermal inactivation of the serum interferes with the immunoanalysis of antibodies to SARS-CoV-2	-
224	Colaneri M. 2020	-	-	-	Environmental contamination by SARS-CoV-2 in a hospital environment	-



225	United States Environmental Protection Agency Washington, D.C. 20460 Office Of Chemical Safety And Pollution Prevention List N	-	-	-	Online list of disinfectants against SARS-CoV-2	-
226	Gurzawska-Comis K. 2020	-	-	-	Guidelines on the prevention of nosocomial infection during the pandemic by COVID-19	-
227	Poggio C. 2020	-	-	-	Guidelines de desinfecção de superfícies no contexto pandémico pela COVID-19	-
228	Bali RK. 2020	-	-	-	Guidelines for disinfecting surfaces in the pandemic context by COVID-19	-
229	Dexter F. 2020	-	-	-	Instrument disinfection combined between disinfectants and UV light	-
230	Haralur SB, Al-Dowah OS, et al. 2012	-	-	-	Effect of alginate disinfection techniques on bacterial colony count	-
231	Azevedo MJ, Correia I, et al. 2019	-	-	-	Disinfection of added silicone impressions	-
232	Richard A. Bass, K.D. Plummer, et al. 1992	-	-	-	The effect of surface decontamination on the plaster model	-
233	Soares, C.R. and M. Ueti, et al. 2001	-	-	-	Influence of different disinfection methods on the physical properties of type IV and V plaster	-
234	Pisulkar SK, Dahane T, et al. 2018	-	-	-	Effect of saliva contamination of plaster models in terms of bacterial colony count	-
235	Da Silva FC, Kimpara ET, et al. 2008	-	-	-	Effectiveness of 6 disinfectants in removing acrylic biofilm from prostheses	-
236	CDC. 2020	-	-	-	Sequence of placement of PPE	-
237	CDC. 2019	-	-	-	Decontamination and reuse of masks / respirators	-
238	COEC. 2020	-	-	-	Emergency medicine in dentistry	-
239	CDC. 2019	-	-	-	Eye Protection	-

240	CDC. 2019	-	-	-	Cleaning and disinfection	-
241	Auer R, Cornelli G, et al. 2020	-	-	-	Changes in forms of payment in the face of the SARS-CoV-2 pandemic	-
242	Volgenant CMC. 2021	-	-	-	Guidelines for the prevention and reduction of the risk of transmissibility of SARS-CoV-2. Approach includes limiting aerosol production.	-
243	CDC. 2001	-	-	-	Sterilization and disinfection of dental instruments	-
244	Schwartz, A, et al. 2020	-	-	-	Decontamination and Reuse of N95	-
245	Hoffmann M, Kleine-Webber H, et al. 2020	-	-	-	Transmission routes of the SARS-CoV-2	-
246	Wolfel R, Corman VM, et al. 2020	-	-	-	Análise virológica do SARS-CoV-2: replicação ativa RNA	-
247	To K, Tsang O, et al. 2020	-	-	-	Virological analysis of SARS-CoV-2: active RNA will replicate	-
248	Liu L, Wei Q, et al. 2020	-	-	-	Transmission of SARS-CoV-2 through saliva	-
249	Xu H, Zhong L, et al. 2020	-	-	-	Expression of the ACE2 receptor on epithelial cells of the oral mucosa	-
250	Chinese Center for Disease Control and Preventions	-	-	-	Key points for the prevention of health professionals against SARS-CoV-2	-
251	Neher RA, Dyrdak R, et al. 2020	-	-	-	Future prospects - impact of the seasonality of the disease	-
252	Dion M, AbdelMalik P, et al. 2015	-	-	-	Big data in healthcare	Future prospects
253	Hollander JE, Carr BG. 2020	-	-	-	Telemedicine article	Include in future perspectives

254	Perry DJ. 2007	-	-	-	Levels of Evidence	-
255	Kirchhof, P. 2016	-	-	-	Recommendation classes	-

## Section 4 | Articles Excluded

Table 4 | Articles Excluded

Author (1st and 2nd), Year	Criterion	DOI/PMID
Kang L, Li Y <i>et al.</i> 2020	J	10.1016/S2215-0366(20)30047-X
Koh, D. 2020	J	10.1093/occmed/kqaa036
Tang B, Bragazzi NL, <i>et al.</i> 2020	J	10.1016/j.idm.2020.02.001
To KK, Tsang OT <i>et al.</i> 2020	A	10.1093/cid/ciaa149
Tay JY, Lim PL <i>et al.</i> 2020	A	10.1093/cid/ciaa179
Catton H. 2020	J	10.1111/inr.12578
Callisher C, Carroll D, <i>et al.</i> 2020	J	10.1016/S0140-6736(20)30418-9
Bouadma L, Lescure FX, <i>et al.</i> 2020	A	10.1007/s00134-020-05967-x
Ayittey FK, Ayttey MK <i>et al.</i> 2020	J	10.1002/jmv.25706
Chen Q, Liang M <i>et al.</i> 2020	J	10.1016/S2215-0366(20)30078-X
Kmietowicz Z. 2020	J	10.1136/bmj.m707
Imai H. 2020	J	10.1111/pcn.12995
Liu J, Liao X, <i>et al.</i> 2020	J	10.3201/eid2606.200239
Joob B, Wiwanitkit <i>et al.</i> 2020	J	10.1016/j.jhin.2020.02.016
Zhang S, Diao MY <i>et al.</i> 2020	J	10.1007/s00134-020-05977-9
Zhou P, Huang Z <i>et al.</i> 2020	A	10.1017/ice.2020.60
Xie J, Tong Z <i>et al.</i> 2020	A	10.1007/s00134-020-05979-7
Cao Y, Li Q <i>et al.</i> 2020	K	10.1111/acem.13951
Li X, Qian Y <i>et al.</i> 2020	A	10.1148/radiol.2020200661
Favre G, Pomar L, <i>et al.</i> 2020	E	10.1016/S1473-3099(20)30157-2
Lee IK, Wang CC, <i>et al.</i> 2020	J	10.1016/j.jhin.2020.02.022
Petersen E, Hui D, <i>et al.</i> 2020	J	10.1016/j.ijid.2020.02.052
Wang J, Zhou M, <i>et al.</i> 2020	J	10.1016/j.jhin.2020.03.002
Haines A, de Barros EF, <i>et al.</i> 2020	J	10.1016/S0140-6736(20)30735-2

Al-Tawfiq JA, Auwaerter PG, <i>et al.</i> 2019	J	10.1016/j.jhin.2018.05.021
Park SY, Lee JS, <i>et al.</i> 2019	H	10.1016/j.jhin.2018.09.005
Song Z, Xu Y, <i>et al.</i> 2019	J	10.3390/v11010059
Aldohyan M, Al-Rawashdeh N, <i>et al.</i> 2019	J	10.1186/s12879-019-3898-2
Mobaraki K, Ahmadzadeh J. 2019	J	10.1186/s12879-019-3987-2
Xie J, Hu Y, <i>et al.</i> 2019	J	10.1177/0046958019847865
Corman VM, Lienau J, <i>et al.</i> 2019	I	10.1007/ s00108-019-00671-5
Hong J, Chen J, <i>et al.</i> 2019	J	10.3390/ijerph16193758
Zhao Y, Chen Z, <i>et al.</i> 2019	J	10.1007/s11596-019-2114-3
Wang P, Anderson N, <i>et al.</i> 2020	J	10.1093/clinchem/hvaa080
Li W, Zhou Q, <i>et al.</i> 2020	J	10.21037/apm.2020.02.33
Chen Q, Allot A, <i>et al.</i> 2020	J	10.1038/ d41586-020-00694-1
Klompas M. 2020	J	10.7326/M20-0751
Adams JG, Walls RM. 2020	J	10.1001/jama.2020.3972
Zandifar A, Badrfam R. 2020	E	10.1016/j.ajp.2020.101990
Schwartz J, King CC, <i>et al.</i> 2020	J	10.1093/cid/ciaa255
Kwon KT, Ko JH, <i>et al.</i> 2020	J	10.3346/jkms.2020.35.e123
Li Z, Ge J, <i>et al.</i> 2020	J	10.1016/j.bbi.2020.03.007
Chen C, Zhao B. 2020	J	10.1016/j.jhin.2020.03.008
Lynch C, Mahida N, <i>et al.</i> 2020	J	10.1016/j.jhin.2020.03.010
Rimmer A. 2020	J	10.1136/bmj.m1055
Stein R. 2020	J	10.1111/ijcp.13501
Golda A, Malek N, <i>et al.</i> 2011	E	10.1099/vir.0.028381-0
Scully C, Samaranayake LP. 2016	J	10.1111/odi.12356
Sukumaran A, Patil S	J	25095856
Kharma MY, Alalwani MS, <i>et al.</i> 2015	J	10.4103/2231-0762.159951
Ashok N, Rodrigues JC, <i>et al.</i> 2016	J	10.7860/JCDR/ 2016/17519.7790
Baseer MA, Ansari SH, <i>et al.</i> 2016	J	10.4317/jced.52811
Kim HJ, Ko JS, <i>et al.</i> 2020	J	10.4097/kja.20110

Calvo C, García López-Hortelano M, <i>et al.</i> 2020	A/F/H	10.1016/ j.anpedi.2020.02.001
Mossa-Basha M, Meltzer CC, <i>et al.</i> 2020	J	10.1148/radiol.2020200988
Althomairy SA, Baseer MA, <i>et al.</i> 2018	J	10.4103/ jispcd.JISPCD_9_18
Elrggal ME, Karami NA, <i>et al.</i> 2018	J	10.1007/ s10389-018-0917-5
Gaffar BO, El Tantawi M, <i>et al.</i> 2019	J	10.15537/smj.2019.7.24304
Aminnejad R, Salimi A, <i>et al.</i> 2020	J	10.1007/ s12630-020-01627-2
Oxford JS, Bossuyt S, <i>et al.</i> 2003	J	10.1046/ j.1365-2567.2003.01684.x
Fang LS. 2003	J	14552214
Huber SJ, Wynia MK. 2004	J	10.1162/152651604773067 497
Chang HJ, Huang N, <i>et al.</i> 2004	J	10.2105/ajph.94.4.562
Yu H, Sun X, <i>et al.</i> 2020	J	10.3390/ijerph17051770
Power J. 2004	J	15239585
Li RW, Leung KW, <i>et al.</i> 2004	J	10.1038/sj.bdj.4811469
Testarelli L, D' Aversa L, <i>et al.</i> 2004	J	15278018
Galligan JM. 2004	J	15481238
Adhikari SP, Meng S, <i>et al.</i> 2020	A	10.1186/ s40249-020-00646-x
Yip HK, Tsang PC, <i>et al.</i> 2007	J	17405470
Qian F, Lim MK. 2010	J	10.1007/ s00038-009-0099-6
Holý O, Matoušková I. 2012	J	10.5114/wo.2012.29298
Iacobucci G. 2020	J	10.1136/bmj.m1097
Rimmer A. 2020	J	10.1136/bmj.m1099
Ebrahim SH, Ahmed QA, <i>et al.</i> 2020	J	10.1136/bmj.m1066
Kain NA, Jardine CG. 2020	J	32165480
Silva AAMD. 2020	J	10.1590/1980-54972020002 1.
De Ceukelaire W, Bodini C. 2020	J	10.1177/002073142091672 5.
Wang H, Wang S, <i>et al.</i> 2020	J	10.1186/ s13054-020-2832-8

Rimmer A. 2020	J	10.1136/bmj.m1104
Tanne JH, Hayasaki E, <i>et al.</i> 2020	J	10.1136/bmj.m1090
Thomas-Rüddel D, Winning J, <i>et al.</i> 2020	I	10.1007/ s00101-020-00758-x
Giovanetti M, Angeletti S, <i>et al.</i> 2020	J	10.1002/jmv.25773
Koenig KL, Bej CK, <i>et al.</i> 2020	J	10.5811/ westjem.2020.1.46760.
Bachireddy C, Chen C, <i>et al.</i> 2020	J	10.1001/jama.2020.4272.
Lippi G, Plebani M. 2020	J	10.1515/cclm-2020-0240.
Kakimoto K, Kamiya H, <i>et al.</i> 2020	J	10.15585/ mmwr.mm6911e2.
Ng Y, Li Z, <i>et al.</i> 2020	J	10.15585/ mmwr.mm6911e1.
Salathé M, Althaus CL, <i>et al.</i> 2020	J	10.4414/smw.2020.20225
Cleveland J, Gray ST, <i>et al.</i> 2020	E	10.1016/j.ada.2016.03.020
Gorbalenya AE. 2020	J	10.1101/2020.02.07.937862
To K, Tsang O <i>et al.</i> 2020	J	10.1093/cid/ciaa149.
Kwok YLL, Gralton J, <i>et al.</i> 2020	J	10.1016/j.ajic.2014.10.015
Wahba L, Jain N <i>et al.</i> 2020	B	10.1093/cid/ciaa149
Schwartz IS, Friedman DZP, <i>et al.</i> 2020	E	10.1093/cid/ciaa007
Crooks K, Massey PD, <i>et al.</i> 2018	J	10.5365/wpsar.2018.9.5.005
Singh I, Munjal S, <i>et al.</i> 2019	J	10.4103/ jfmprc.jfmprc_381_19
Patel AB, Verma A. 2020	D/H	10.1001/jama.2020.4812
Yamamoto Y, Saruta J, <i>et al.</i> 2019	H	10.1080/00016357.2019.16 09697
Fessell D, Cherniss C, <i>et al.</i> 2020	J	10.1016/j.jacr.2020.03.013
Jairaj A, Shirisha P, <i>et al.</i> 2018	C	10.4103/ jispdc.JISPDC_347_18
Kumar R, Kratzer D, <i>et al.</i> 2019	J	10.1002/marc.201800530
Gagliano A, Villani PG, Cò FM, <i>et al.</i> 2020	F	10.1017/dmp.2020.51
Roode GJ, Bütow KW. 2018	E	10.3121/cmr.2018.1385

Nicol GE, Piccirillo JF, <i>et al.</i> 2020	J	10.1111/jgs.16443
McKimm-Breschkin JL, Jiang S, <i>et al.</i> 2018	C/D/H	10.1016/j.antiviral.2017.11.013
Naleway AL, Henninger ML, <i>et al.</i> 2018	C	10.1111/jphd.12256
Jiang S, Ojcius DM. 2017	C/D	10.1016/j.micinf.2017.09.013
Zhou Y, Jiang S, <i>et al.</i> 2018	E	10.1111/idj.12335
Sperr M, Kundi M, <i>et al.</i> 2018	E	10.1902/jop.2017.170333
Gambhir RS, Pannu PR, <i>et al.</i> 2016	J	10.7860/JCDR/2016/20387.8536
Tsuruhara A, Aso K, <i>et al.</i> 2017	J	10.1093/intimm/dxx001
Bulgakova VA, Poromov AA, <i>et al.</i> 2017	I	10.17116/terarkh201789162-71
Mishra P, Bhadauria US. 2016	J	27344473
Cusinato R, Pacenti M, <i>et al.</i> 2016	J	10.1016/j.jviromet.2016.03.001
Buckee CO, Balsari S, <i>et al.</i> 2020	J	10.1126/science.abb8021
Lillemoe J, Caplan AL. 2016	J	30512265
Freund Y. 2020	J	10.1097/MEJ.0000000000000699
Li H, Zhou Y, <i>et al.</i> 2020	H	10.1128/AAC.00483-20
Joob B, Wiwanitkit V. 2020	J	10.1016/j.bbi.2020.03.020
Li S, Wang Y, Xue J, <i>et al.</i> 2020	J	10.3390/ijerph17062032
Bouckaert N, Schokkaert E. 2016	J	10.1007/s10198-015-0709-6
Feng S, Shen C, <i>et al.</i> 2020	J	10.1016/S2213-2600(20)30134-X
Pylińska-Dąbrowska D, Starzyńska A, <i>et al.</i> 2020	G	10.3390/jcm9103344
Sebastian P, Jorge P, <i>et al.</i> 2021	G	10.1016/j.jobcr.2021.01.006
Jungo S, Moreau N, <i>et al.</i> 2021	A	10.1371/journal.pone.0246586
Estrich CG, Mikkelsen M, <i>et al.</i> 2020	A	10.1016/j.adaj.2020.09.005
Allen-Tejerina AM, Rallis KS, <i>et al.</i> 2020	B	10.1016/j.eururo.2020.10.003
Yee R, Truong TT, <i>et al.</i> 2021	G	10.1128/JCM.02686-20
Costa Dos Santos Junior G, Pereira CM, <i>et al.</i> 2020	D	10.1021/acs.analchem.0c04679



Carraturo F, Del Giudice C, <i>et al.</i> 2020	B	10.1016/ j.envpol.2020.115010
Yang H, Hu J, <i>et al.</i> 2020	G	10.1016/ j.pdpdt.2020.101943
Barbato L, Bernardelli F, <i>et al.</i> 2020	G	10.1111/odi.13646
Vargas-Buratovic JP, Verdugo-Paiva F, <i>et al.</i> 2020	G	10.5867/ medwave.2020.05.7916.
Ashtiani RE, Tehrani S, <i>et al.</i> 2020	G	10.1111/jopr.13261
Shi AH, Guo W, <i>et al.</i> 2020	G	PMID: 32582908
Hertrampf K, Flörke C, <i>et al.</i> 2020	G	10.1111/odi.13529
Zhao B, An N, <i>et al.</i> 2020	G	10.1017/ice.2020.29
Rupf S, Hannig M. 2020	G	10.1007/ s00784-020-03351-z
Harmooshi NN, Shirbandi K, <i>et al.</i> 2020	G	10.1007/ s11356-020-09733-w
Khamis AH, Jaber M, <i>et al.</i> 2020	D	10.1016/j.jfma.2020.12.003
Abduljabbar T, Alhamdan RS, <i>et al.</i> 2020	D	10.1055/s-0040-1716986
Guo F, Tang B, <i>et al.</i> 2020	G	10.3389/fmed.2020.577468
Esteki R, Asgari N, <i>et al.</i> 2020	D	<a href="https://doi.org/10.29089/2020.20.00115">https://doi.org/ 10.29089/2020.20.00115</a>
Xiong X, Wu Y, <i>et al.</i> 2020	G	10.1016/ j.ajodo.2020.07.005
Cueno ME, Imai K. 2021	D	10.3389/fmed.2020.594439
Davies A, Howells R, <i>et al.</i> 2020	G	10.1111/ipd.12773
Saltaji H, Sharaf KA. 2020	G	10.1016/ j.ajodo.2020.04.006
Misra RS, Johnston CJ, <i>et al.</i> 2015	J	10.1667/RR13917.1
Lai J, Ma S, <i>et al.</i> 2020	J	10.1001/ jamanetworkopen.2020.397 6
Dong L, Bouey J. 2020	J	10.3201/eid2607.200407
Perlis RH. 2020	J	10.1001/ jamanetworkopen.2020.4006
Broadbent L, Groves H, <i>et al.</i> 2015	C/E/H	10.1111/irv.12313
Ficarra V, Novara G, <i>et al.</i> 2020	E	10.23736/ S0393-2249.20.03846-1
Choi KR, Skrine Jeffers K, <i>et al.</i> 2020	F	10.1111/jan.14369
Agarwal A, Nagi N, <i>et al.</i> 2020	J	10.4103/ijmr.IJMR_518_20

Colvard MD, Hirst JL, <i>et al.</i> 2014	C	10.1017/dmp.2014.44
Jin GB, Winans B, <i>et al.</i> 2014	J	10.1002/eji.201343980
Jerant A, Fiscella K, <i>et al.</i> 2013	J	10.3122/jabfm.2013.06.130054
Tjäderhane L, Koivumäki S, <i>et al.</i> 2013	J	10.1177/0022034513504783
Zucker J, Lee J, <i>et al.</i> 2013	J	10.1353/hpu.2013.0009
Wheeler JL, Martin KC, <i>et al.</i> 2013	J	10.4049/jimmunol.1201341
Mahanonda R, Sa-Ard-lam N, <i>et al.</i> 2012	J	10.1002/eji.201141657
The Lancet. 2020	J	10.1016/S0140-6736(20)30644-9
Lima CKT, Carvalho PMM, <i>et al.</i> 2020	J	10.1016/j.psychres.2020.112915
Pang HB, Xu LM, <i>et al.</i> 2020	J	10.12116/j.issn.1004-5619.2020.01.007
Qiu H, Wang HJ, <i>et al.</i> 2020	J	10.12116/j.issn.1004-5619.2020.01.006
Mao DM, Zhou N, <i>et al.</i> 2020	J	10.12116/j.issn.1004-5619.2020.01.003
Willan J, King AJ, <i>et al.</i> 2020	J	10.1136/bmj.m1117
Wicker S, Rabenau HF, <i>et al.</i> 2011	C	10.1016/j.ijheh.2011.08.005
Pyrk K, Strzyz P, <i>et al.</i> 201	J	10.1099/vir.0.032094-0
Bilder L, Machtei EE, <i>et al.</i> 2011	A	10.1177/0022034511413283
Malamud D, Abrams WR, <i>et al.</i> 2011	J	10.1177/0022034511399282
Flaer PJ, Younis MZ, <i>et al.</i> 2010	J	20973377
Kamate SK, Agrawal A, <i>et al.</i> 2009	J	20130372
Molinari JA. 2010	J	20333864
Mays JW, Bailey MT, <i>et al.</i> 2010	J	10.4049/jimmunol.0900183
Merchant VA. 2009	J	19877489
Dewey C, Hingle S, <i>et al.</i> 2020	J	10.7326/M20-1033
Li P, Fu JB, <i>et al.</i> 2020	D	10.1016/j.ijid.2020.03.027
Rueckmann E, Shah MN, <i>et al.</i> 2009	C	10.1080/10903120802471949

McNeary L, Maltser S, <i>et al.</i> 2020	J	10.1002/pmrj.12369
Chen D, Yang H, <i>et al.</i> 2020	E	10.1002/ijgo.13146
Luo M, Cao S, <i>et al.</i> 2020	J	10.1097/ ALN.00000000000003288
Zhang HF, Bo LL, <i>et al.</i> 2020	J	10.1097/ ALN.00000000000003300
Zhu W, Wang Y, <i>et al.</i> 2020	J	10.1097/ ALN.00000000000003299
Chen X, Liu Y, <i>et al.</i> 2020	J	10.1097/ ALN.00000000000003301
Greenland JR, Michelow MD, <i>et al.</i> 2020	J	10.1097/ ALN.00000000000003303
Owens MD, Beckles GL, <i>et al.</i> 2008	E	10.1089/jwh.2008.1125
Banerjea R, Findley PA, <i>et al.</i> 2008	J	10.1080/03630240802099261
Fiscella K, Holt K. 2007	J	10.3122/jabfm.2007.06.070053
Fiscella K, Holt K, <i>et al.</i> 2006	J	10.1186/1472-6963-6-122
Shah MN, Clarkson L, <i>et al.</i> 2006	J	10.1111/ j.1532-5415.2006.00736.x
Oxford JS, Lambkin R. 2006	E/H	10.1016/ j.ijantimicag.2006.02.007
Szilagyi PG, Shone LP, <i>et al.</i> 2005	J	10.1016/j.ypped.2004.05.010
Marin MG, Zitter JN. 2004	J	10.1016/j.ypped.2004.03.017
Falsey AR, Walsh EE, <i>et al.</i> 2002	A	10.1086/339881
Shuman SK, McCusker ML, <i>et al.</i> 1993	E	10.14219/ jada.archive.1993.0177
Fernandez JB, Glotzer DL, <i>et al.</i> 2008	J	10.5055/ajdm.2008.0019
Osterholm MT. 2006	J	16573182
Wessner M. 2005	J	16379153
Palenik CJ, Govoni M. 2004	J	15641330
McClarey M. 2000	J	12026448
Wang Y, Zhou CC, <i>et al.</i> 2020	I	10.12182/20200360101
Bol P. 2001	I	11441718
Janes GR, Blackman DK, <i>et al.</i> 199	J	10634271
Vingilis ER, Brown U, <i>et al.</i> 1999	J	PMC6979913

Carter WB, 1992	J	1487582
Ship JA. 1992	J	1504940
Palenik CJ, Miller CH. 1990	J	2284556
Oehring H, Welker D, <i>et al.</i> 1990	J	2150255
Oehring H, Welker D, <i>et al.</i> 1990	J	2149903
Dolin R. 1990	C	PMC2376495
Weintraub TL, Iftimovici R. 1981	J	7245605
Waldman HB. 1977	J	275493
Constantin I, Severineanu V, <i>et al.</i> 1977	I	266299
Lang W. 1973	H	4582161
KALMI. 1952	J	13048973
Alwan NA, Bhopal R, <i>et al.</i> 2020	J	10.1016/S0140-6736(20)30667-X
Pan X, Ojcius D, <i>et al.</i> 2020	D	10.1016/j.micinf.2020.02.004
Liu K. 2020	J	10.1126/science.367.6484.1398
Cohen J, Kupferschmidt K. 2020	D	10.1126/science.367.6484.1287
Wang J, Xu C, <i>et al.</i> 2020	D/E	10.1016/S0140-6736(20)30561-4
Pung R, Chiew CJ, <i>et al.</i> 2020	D	10.1016/S0140-6736(20)30528-6
Morishima T. 2006	J	17037365
Hartmann-Boyce J, Mahtani K. 2020	J	<a href="https://www.cebm.net/COVID-19/supporting-people-with-long-term-conditions-ltcs-during-national-emergencies/">https://www.cebm.net/COVID-19/supporting-people-with-long-term-conditions-ltcs-during-national-emergencies/</a>
Health Protection Scotland 2020	J	<a href="https://www.hps.scot.nhs.uk/web-resources/container/COVID-19-guidance-for-first-responders-and-others-in-close-contact-with-symptomatic-people-with-potential-COVID-19/">https://www.hps.scot.nhs.uk/web-resources/container/COVID-19-guidance-for-first-responders-and-others-in-close-contact-with-symptomatic-people-with-potential-COVID-19/</a>
Intensive Care medicine 2020	J	<a href="https://static1.squarespace.com/static/5e6613a1dc75b87df82b78e1/t/5e6d0530d0babc2b370bf29b/1584203289238/COVID-19_Airwayshort.pdf">https://static1.squarespace.com/static/5e6613a1dc75b87df82b78e1/t/5e6d0530d0babc2b370bf29b/1584203289238/COVID-19_Airwayshort.pdf</a>
Hartmann-Boyce J, Narthan D, <i>et al.</i> 2020	J	<a href="https://www.cebm.net/COVID-19/maximising-mobility-in-the-older-people-when-isolated-with-COVID-19/">https://www.cebm.net/COVID-19/maximising-mobility-in-the-older-people-when-isolated-with-COVID-19/</a>

Health Protection Scotland 2020	J	<a href="https://www.hps.scot.nhs.uk/web-resources-container/COVID-19-guidance-for-infection-prevention-and-control-in-healthcare-settings/">https://www.hps.scot.nhs.uk/web-resources-container/COVID-19-guidance-for-infection-prevention-and-control-in-healthcare-settings/</a>
Health Protection Scotland 2020	J	<a href="https://www.hps.scot.nhs.uk/web-resources-container/COVID-19-guidance-for-secondary-care/">https://www.hps.scot.nhs.uk/web-resources-container/COVID-19-guidance-for-secondary-care/</a>
Health Protection Scotland 2020	A	<a href="https://hpspubsrepo.blob.core.windows.net/hps-website/nss/2935/documents/1_COVID-19-guidance-for-hpt.pdf">https://hpspubsrepo.blob.core.windows.net/hps-website/nss/2935/documents/1_COVID-19-guidance-for-hpt.pdf</a>
Khunti K, Greenhalgh T <i>et al.</i> 2020	J	<a href="https://www.cebm.net/COVID-19/what-is-the-efficacy-of-eye-protection-equipment-compared-to-no-eye-protection-equipment-in-preventing-transmission-of-COVID-19-type-respiratory-illnesses-in-primary-and-community-care/">https://www.cebm.net/COVID-19/what-is-the-efficacy-of-eye-protection-equipment-compared-to-no-eye-protection-equipment-in-preventing-transmission-of-COVID-19-type-respiratory-illnesses-in-primary-and-community-care/</a>
INTERIM ClínicaL GUIDANCE FOR ADULTS WITH SUSPECTED OR CONFIRMED COVID-19 IN BELGIUM Abril 2020	F	<a href="https://epidemiology.wiv-isp.be/ID/Documents/Covid19/COVID-19_InterimGuidelines_Treatment_ENG.pdf">https://epidemiology.wiv-isp.be/ID/Documents/Covid19/COVID-19_InterimGuidelines_Treatment_ENG.pdf</a>
Ministry Of health ontario 2020	J	<a href="http://www.health.gov.on.ca/en/pro/programs/publichealth/coronavirus/docs/2019_primary_care_guidance.pdf">http://www.health.gov.on.ca/en/pro/programs/publichealth/coronavirus/docs/2019_primary_care_guidance.pdf</a>
University of Oxford 2020	J	<a href="https://bjglife.com/wp-content/uploads/2020/03/Video-consultations-a-guide-for-practice.pdf">https://bjglife.com/wp-content/uploads/2020/03/Video-consultations-a-guide-for-practice.pdf</a>
He Knowledge and Library Services Team 2020	J	<a href="https://phelibrary.koha-ptfs.co.uk/coronavirusinformation/">https://phelibrary.koha-ptfs.co.uk/coronavirusinformation/</a>

## Section 5 | Level of Evidence and Classes of Recommendation

*Evidence Levels Adapted and translated from Perry, DJ, et al. 2007<sup>[184]</sup>*

Level of Evidence	Descriptive
<b>Level Ia</b>	Evidence from systematic reviews or meta-analyses of randomized controlled trials
<b>Level Ib</b>	Evidence from, at least, one randomized controlled trial
<b>Level IIa</b>	Evidence from, at least, one well-designed non-randomized controlled trial
<b>Level IIb</b>	Evidence from, at least, one other type of quasi-experimental well-designed trial
<b>Level III</b>	Evidence from descriptive non-experimental well-designed trials, such as comparative studies, correlation studies and case studies
<b>Level IV</b>	Evidence from reports of expert committees or experiments performed by respected entities

*Classes of recommendation Adapted and translated from Kirchhof P, et al. 2016<sup>[185]</sup>*


Class of recommendation	Descriptive	Indication
<b>Class I</b>	Evidence and/or general agreement that a given procedure or treatment is beneficial, useful, and effective	<b>Recommended/Indicated</b>
<b>Class II</b>	Conflicting evidence and/or a divergence of opinion about the usefulness/efficacy of a procedure or treatment	
<b>Class IIa</b>	Evidence/opinion is in favor of usefulness/efficacy	<b>Should be considered</b>
<b>Class IIb</b>	Usefulness/efficacy is less well established by evidence/opinion	<b>May be considered</b>
<b>Class III</b>	Evidence and/or general agreement that a procedure/treatment is not useful/effective and in some cases may be harmful	<b>Not recommended</b>

## Appendix III

Support Material



Section 1 | Hand hygiene technique using an alcohol-base antiseptic solution (ABAS) or water and soap<sup>[84]</sup>



## Section 2 | Sequence to put on and remove PPE

### Put on PPE

Protection of degree 1 (for moderate risk level)	Protection of degree 2 (for high risk level)	Protection of degree 3 (for very high risk level)
<p>1st Hand antisepsis;</p> <p>2nd Adapt surgical mask and check if it is adjusted;</p> <p>3rd Put on impervious gown;</p> <p>4th Put on surgical cap or bouffant cap;</p> <p>5th Hand antisepsis with an ABAS;</p> <p>6th Put on well adapted gloves covering the gown.</p>	<p>1st Hand antisepsis;</p> <p>2nd Adapt respirator and perform adjustment test;</p> <p>3rd Put on impervious gown;</p> <p>4th Put on disposable shoe-covers;</p> <p>5th Hand antisepsis with an ABAS;</p> <p>6th Put on goggles (if applicable);</p> <p>7th Put on a surgical cap or surgical hood;</p> <p>8th Put on facial shield (if applicable);</p> <p>9th Put on well adapted gloves covering the gown;</p> <p>10th Check if all the equipment is well adjusted.</p>	<p>1st Hand antisepsis;</p> <p>2nd Adapt respirator and perform adjustment test;</p> <p>3rd Put on impervious gown or impervious coveralls;</p> <p>4th Put on disposable shoe-covers;</p> <p>5th Hand antisepsis with an ABAS;</p> <p>6th Put on a pair of gloves covering the gown/coverall;</p> <p>7th Put on goggles (if applicable);</p> <p>8th Put on hood of coverall or surgical gown;</p> <p>9th Put on facial shield;</p> <p>10th Disinfect inner gloves with ABAS;</p> <p>11th Remove outer pair of gloves;</p> <p>12th Check if all the equipment is well adjusted.</p>

## Removal of PPE

Protection of degree 1 (for moderate risk level)	Protection of degree 2 (for high risk level)	Protection of degree 3 (for very high risk level)
<p>1st Remove impervious gown;</p> <p>2nd Remove gloves together with the gown;</p> <p>3rd Hand antisepsis with an ABAS;</p> <p>4th Remove surgical cap or bouffant cap;</p> <p>5th Remove surgical mask;</p> <p>6th Hand antisepsis with an ABAS;</p>	<p>1st Remove impervious gown;</p> <p>2nd Remove gloves together with the gown;</p> <p>3rd Hand antisepsis with an ABAS;</p> <p>4th Remove eye protection and cap/bouffant (from the back to the front);</p> <p>5th Remove all shoe-covers:</p> <p>6th Hand antisepsis with an ABAS;</p> <p>7th Remove respirator;</p> <p>8th Hand antisepsis with an ABAS;</p>	<p>1st Remove outer pair of gloves;</p> <p>2nd Disinfect inner gloves with ABAS;</p> <p>3th Remove eye protection and cap/bouffant (from the back to the front) or the hood of the coverall;</p> <p>4th Remove the gown and simultaneously the inner pair of gloves; or remove the coverall and then the gloves;</p> <p>5th Remove all shoe-covers:</p> <p>6th Hand antisepsis with an ABAS;</p> <p>7th Remove respirator;</p> <p>8th Hand antisepsis with an ABAS;</p>

## Section 3 | *Disinfectant Solutions*

### **70% Ethanol Solution**

To obtain a 70% ethanol solution based on a 96% ethanol stock-solution, you need:

729,2 mL of ethanol at 96%

270,8 mL distilled water

### **0.1% Sodium Hypochlorite Solution**

To obtain 1 L of 0.1% hypochlorite solution, based on a 5% hypochlorite solution stock-solution, you need:

20 mL of Sodium Hypochlorite at 5%

980 mL distilled water

### **0.5% Sodium Hypochlorite Solution**

To obtain 1 L of 0.5% hypochlorite solution, based on a 5% hypochlorite solution stock-solution, you need:

100 mL of Sodium Hypochlorite at 5%

900 mL distilled water

### **1% Sodium Hypochlorite Solution**

To obtain 1 L of 1% hypochlorite solution, based on a 5% hypochlorite solution stock-solution, you need:

200 mL of Sodium Hypochlorite at 5%

800 mL distilled water

## **0.2% Povidone-iodine Solution**

To obtain 1L of 0.2% povidone-iodine solution based on a 10% povidone-iodine stock-solution, you need:

20 mL of 10% povidone-iodine

980 mL distilled water

## **1% Hydrogen Peroxide Solution**

To obtain 1 L of 1% hydrogen peroxide solution, based on a 3% hydrogen peroxide solution stock-solution (equivalent to 10 volume Oxygenated Water), you need:

333.3 mL of 3% hydrogen peroxide

666,7 mL distilled water

## **1% Hydrogen Peroxide Solution**

To obtain 1 L of 1% hydrogen peroxide solution, based on a 9% hydrogen peroxide solution stock-solution (equivalent to 30 volume Oxygenated Water), you need:

111 mL of 9% hydrogen peroxide

889 mL distilled water

## Bibliography

1. Guerra, F., et al., COVID-19. Normas de Orientação Clínica - Medicina Dentária. Universidade de Coimbra, 2020. ISBN: 978-989-26-1985-9. DOI: 10.14195/978-26-1985-9.
2. National Institute for Health and Care Excellence, Developing NICE guidelines: the manual Process and methods. Published: 31 October 2014. Last updated: 15 October 2020. <https://www.nice.org.uk/process/pmg20/chapter/updating-guidelines> (acedido a 10 de março de 2021).
3. Rosenfeld, R.M., et al., Clinical practice guideline development manual: A quality-driven approach for translating evidence into action. Otolaryngology–Head and Neck Surgery, 2009. 140:S1-S43. DOI: 10.1016/j.otohns.2009.04.015.
4. National Institute for Health and care Excellence, Interim process and methods for developing rapid guidelines on COVID-19. Published date: 20 March 2020. <https://www.nice.org.uk/process/pmg35/chapter/scoping> (acedido a 11 de março de 2021).
5. El Zowalaty, M.E. and Jarhult, J.D., From SARS to COVID-19: A previously unknown SARS- related coronavirus (SARS-CoV-2) of pandemic potential infecting humans - Call for a One Health approach. One Health, 2020. 9: p. 100124.
6. Xu, J., *et al.*, 2019 novel Coronavirus outbreak: a quiz or final exam? Front Med, 2020.
7. Sohrabi, C., *et al.*, World Health Organization declares global emergency: A review of the 2019 novel coronavirus (COVID-19). Int J Surg, 2020. 76: p. 71-76.
8. Amodio, E., *et al.*, Outbreak of Novel Coronavirus (SARS-Cov-2): First Evidences From International Scientific Literature and Pending Questions. Healthcare (Basel), 2020. 8(1).
9. Chan, K.W., *et al.*, COVID-19: An Update on the Epidemiological, Clinical, Preventive and Therapeutic Evidence and Guidelines of Integrative Chinese-Western Medicine for the Management of 2019 Novel Coronavirus Disease. Am J Chin Med, 2020: p. 1-26.
10. Coronaviridae Study Group of the International Committee on Taxonomy of Viruses, The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. Nat Microbiol, 2020. 5(4): p. 536-544.
11. Guan, W.J., *et al.*, Clinical Characteristics of Coronavirus Disease 2019 in China. N Engl J Med, 2020.

12. Spagnuolo, G., et al., COVID-19 Outbreak: An Overview on Dentistry. *Int J Environ Res Public Health*, 2020. 17(6):2094. DOI: 10.3390/ijerph17062094.
13. Ungaro, R.C., et al., What Should Gastroenterologists and Patients Know About COVID-19? *Clin Gastroenterol Hepatol*, 2020.
14. Borges do Nascimento, I.J., et al., Novel Coronavirus Infection (COVID-19) in Humans: A Scoping Review and Meta-Analysis. *J Clin Med*, 2020. 9(4).
15. World Health Organization, Clinical management of severe acute respiratory infection when COVID-19 is suspected. [https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected). (acedido a 20 de abril de 2020).
16. Xia, N., et al., Serological Test is an Efficient Supplement of RNA Detection for Confirmation of SARS-CoV-2 Infection. *Preprints 2020*, 2020030184 (doi: 10.20944/preprints202003.0184.v1).
17. Meng, L., et al., Coronavirus Disease 2019 (COVID-19): Emerging and Future Challenges for Dental and Oral Medicine. *J Dent Res*, 2020. 99(5). DOI: <https://doi.org/10.1177/0022034520914246>.
18. Sahni, V. and Gupta, S., COVID-19 & Periodontitis: The cytokine connection. *Medical Hypotheses*, 2020. 144. DOI: <https://doi.org/10.1016/j.mehy.2020.109908>.
19. Takahashi, Y., et al., Aspiration of periodontopathic bacteria due to poor oral hygiene potentially contributes to the aggravation of COVID-19. *J Oral Sci*, 2020. 63(3):1-3. DOI: 10.2334/josnusd.20-0388.
20. Manzalawi, R., et al., Gingival bleeding associated with COVID-19 infection. *Clinical Case Reports*, 2020. DOI: <https://doi.org/10.1002/ccr3.3519>.
21. Di Spirito, F., et al., Oral manifestations and the role of the oral healthcare workers in COVID-19. *Oral Diseases*, 2020. DOI: <https://doi.org/10.1111/odi.13688>.
22. Badran, Z., et al., Periodontal pockets: A potential reservoir for SARS-CoV-2?. *Medical Hypotheses*, 2020. 143. DOI: <https://doi.org/10.1016/j.mehy.2020.109907>.
23. Xiang, Z., et al., Potential implications of SARS-CoV-2 oral infection in the host microbiota. *J Oral Microbiol*, 2020. 13(1):1853451. DOI: 10.1080/20002297.2020.1853451.
24. Amorim dos Santos, J., et al., Oral Manifestations in Patients with COVID-19: A Living Systematic Review. *Journal of Dental Research*, 2020. DOI: <https://doi.org/10.1177/0022034520957289>.
25. Bemquerer, L.M., et al., The oral cavity cannot be forgotten in the COVID-19 era: Is there a connection between dermatologic and oral manifestations? *Journal of the*

American Academy of Dermatology, 2021. 84:E143-E145. DOI: <https://doi.org/10.1016/j.jaad.2020.11.034>.

26. Corchuelo, J. and Ullo, F.C., et al., Oral manifestations in a patient with a history of asymptomatic COVID-19: Case report. *International Journal of Infectious Diseases*, 2020. 100:154-157. DOI: <https://doi.org/10.1016/j.ijid.2020.08.071>.

27. Brandão, T.B., et al., Oral lesions in patients with SARS-CoV-2 infection: could the oral cavity be a target organ? *Oral Surgery Oral Medicina Oral Pathology Oral Radiology*, 2021. 131:E45-E51. DOI: <https://doi.org/10.1016/j.oooo.2020.07.014>.

28. Riad, A., et al., The manifestation of oral mucositis in COVID-19 patients: A case-series. *Dermatol Ther*, 2020. 33(6):e14479. DOI: 10.1111/dth.14479.

29. Pérez-Sayáns, M., et al., Can “COVID-19 tongue” be considered a pathognomonic finding in SARS-CoV-2 infection?. *Oral Diseases*, 2021. DOI: <https://doi.org/10.1111/odi.13807>.

30. Hathway, R.W., COVID tongue. *Br Dent J*, 2021. 230(3):114. DOI: 10.1038/s41415-021-2666-z.

31. Santosh, A.B.R., et al., Asymptomatic COVID-19 and saliva: Let's ask "Do you feel that saliva in your mouth had reduced in recent times?". *Int J Clin Pract*, 2020. 74(12):e13657. DOI: 10.1111/IJCP.13657.

32. Sarode, G.S., et al., Clinical status determines the efficacy of salivary and nasopharyngeal samples for detection of SARS-CoV-2. *Clinical Oral Investigations*, 2020. 24:4661-4662. DOI: <https://doi.org/10.1007/s00784-020-03630-9>.

33. Medeiros da Silva, R.C., et al., Saliva as a possible tool for the SARS-CoV-2 detection: A review. *Travel Medicine and Infectious Disease*, 2020. 38. DOI: <https://doi.org/10.1016/j.tmaid.2020.101920>.

34. Fernandes, L.L., et al., Saliva in the Diagnosis of COVID-19: A Review and New Research Directions. *J Dent Res*, 2020. 99(13):1435-1443. DOI: 10.1177/0022034520960070.

35. Michailidou, E., et al., Salivary diagnostics of the novel coronavirus SARS-CoV-2 (COVID-19). *Oral Diseases*, 2020. [Epub ahead of print]. DOI: 10.1111/odi.13729.

36. Kashiwagi, K., et al., Immunochromatographic test for the detection of SARS-CoV-2 in saliva. *Journal of Infection and Chemotherapy*, 2021. 27(2):384-386. DOI: <https://doi.org/10.1016/j.jiac.2020.11.016>.

37. Hamid, H., et al., COVID-19 Pandemic and Role of Human Saliva as a Testing Biofluid in Point-of-Care Technology. *Eur J Dent*, 2020. 14(1):S123-S129. DOI: 10.1055/s-0040-1713020.



38. Chen, L., et al., Detection of SARS-CoV-2 in saliva and characterization of oral symptoms in COVID-19 patients. *Cell Prolif*, 2020. 53(12):e12923. DOI: 10.1111/cpr.12923.
39. Hung, K., et al., New COVID-19 saliva-based test: How good is it compared with the current nasopharyngeal or throat swab test? *J Chin Med Assoc*, 2020. 83(10):891-894. DOI: 10.1097/JCMA.0000000000000396.
40. Azzi, L., et al., Diagnostic Salivary Tests for SARS-CoV-2. *J Dent Res*, 2021. 100(2):115-123. DOI: 10.1177/0022034520969670.
41. Fakheran, O., et al., Saliva as a diagnostic specimen for detection of SARS-CoV-2 in suspected patients: a scoping review. *Infect Dis Poverty*, 2020. 9(1):100. DOI: <https://doi.org/10.1186/s40249-020-00728-w>.
42. Altawalrah, H., et al., Saliva specimens for detection of severe acute respiratory syndrome coronavirus 2 in Kuwait: A cross-sectional study. *J Clin Virol*, 2020. 132:104652. DOI: <https://doi.org/10.1016/j.jcv.2020.104652>.
43. Czumbel, L.M., et al., Saliva as a Candidate for COVID-19 Diagnostic Testing: A Meta-Analysis. *Front Med (Lausanne)*, 2020. 7:465. DOI: 10.3389/fmed.2020.00465.
44. Gupta, S., et al., SARS-CoV-2 Detection in Gingival Crevicular Fluid. *J Dent Res*, 2021. 100(2):187-193. DOI: 10.1177/0022034520970536.
45. Droegemeier, K., Rapid Expert Consultation Update on SARS-CoV-2 Surface Stability and Incubation for the COVID-19 Pandemic. National Academies of Sciences, Engineering, and Medicine, 2020.
46. Bai, Y., *et al.*, Presumed Asymptomatic Carrier Transmission of COVID-19. *JAMA*, 2020.
47. Wang, J., *et al.*, Preparedness is essential for malaria-endemic regions during the COVID-19 pandemic. *Lancet*, 2020. 395(10230): p. 1094-1096.
48. Zou, L., *et al.*, SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. *N Engl J Med*, 2020. 382(12): p. 1177-1179.
49. National Research Council 2020. Rapid Expert Consultation on the Possibility of Bioaerosol Spread of SARS-CoV-2 for the COVID-19 Pandemic (April 1, 2020). Washington, DC: The National Academies Press. <https://doi.org/10.17226/25769>.
50. Protocol for prevention and control of COVID-19, CDC - China 2020. <http://www.nhc.gov.cn/jkj/s3577/202003/4856d5b0458141fa9f376853224d41d7.shtml>. (acedido a 17 de abril de 2020).
51. Huang, C., *et al.*, Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*, 2020. 395(10223): p. 497-506.

52. Santarpia, J.L., *et al.*, Transmission Potential of SARS-CoV-2 in Viral Shedding Observed at the University of Nebraska Medical Center. medRxiv, 2020.
53. Li, J.Y., *et al.*, The epidemic of 2019-novel-coronavirus (2019-nCoV) pneumonia and insights for emerging infectious diseases in the future. *Microbes Infect*, 2020. 22(2): p. 80-85.
54. Peng, X., *et al.*, Transmission routes of 2019-nCoV and controls in dental practice. *Int J Oral Sci*, 2020. 12(1):9. DOI: 10.1038/s41368-020-0075-9.
55. Zhu, F., *et al.*, Safety, tolerability, and immunogenicity of a recombinant adenovirus type-5 vectored COVID-19 vaccine: a dose-escalation, open-label, non-randomised, first-in-human trial. *Lancet*, 2020. 395(10240):1845-1854. DOI: 10.1016/S0140-6736(20)31208-3.
56. Zhao, J., *et al.*, COVID-19: Coronavirus Vaccine Development Updates. *Front Immunol*, 2020. 11: 602256. DOI: 10.3389/fimmu.2020.602256.
57. Awadasseid, A., *et al.*, Current advances in the development of SARS-CoV-2 vaccines. *Int J Biol Sci*, 2021. 17(1): 8–19. DOI: 10.7150/ijbs.52569.
58. Kim, K.D., *et al.*, Progress and Challenges in the Development of COVID-19 Vaccines and Current Understanding of SARS-CoV-2- Specific Immune Responses. *J Microbiol Biotechnol*, 2020. 30(8):1109-1115. doi: 10.4014/jmb.2006.06006.
59. Norma 021/2020: Campanha de Vacinação contra a COVID-19 Vacina COMIRNATY®. Data: 23/12/2020. Atualizada a 01/03/2021. (<https://covid19.min-saude.pt/wp-content/uploads/2021/03/i027220.pdf>)
60. Norma 001/2021: Campanha de Vacinação contra a COVID-19 COVID-19 Vaccine MODERNA®. Data: 14/01/2021. (<https://covid19.min-saude.pt/wp-content/uploads/2021/01/norma-01-2021-14012021.pdf>)
61. Norma 003/2021: Campanha de Vacinação contra a COVID-19 COVID-19 Vaccine AstraZeneca®. Data: 08/02/2021. Atualizada a 10/03/2021. (<https://covid19.min-saude.pt/wp-content/uploads/2021/03/i027257.pdf>).
62. Centers for Disease Control and Prevention. Interim Clinical Considerations for Use of COVID-19 Vaccines Currently Authorized in the United States. Last updated March 5, 2021. <https://www.cdc.gov/vaccines/covid-19/info-by-product/clinical-considerations.html> (acedido a 11 de março de 2021).
63. Bernal, J.L., *et al.*, Early effectiveness of COVID-19 vaccination with BNT162b2 mRNA vaccine and ChAdOx1 adenovirus vector vaccine on symptomatic disease, hospitalisations and mortality in older adults in England. medRxiv, 2021. DOI: 10.1101/2021.03.01.21252652.

64. Brandolin, B.A., et al., The inconspicuous nature of COVID-19 and its impact to dentistry. *Semin Orthod*, 2020. 26(4): 176–182. DOI: 10.1053/j.sodo.2020.11.002.
65. Banakar, M., et al., COVID-19 transmission risk and protective protocols in dentistry: a systematic review. *BMC Oral Health*, 2020. 20:275. DOI: 10.1186/s12903-020-01270-9.
66. Bizzoca, M.E., et al., Covid-19 Pandemic: What Changes for Dentists and Oral Medicine Experts? A Narrative Review and Novel Approaches to Infection Containment. *Int J Environ Res Public Health*, 2020. 17(11): 3793. DOI: 10.3390/ijerph17113793.
67. Kathree, B.A., et al., COVID-19 and its impact in the dental setting: A scoping review. *PLoS One*, 2020. 15(12): e0244352. DOI: 10.1371/journal.pone.0244352.
68. Coulthard, P., et al., The COVID-19 pandemic and dentistry: the clinical, legal and economic consequences - part 2: consequences of withholding dental care. *Br Dent J*, 2020. 229(12): 801–805. DOI: 10.1038/s41415-020-2406-9.
69. Wijesooriya, N.R., et al., COVID-19 and telehealth, education, and research adaptations. *Paediatr Respir Rev*, 2020. 35:38–42. DOI: 10.1016/j.prrv.2020.06.009.
70. Crawford, E. And Taylor, N. The effective use of an e-dentistry service during the COVID-19 crisis. *J Orthod*. 2020. 47(4):330-337. DOI: 10.1177/1465312520949557.
71. Ghai, S. Teledentistry during COVID-19 pandemic. *Diabetes Metab Syndr*, 2020. 14(5):933-935. DOI: 10.1016/j.dsx.2020.06.029.
72. Telles-Araujo, G.T., et al., Teledentistry support in COVID-19 oral care. *Clinics (Sao Paulo)*, 2020. 75:e2030. DOI: 10.6061/clinics/2020/e2030.
73. Tapia, R.O.C., et al., Oral mucosal lesions in patients with SARS-CoV-2 infection. Report of four cases. Are they a true sign of COVID-19 disease? *Special Care in Dentistry*, 2020. DOI: 10.1111/scd.12520.
74. Petruzzi, M. And Benediis, M., WhatsApp: a telemedicine platform for facilitating remote oral medicine consultation and improving clinical examinations. *Oral Surgery Oral Medicina Oral Pathology Oral Radiology*, 2016. 121(3):248-254. DOI: <https://doi.org/10.1016/j.oooo.2015.11.005>.
75. Saccomanno, S., et al., Perspectives of tele-orthodontics in the COVID-19 emergency and as a future tool in daily practice. *Eur J Paediatr Dent*, 2020. 21(2):157-162. DOI: 10.23804/ejpd.2020.21.02.12.
- 76 - Farronato, M., et al., A Call for Action to Safely Deliver Oral Health Care during and Post COVID-19 Pandemic. *Int J Environ Res Public Health*, 2020. 17(18):E6704. DOI: 10.3390/ijerph17186704

77. Centers for Disease Control and Prevention, Guidelines for Preventing the Transmission of Mycobacterium tuberculosis in Health-Care Settings, 2005. <https://www.cdc.gov/mmwr/preview/mmwrhtml/rr5417a1.htm>. (accedido a 19 de abril de 2020).
78. ACSS, Especificações Técnicas para Instalações de AVAC – ET 06/2008. [http://www.acss.min-saude.pt/wp-content/uploads/2016/09/Especificacoes\\_Tecnicas\\_06\\_2008.pdf](http://www.acss.min-saude.pt/wp-content/uploads/2016/09/Especificacoes_Tecnicas_06_2008.pdf). (accedido a 19 de abril de 2020).
79. Centers for Disease Control and Prevention, Guidelines for Environmental Infection Control in Health-Care Facilities (2003). Appendix B. Air. <https://www.cdc.gov/infectioncontrol/guidelines/environmental/appendix/air.html#tableb1>. (accedido a 19 de abril de 2020).
80. New York State Department of Health. New York State Department of Health. 2017. <https://www.health.ny.gov/facilities/cons/docs/3-14-0.pdf> (accedidos a 19 de abril de 2020).
81. Veterans Affairs. Dental Service design guide. 2014. <https://www.cfm.va.gov/til/dGuide/dgDental.pdf>. (accedido a 19 de abril de 2020).
82. Harrel, S.K. and Molinari J., Aerosols and splatter in dentistry: a brief review of the literature and infection control implications. *J Am Dent Assoc*, 2004. 135(4): p. 429-37.
83. Helmis, C.G., *et al.*, Indoor air quality in a dentistry clinic. *Sci Total Environ*, 2007. 377(2-3): p. 349-65.
84. Li, Y., *et al.*, Role of ventilation in airborne transmission of infectious agents in the built environment - a multidisciplinary systematic review. *Indoor Air*, 2007. 17(1): p. 2-18.
85. Raghunath, N., *et al.*, Aerosols in Dental Practice- A Neglected Infectious Vector. *Microbiology Research Journal International*, 2016. 14.
86. Qian, H. and Zheng, X., Ventilation control for airborne transmission of human exhaled bio-aerosols in buildings. *J Thorac Dis*, 2018. 10(Suppl 19): p. S2295-S2304.
87. Silva, M.G., Cientista da UC analisa os modos de transmissão da COVID-19 à luz dos conceitos de Qualidade do Ar Interior. <http://noticias.uc.pt/universo-uc/cientista-da-uc-analisa-os-modos-de-transmissao-da-covid-19-a-luz-dos-conceitos-de-qualidade-do-ar-interior/>. (accedido a 18 de abril de 2020). 2020.
88. OSHA, Guidance on Preparing Workplaces for COVID-19. 2020: p. 18-20.
89. World Health Organization, Infection prevention and control during health care when novel coronavirus (nCoV) infection is suspected Interim guidance. ISBN 978-92-4-000091-9. 2020.
90. Bioseguridad para la atención odontológica: COVID 19. Protocolo para gestión y cumplimiento de normas de higiene y bioseguridad. Helga Lang. <https://>

www.colegiohigienistascv.es/wp-content/uploads/2020/04/PROTOCOLO-DE-BIOSEGURIDAD-pdf.pdf.pdf (acedido a 15 de Abril de 2020).

91. Zhang, W. and Jiang, X., Measures and suggestions for the prevention and control of the novel coronavirus in dental institutions. *Frontiers of Oral and Maxillofacial Medicine*, 2020. 2.

92. Huh, S., How to train health personnel to protect themselves from SARS-CoV-2 (novel coronavirus) infection when caring for a patient or suspected case. *Journal of Educational Evaluation for Health Professions*, 2020. 17.

93. Van Doremalen N, *et al.*, Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *Engl J Med.*, 2020. Mar.

94. Orientação 022/2020: COVID-19:Procedimentos em Clínicas, Consultórios ou Serviços de Saúde Oral dos Cuidados de Saúde Primários, Sector Social e Privado. Data: 01/05/2020. Atualizada a 20/07/2020. (<https://www.dgs.pt/directrizes-da-dgs/orientacoes-e-circulares-informativas/orientacao-n-0222020-de-01052020-pdf.aspx>).

95. Lurie, N. and Carr, B.G., The Role of Telehealth in the Medical Response to Disasters. *JAMA Intern Med*, 2018. 178(6): p. 745-746. DOI: 10.1001/jamainternmed.2018.1314.

96. Reeves, J.J., *et al.*, Rapid Response to COVID-19: Health Informatics Support for Outbreak Management in an Academic Health System. *Journal of the American Medical Informatics Association*, 2020. 27(6):853-859. DOI: 10.1093/jamia/ocaa037.

97. Lv, N., *et al.*, Management of oral medicine emergencies during COVID-19: A study to develop practice guidelines. *J Dent Sci*, 2020. 16(1):493-500. Online ahead of print. DOI: 10.1016/j.jds.2020.07.016.

98. León, S. and Giacaman, R.A., COVID-19 and Inequities in Oral Health Care for Older People: An Opportunity for Emerging Paradigms. *JDR Clin Trans Res*, 2020. 5(4):290-292. DOI: 10.1177/2380084420934742.

99. Norma 007/2020: Prevenção e Controlo de Infecção por SARS-CoV-2 (COVID-19): Equipamentos de Proteção Individual (EPI). Data: 29/03/2020. (<https://www.dgs.pt/directrizes-da-dgs/normas-e-circulares-normativas/norma-n-0072020-de-29032020-pdf.aspx>).

100. Smales, F.C. and Samaranyake, L.P., Maintaining dental education and specialist dental care during an outbreak of a new coronavirus infection. Part 2: Control of the disease, then elimination. *Br Dent J*, 2003. 195(12):679-81. DOI: 10.1038/sj.bdj.4810819.

101. Li, R.W., *et al.*, Severe acute respiratory syndrome (SARS) and the GDP. Part II: implications for GDPs. *Br Dent J*, 2004. 197(3):130-4. DOI: 10.1038/sj.bdj.4811522.

102. Samaranayake, L.P. and Peiris, M., Severe acute respiratory syndrome and dentistry: a retrospective view. *J Am Dent Assoc*, 2004. 135(9):1292-302. DOI: 10.14219/jada.archive.2004.0405.
103. Huang, Z., et al., The Battle Against Coronavirus Disease 2019 (COVID-19): Emergency Management and Infection Control in a Radiology Department. *J Am Coll Radiol*, 2020. 17(6):710-716. DOI: 10.1016/j.jacr.2020.03.011.
104. Lai, T.H.T., et al., Stepping up infection control measures in ophthalmology during the novel coronavirus outbreak: an experience from Hong Kong. *Graefes Arch Clin Exp Ophthalmol*, 2020. 258(5):1049-1055. DOI: 10.1007/s00417-020-04641-8.
105. Maragakis, L.L., Coronavirus Symptoms: Frequently Asked Questions. <https://www.hopkinsmedicine.org/health/conditions-and-diseases/coronavirus/coronavirus-symptoms-frequently-asked%20questions>. (acedido a 17 de abril de 2020). Johns Hopkins Medicine, 2020.
106. Repici, A., et al., Coronavirus (COVID-19) outbreak: what the department of endoscopy should know. *Gastrointest Endosc*, 2020. 92(1):192-197. DOI: 10.1016/j.gie.2020.03.019.
107. Zhang, Y., et al., Hospital response to the COVID-19 outbreak: The experience in Shanghai, China. *J Adv Nurs*, 2020. 76(7):1483-1485. DOI: 10.1111/jan.14364.
108. Norma 020/2020: COVID-19: Definição de Caso de COVID-19. Data: 09/11/2020. ([https://covid19.min-saude.pt/wp-content/uploads/2020/11/Norma\\_020\\_2020.pdf](https://covid19.min-saude.pt/wp-content/uploads/2020/11/Norma_020_2020.pdf)).
109. Norma 004/2020: COVID-19: Abordagem do Doente com Suspeita ou Confirmação de COVID-19. Data: 23/03/2020. Atualizada a 14/10/2020. ([https://covid19.min-saude.pt/wp-content/uploads/2020/12/Norma-004\\_2020.pdf](https://covid19.min-saude.pt/wp-content/uploads/2020/12/Norma-004_2020.pdf)).
110. COVID-19: Fase de Mitigação. Procedimentos para Estruturas Residenciais para Idosos (ERPI), Unidades de Cuidados Continuados Integrados (UCCI) da Rede Nacional de Cuidados Continuados (RNCCI) e outras respostas dedicadas a pessoas idosas; instituições de acolhimento de crianças e jovens em risco. Orientação 009/2020 de 11/03/2020. Atualizada a 07/04/2020.
111. ADA, Interim Guidance for Minimizing Risk of COVID-19 Transmission. ADA.org/InterimGuidance. [https://www.ada.org/~media/CPS/Files/COVID/ADA\\_COVID\\_Int\\_Guidance\\_Treat\\_Pts.pdf](https://www.ada.org/~media/CPS/Files/COVID/ADA_COVID_Int_Guidance_Treat_Pts.pdf) (acedido a 15 de Abril de 2020).
112. European Centre for Disease Prevention and Control, Novel coronavirus (SARS-CoV-2) Discharge criteria for confirmed COVID-19 cases – When is it safe to discharge COVID-19 cases from the hospital or end home isolation? <https://www.ecdc.europa.eu/>

sites/default/files/documents/COVID-19-Discharge-criteria.pdf (acedido a 15 de Abril de 2020).

113. Orientação 010/2020: Infeção por SARS-CoV-2 (COVID-19) - Distanciamento Social e Isolamento. Data: 16/03/2020. (<https://www.dgs.pt/directrizes-da-dgs/orientacoes-e-circulares-informativas/orientacao-n-0102020-de-16032020-pdf.aspx>)

114. OMD, Devo atender um doente infetado com COVID-19, ou que representa um forte risco de contágio? <https://www.omb.pt/covid-19/perguntas/7iietr/>. (acedido a 26 de abril de 2020).

115. OMD, COVID-19: cuidados preventivos e consultas de medicina dentária. <https://www.omb.pt/2020/03/cuidados-preventivos-consultas/>. (acedido a 26 de abril de 2020).

116. Infeção por SARS-CoV-2 (COVID-19). Medidas de prevenção da transmissão em estabelecimentos de atendimento ao público. Orientação nº 011/2020 de 17/03/2020.

117. Boyce, J.M., *et al.*, Guideline for Hand Hygiene in Health-Care Settings. Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. Society for Healthcare Epidemiology of America/Association for Professionals in Infection Control/Infectious Diseases Society of America. MMWR Recomm Rep, 2002. 51(RR-16): p. 1-45, quiz CE1-4.

118. Guangming Ye, H.L., *et al.*, Environmental contamination of the SARS-CoV-2 in healthcare premises: An urgent call for protection for healthcare workers. medRxiv, 2020.

119. Centers for Disease Control and Prevention, Interim U.S. Guidance for Risk Assessment and Public Health Management of Healthcare Personnel with Potential Exposure in a Healthcare Setting to Patients with Coronavirus Disease (COVID-19). 2020: p. 1-7.

120. Consejo General de Dentistas de España, Plan Estratégico de Acción para Periodo Posterior a la Crisis por El Covid 19. 2020 (Version 13 abril de 2020).

121. Centers for Disease Control and Prevention, Best Practices for Environmental Cleaning in Healthcare Facilities : in Resource-Limited Settings. 2019: p. 1-91.

122. Lotfinejad, N., *et al.*, Hand hygiene and the novel coronavirus pandemic: The role of healthcare workers. J Hosp Infect, 2020.

123. Allegranzi, B., *et al.*, Burden of endemic health-care-associated infection in developing countries: systematic review and meta-analysis. Lancet, 2011. 377(9761): p. 228-41.

124. Jefferson, T., *et al.*, Physical interventions to interrupt or reduce the spread of respiratory viruses. Cochrane Database Syst Rev, 2011(7): p. CD006207.

125. Basile, C.C., *et al.*, Recommendations for the prevention, mitigation and containment of the emerging SARS-CoV-2 (COVID-19) pandemic in haemodialysis centres. *Nephrology Dialysis Transplantation*, 2020. in press.
126. World Health Organization, Guidelines on Hand Hygiene in Health Care: First Global Patient Safety Challenge Clean Care is Safer Care. ISBN 978 9241597906. 2009.
127. Picheansathian, W., A systematic review on the effectiveness of alcohol-based solutions for hand hygiene. *Int J Nurs Pract*, 2004. 10(1): p. 3-9.
128. Centers for Disease Control and Prevention, Guidelines for Infection Control in Dental Health-Care Settings. *Morbidity and Mortality Weekly Report*, 2003. 52.
129. Sebastiani, F.R., *et al.*, Infection Control in the Dental Office. *Dent Clin North Am*, 2017. 61(2): p. 435-457.
130. Kampf, G. and Löffler, H., Dermatological aspects of a successful introduction and continuation of alcohol-based hand rubs for hygienic hand disinfection. *The Journal of Hospital Infection*, 2003. 55.
131. Infeção por SARS-CoV-2 (COVID-19) - Limpeza e desinfecção de superfícies em estabelecimentos de atendimento ao público ou similares. Orientação nº 014/2020 de 21/03/2020.
132. Centers for Disease Control and Prevention, Healthcare Infection control Practices Advisory Committee: Guidelines for Environmental Infection Control in Health-care Facilities. 2003: p. 1-235.
133. Kundrapu S, *et al.*, Daily Disinfection of High-Touch Surfaces in Isolation Rooms to Reduce Contamination of Healthcare Workers' Hands. *Infect Control Hosp Epidemiol.*, 2012. 33(10): p. 4.
134. European Centre for Disease Prevention and Control, Disinfection of environments in healthcare and non- healthcare settings potentially contaminated with SARS-CoV-2. Stockholm;, 2020.
135. Consejo Dentistas - Organización Colegial de Dentistas de España. Plan estratégico de acción para el periodo posterior a la crisis creada por el COVID-19. 2020.
136. Consejo Dentistas - Organización Colegial de Dentistas de España. Recomendaciones de buenas prácticas para la atención de urgencias. <https://consejodentistas.es/comunicacion/actualidad-consejo/notas-de-prensa-consejo/item/1739-recomendaciones-de-buenas-practicas-para-la-atencion-de-urgencias.html>. (accedido a 19 de abril de 2020).
137. Handbook of COVID-19 Prevention and Treatment. <https://www.alnap.org/help-library/handbook-of-covid-19-prevention-and-treatment>. (accedido a 16 de Abril de 2020).



138. Chughtai, A.A., *et al.*, Policies on the use of respiratory protection for hospital health workers to protect from coronavirus disease (COVID-19). *Int J Nurs Stud*, 2020. 105: p. 103567.
139. Giwa, A.L., *et al.*, Novel 2019 coronavirus SARS-CoV-2 (COVID-19): An updated overview for emergency clinicians. *Emerg Med Pract*, 2020. 22(5): p. 1-28.
140. Leung NHL, *et al.*, Respiratory virus shedding in exhaled breath and efficacy of face masks. *Nature Medicine*, 2020: p. 1-5.
141. Murthy, S., *et al.*, Care for Critically Ill Patients With COVID-19. *JAMA*, 2020.
142. Ng, K., *et al.*, COVID-19 and the Risk to Health Care Workers: A Case Report. *Ann Intern Med*, 2020.
143. Wang, Q. and Yu C., The role of masks and respirator protection against SARS-CoV-2. *Infect Control Hosp Epidemiol*, 2020: p. 1-2.
144. Chinese Center for Disease Control and Prevention, Norms for Prevention and Control Practice in Grass-roots Medical and Health Institutions. (disponível em formato PDF) 2020.
145. Centers for Disease Control and Prevention, Interim U.S. Guidance for Risk Assessment and Public Health Management of Healthcare Personnel with Potential Exposure in a Healthcare Setting to Patients with Coronavirus Disease 2019 (COVID-19). <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-risk-assessment-hcp.html#table1%0A%0ANormas%20preven%C3%A7%C3%A3o-CDC%20Chin%C3%AAspdf>. (acedido a 15 de Abril de 2020). 2020.
146. Rimmer, A., Covid-19: What's the current advice for UK doctors? *BMJ*, 2020. 368: p. m978.
147. Rombola, G., *et al.*, Practical indications for the prevention and management of SARS-CoV-2 in ambulatory dialysis patients: lessons from the first phase of the epidemics in Lombardy. *J Nephrol*, 2020. 33(2): p. 193-196.
148. Krithikadatta, J., *et al.*, Endodontic and Dental Practice during COVID-19 Pandemic: Position Statement from International Federation of Endodontic Associations (IFEA) & Indian Endodontic Society (IES). [http://www.aede.info/pdf/newsletters/IFEA\\_IES\\_Endodontic\\_and\\_Dental\\_Practice\\_during\\_COVID-19.pdf](http://www.aede.info/pdf/newsletters/IFEA_IES_Endodontic_and_Dental_Practice_during_COVID-19.pdf). (acedido a 15 de Abril de 2020).
149. Cowling, B.J. and Aiello, A., Public health measures to slow community spread of COVID-19. *J Infect Dis*, 2020.
150. Infecção por SARS-CoV-2 (COVID-19) – Recolha, Transporte e Tratamento dos Resíduos Hospitalares. Orientação nº 012/2020 de 19/03/2020.

151. World Health Organization, Guidance for health workers. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/health-workers>. (acedido a 16 de Abril de 2020).
152. COVID-19: infection prevention and control (IPC). <https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control>. (acedido a 20 de abril de 2020).
153. European Centre for Disease Prevention and Control, Infection prevention and control and preparedness for COVID-19 in healthcare settings - second update. <https://www.ecdc.europa.eu/en/publications-data/infection-prevention-and-control-and-preparedness-covid-19-healthcare-settings>. (acedidio a 20 de abril de 2020).
154. Association of Occupational Health Professionals in Healthcare, Infections by Bio-aerosols. <http://ldh.la.gov/assets/oph/Center-PHCH/Center-CH/infectious-epi/HAI/InfectionsbyAerosolsDropletsHandout.pdf>. (acedido a 20 de abril de 2020).
155. Verbeek, J.H., *et al.*, Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. *Cochrane Database Syst Rev*, 2019. 7: p. CD011621.
156. Long Y, *et al.*, Effectiveness of N95 respirators versus surgical masks against influenza: A systematic review and meta-analysis. *J Evid Based Med.*, 2020. 2(February): p. 1-9.
157. Centers for Disease Control and Prevention, Guidance for Providing Dental Care During COVID-19. 2020.
158. CAL/OSHA, The California Workplace Guide to Aerosol Transmissible Diseases. [https://www.dir.ca.gov/dosh/dosh\\_publications/ATD-Guide.pdf](https://www.dir.ca.gov/dosh/dosh_publications/ATD-Guide.pdf). (acedido a 20 de abril de 2020).
159. Pan, Y., *et al.*, Transmission routes of SARS-CoV-2 and protective measures in dental clinics during the COVID-19 pandemic. *Am J Dent*, 2020 Jun;33(3):129-134. PMID: 32470237.
160. European Centre for Disease Prevention and Control, Guidance for wearing and removing personal protective equipment in healthcare settings for the care of patients with suspected or confirmed COVID-19. <https://www.ecdc.europa.eu/en/publications-data/guidance-wearing-and-removing-personal-protective-equipment-healthcare-settings>. (acedido a 20 de abril de 2020).
161. European Centre for Disease Prevention and Control, Personal protective equipment (PPE) needs in healthcare settings for the care of patients with suspected or confirmed novel coronavirus (2019-nCoV). <https://www.ecdc.europa.eu/en/publications->

- data/personal-protective-equipment-ppe-needs-healthcare-settings-care-patients. (accedido a 20 de abril de 2020).
162. European Centre for Disease Prevention and Control, Safe use of personal protective equipment in the treatment of infectious diseases of high consequence. <https://www.ecdc.europa.eu/en/publications-data/safe-use-personal-protective-equipment-treatment-infectious-diseases-high>. (accedido a 20 de abril de 2020).
163. Tang, J.W., *et al.*, Factors involved in the aerosol transmission of infection and control of ventilation in healthcare premises. *J Hosp Infect*, 2006. 64(2): p. 100-14.
164. Asadi, S., *et al.*, Aerosol emission and superemission during human speech increase with voice loudness. *Sci Rep*, 2019. 9(1): p. 2348.
165. Sociedad Española de Patología Digestiva and Asociación Española de Gastroenterología, Recommendations by the SEPD and AEG, both in general and on the operation of gastrointestinal endoscopy and gastroenterology units, concerning the current SARS-CoV-2 pandemic (March, 18). *Rev Esp Enferm Dig*, 2020. 112(4): p. 319-322.
166. Chin, A., *et al.*, Stability of SARS-CoV-2 in different environmental conditions. *medRxiv*, 2020.
167. Moriarty L.F., *et al.* Public Health Responses to COVID-19 Outbreaks on Cruise Ships — Worldwide, February–March 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:347-352. DOI: [http://dx.doi.org/10.15585/mmwr.mm6912e3external icon](http://dx.doi.org/10.15585/mmwr.mm6912e3external%20icon).
168. Ong, S.W.X., *et al.*, Air, Surface Environmental, and Personal Protective Equipment Contamination by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) From a Symptomatic Patient. *JAMA*, 2020.
169. Jiang, Y., *et al.*, Clinical Data on Hospital Environmental Hygiene Monitoring and Medical Staff Protection during the Coronavirus Disease 2019 Outbreak. *medRxiv*, 2020.
170. Zayas, G., *et al.*, Cough aerosol in healthy participants: fundamental knowledge to optimize droplet-spread infectious respiratory disease management. *BMC Pulmonary Medicine*, 2012. 12.
171. Bourouiba, L., Turbulent Gas Clouds and Respiratory Pathogen Emissions: Potential Implications for Reducing Transmission of COVID-19. *JAMA*, 2020.
172. Bentley, C.D., N.W. Burkhart, and J.J. Crawford, Evaluating Spatter and Aerosol Contamination during Dental Procedures. *JADA*, 1994. 125.
173. Bennett, A.M., *et al.*, Microbial aerosols in general dental practice. *British Dental Journal*, 2000. 189.

174. Hokett, S.D., *et al.*, Assessing the effectiveness of direct digital radiography barrier sheaths and finger cots. *J Am Dent Assoc*, 2000. 131(4): p. 463-7.
175. Swaminathan, Y. and Thomas, J.T., "Aerosol"-A Prospective Contaminant of Dental Environment! *IOSR Journal of Dental and Medical Sciences*, 2013. 11.
176. Otter, J.A., *et al.*, Transmission of SARS and MERS coronaviruses and influenza virus in healthcare settings: the possible role of dry surface contamination. *J Hosp Infect*, 2016. 92(3): p. 235-50.
177. Rutala, W.A. and Weber, D.J., *Guideline for Disinfection and Sterilization in Healthcare Facilities*, 2008. <https://www.cdc.gov/infectioncontrol/pdf/guidelines/disinfection-guidelines-H.pdf>. (acedido a 19 de abril de 2020). 2019.
178. Kampf, G., *et al.*, Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *J Hospital Infect*, 2020. 104(3): p. 246-251.
179. Diário da República Eletrónico, Portaria n.º 353-A/2013.
180. ISO 7730:2005, Ergonomics of the thermal environment — Analytical determination and interpretation of thermal comfort using calculation of the PMV and PPD indices and local thermal comfort criteria. <https://www.iso.org/standard/39155.html>. (acedido a 19 de abril de 2020).
181. Ijaz, M.K., *et al.*, Survival characteristics of airborne human coronavirus 229E. *J Gen Virol*, 1985. 66 ( Pt 12): p. 2743-8.
182. Offner, D., *et al.*, Contamination of dental unit waterlines: assessment of three continuous water disinfection systems. *BDJ Open*, 2016. 2: p. 16007.
183. Ji, X.Y., *et al.*, Three key factors influencing the bacterial contamination of dental unit waterlines: a 6-year survey from 2012 to 2017. *International Dental Journal*, 2018. 69.
184. Lizzadro, J., *et al.*, Comparison between Two Types of Dental Unit Waterlines: How Evaluation of Microbiological Contamination Can Support Risk Containment. *Int J Environ Res Public Health*, 2019. 16(3).
185. Desarda, H., *et al.*, Efficacy of High-volume Evacuator in Aerosol Reduction: Truth or Myth? A Clinical and Microbiological Study. *J Dent Res Dent Clin Dent Prospects*, 2014. 8(3): p. 176-9.
186. Avasthi, A., High Volume Evacuator (HVE) in reducing aerosol- an exploration worth by clinicians. *Journal of Dental Health Oral Disorders & Therapy*, 2018. 9.
187. Van der Sluijs, M., *et al.*, The effect on clinical parameters of periodontal inflammation following non-surgical periodontal therapy with ultrasonics and chemotherapeutic cooling solutions: a systematic review. *J Clin Periodontol*, 2016. 43(12): p. 1074-1085.

188. Jawade, R., *et al.*, Comparative Evaluation of Two Different Ultrasonic Liquid Coolants on Dental Aerosols. *J Clin Diagn Res*, 2016. 10(7): p. ZC53-7.
189. Sethi, K.S., *et al.*, Comparative evaluation of the chlorhexidine and cinnamon extract as ultrasonic coolant for reduction of bacterial load in dental aerosols. *J Indian Soc Periodontol*, 2019. 23(3): p. 226-233.
190. Howard, BE. High-Risk Aerosol-Generating Procedures in COVID-19: Respiratory Protective Equipment Considerations. *Otolaryngol Head Neck Surg*, 2020. 163(1):98-103. DOI: 10.1177/0194599820927335.
191. Zimmermann, M. And Nkenke, E., Approaches to the management of patients in oral and maxillofacial surgery during COVID-19 pandemic. *J Craniomaxillofac Surg*, 2020. 48(5):521–526. DOI: 10.1016/j.jcms.2020.03.011.
192. Alharbi, A., *et al.*, Guidelines for dental care provision during the COVID-19 pandemic. *Saudi Dent J*, 2020. 32(4):181-186. DOI: 10.1016/j.sdentj.2020.04.001.
193. Ayyed AB. Dental Practice Infection Control Measurements: Coronavirus Disease (COVID-19) Outbreaks. *Int J Clin Pediatr Dent*, 2020. 13(3):279-283. DOI 10.5005/jp-journals-10005-1770.
194. Falahchai, M., *et al.*, Dental care management during the COVID-19 outbreak. *Spec Care Dentist*, 2020. DOI: 10.1111/scd.12523.
195. Park, SH. Personal Protective Equipment for Healthcare Workers during the COVID-19. *Infect Chemother*, 2020. 52(2):165-182. DOI:10.3947/ic.2020.52.2.165.
196. Samaranayake LP, *et al.*, The effectiveness and efficacy of respiratory protective equipment (RPE) in dentistry and other health care settings: a systematic review. *Acta Odontol Scand*. 2020. 3:1-14. DOI: 10.1080/00016357.2020.1810769.
197. Jamal, M, *et al.*, Overview of transnational recommendations for COVID-19 transmission control in dental care settings. *Oral Dis*, 2020. DOI: 10.1111/odi.13431.
- 198 Agência Nacional de Vigilância Sanitária, Cartilha de Proteção Respiratória contra Agentes Biológicos para Trabalhadores de Saúde.  
<http://www2.ebserh.gov.br/documents/214604/816023/Cartilha+de+Prote%C3%A7%C3%A3o+Respirat%C3%B3ria+contra+Agentes+Biol%C3%B3gicos+para+Trabalhadores+de+Sa%C3%BAde.pdf/58075f57-e0e2-4ec5-aa96-743d142642f1>. (acedido a 26 de abril de 2020).
199. Gallagher JE, *et al.*, Relevance and paucity of evidence: a dental perspective on personal protective equipment during the COVID-19 pandemic. *Br Dent J*, 2020. 229(2):121-124. DOI: 10.1038/s41415-020-1843-9.

200. Kantor, J., Behavioral considerations and impact on personal protective equipment (PPE) use: Early lessons from the coronavirus (COVID-19) outbreak. *J Am Acad Dermatol*, 2020. 82: p. 1087-1088.
201. Reis, I.N.R., et al., Can preprocedural mouthrinses reduce SARS-CoV-2 load in dental aerosols? *Med Hypotheses*, 2021. 146:110436. DOI: 10.1016/j.mehy.2020.110436.
202. Castro-Ruiz, C. and Vergara-Buenaventura, A., Povidone-Iodine Solution: A Potential Antiseptic to Minimize the Risk of COVID-19? A Narrative Review. *J Int Soc Prev Community Dent*, 2020. 10(6):681-685. DOI: 10.4103/jispcd.JISPCD\_304\_20.
203. Carrouel, F., et al., Antiviral Activity of Reagents in Mouth Rinses against SARS-CoV-2. *J Dent Res*, 2021. 100(2):124-132. DOI: 10.1177/0022034520967933.
204. Ather, A., et al., Efficacy of Mouth Rinses Against SARS-CoV-2: A Scoping Review. *Frontiers in Dental Medicine*, 2021. 2:648547. DOI: 10.3389/fdmed.2021.648547.
205. Yoon, J.G., et al., Clinical Significance of a High SARS-CoV-2 Viral Load in the Saliva. *J Korean Med Sci*, 2020. 35(20): e195. DOI: 10.3346/jkms.2020.35.e195.
205. Martínez Lamas, L., et al., Is povidone-iodine mouthwash effective against SARS-CoV-2? First in vivo tests. *Oral Dis*, 2020 [epub ahead of print 2 Jul 2020]. DOI: 10.1111/odi.13526.
206. Gottsauner, M.J., et al., A prospective clinical pilot study on the effects of a hydrogen peroxide mouthrinse on the intraoral viral load of SARS-CoV-2. *Clin Oral Investig*, 2020. 24(10):3707– 3713. DOI: 10.1007/s00784-020-03549-1.
207. Seneviratne, C.J., et al., Efficacy of commercial mouth-rinses on SARS-CoV-2 viral load in saliva: randomized control trial in Singapore. *Infection*, 2020. [online ahead of print]. DOI: 10.1007/s15010-020-01563-9.
208. Nejatidanesh, F., *et al.*, Risk of Contamination of Different Areas of Dentist's Face During Dental Practices. *Int J Prev Med*, 2013. 4(5): p. 611-5.
209. Caldeira, D., *et al.*, Skin antiseptics in venous puncture-site disinfection for prevention of blood culture contamination: systematic review with meta-analysis. *J Hosp Infect*, 2011. 77(3): p. 223-32.
210. Kamel, C., *et al.*, in *Preoperative Skin Antiseptic Preparations and Application Techniques for Preventing Surgical Site Infections: A Systematic Review of the Clinical Evidence and Guidelines*. 2011: Ottawa (ON).
211. Samaranayake, L.P., *et al.*, The efficacy of rubber dam isolation in reducing atmospheric bacterial contamination. *ASDC J Dent Child*, 1989. 56(6): p. 442-4.

212. El-Din, A.M.T. and Ghoname, N.A.E.-H., Efficacy of rubber dam isolation as an infection control procedure in paediatric dentistry. *Eastern Mediterranean Health Journal*, 1997. 3.
213. Al-Amad, S.H., *et al.*, The effect of rubber dam on atmospheric bacterial aerosols during restorative dentistry. *J Infect Public Health*, 2017. 10(2): p. 195-200.
214. Centers for Disease Control and Prevention, Infection Prevention & Control Guidelines & Recommendations. <https://www.cdc.gov/oralhealth/infectioncontrol/guidelines/index.htm> (accedido a 19 de abril de 2020).
215. Day, C.J., *et al.*, Aerosols and Splatter in Dentistry – A Neglected Menace? *Dental Update*, 2017. 33.
216. Hu, T., *et al.*, Risk of hepatitis B virus transmission via dental handpieces and evaluation of an anti-suction device for prevention of transmission. *Infect Control Hosp Epidemiol*, 2007. 28(1): p. 80-2.
217. Centers for Disease Control and Prevention, Guidelines for infection control in Dental Health Care settings. *J Am Dent Assoc*, 2003. S2(RR17): p. 1-68.
218. Hamedani, S. and Farshidfar, N. The practice of oral and maxillofacial radiology during COVID-19 outbreak. *Oral Radiol*, 2020. 36(4):400-403. DOI: 10.1007/s11282-020-00465-8.
219. Doriguêto, P.V.T., *et al.*, Challenges for the dental radiology clinic in times of the COVID-19 pandemic. *Oral Radiol*, 2020. 36(4):404-405. DOI: 10.1007/s11282-020-00456-9.
220. Ilhan, B., *et al.*, Dental radiographic procedures during COVID-19 outbreak and normalization period: recommendations on infection control. *Oral Radiol*, 2020. 36(4):395-399. DOI: 10.1007/s11282-020-00460-z.
221. Tuttnauer Team. How to better protect dental and medical clinics from airborne contamination? <https://tuttnauer.com/blog/how-to-protect-dental-and-medical-clinics-from-airborne-contamination>. (accedido a 19 de abril de 2020).
222. Centers for Disease Control and Prevention, Interim Infection Prevention and Control Recommendations for Patients with Suspected or Confirmed Coronavirus Disease 2019 (COVID-19) in Healthcare Settings [https://www.cdc.gov/coronavirus/2019-ncov/hcp/infection-control-recommendations.html?CDC\\_AA\\_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Finfection-control%2Fcontrol-recommendations.html](https://www.cdc.gov/coronavirus/2019-ncov/hcp/infection-control-recommendations.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Finfection-control%2Fcontrol-recommendations.html). (accedido a 19 de abril de 2020).

223. Hu, X., *et al.*, Heat inactivation of serum interferes with the immunoanalysis of antibodies to SARS-CoV-2. medRxiv, 2020.
224. Colaneri, M., *et al.*, Lack of SARS-CoV-2 RNA environmental contamination in a tertiary referral hospital for infectious diseases in Northern Italy. J Hosp Infect, 2020.
225. United States Environmental Protection Agency Washington, Prevention List N Products with Emerging Viral Pathogens AND Human Coronavirus claims for use against SARS-CoV-2. 2020.
226. Gurzawska-Comis, K., *et al.* Recommendations for Dental Care during COVID-19 Pandemic. J Clin Med, 2020. 9(6):1833. DOI: 10.3390/jcm9061833.
227. Poggio, C., *et al.*, Copper-Alloy Surfaces and Cleaning Regimens against the Spread of SARS-CoV-2 in Dentistry and Orthopedics. From Fomites to Anti-Infective Nanocoatings. Materials (Basel), 2020. 13(15):3244. DOI: 10.3390/ma13153244.
228. Bali, R.K. and Chaudhry, K., Maxillofacial surgery and COVID-19, The Pandemic !! J Maxillofac Oral Surg, 2020. 19(2):159-161. DOI: 10.1007/s12663-020-01361-8.
229. Dexter, F., *et al.*, Perioperative COVID-19 Defense: An Evidence-Based Approach for Optimization of Infection Control and Operating Room Management. Anesth Analg, 2020.
230. Haralur, S.B., *et al.*, Effect of alginate chemical disinfection on bacterial count over gypsum cast. J Adv Prosthodont, 2012. 4(2): p. 84-8.
231. Azevedo, M.J., *et al.*, A simple and effective method for addition silicone impression disinfection. J Adv Prosthodont, 2019. 11(3): p. 155-161.
232. Bass, R.A., *et al.*, The effect of a surface disinfectant on a dental cast. The Journal of Prosthetic Dentistry. 1992.
233. Soares, C.R. and Ueti, M., [Influence of different methods of chemical disinfection on the physical properties of type IV and V gypsum dies]. Pesqui Odontol Bras, 2001. 15(4): p. 334-40.
234. Pisulkar, S., *et al.*, Analysis of Bacterial Colonization due to Salivary Contamination on Gypsum Casts and Effect of Various Means of Disinfection of Dental Casts on Bacterial Colonization: An in vitro Study. International Journal of Oral Care and Research,, 2018. 6.
235. da Silva, F.C., *et al.*, Effectiveness of six different disinfectants on removing five microbial species and effects on the topographic characteristics of acrylic resin. J Prosthodont, 2008. 17(8): p. 627-33.



236. Centers for Disease Control and Prevention, Sequence for putting on personal protective equipment (PPE). <https://www.cdc.gov/hai/pdfs/ppe/ppe-sequence.pdf> (accedido a 19 de abril de 2020).
237. Centers for Disease Control and Prevention, Decontamination and Reuse of Filtering Facepiece Respirators. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/ppe-strategy/decontamination-reuse-respirators.html>. (accedido a 19 de abril de 2020).
238. COEC. [COVID-19] Recomendaciones e infografías para la atención de urgencias dentales durante la pandemia. <http://www.coec.cat/es/noticies/atencio-urg-dentals-covid19>. (accedido a 19 de abril de 2020).
239. Centers for Disease Control and Prevention, Strategies for Optimizing the Supply of Eye Protection. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/ppe-strategy/eye-protection.html>. (accedido a 19 de abril de 2020).
240. Centers for Disease Control and Prevention, Cleaning and Disinfection for Households. [www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/cleaning-disinfection.html](http://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/cleaning-disinfection.html). (accedido a 19 de abril de 2020).
241. Auer, R., *et al.*, Covid-19, cash, and the future of payments. BIS Bulletin. ISBN 92-9197-357-5. <https://www.bis.org/publ/bisbull03.pdf>. (accedido a 19 de abril de 2020). 2020.
242. Volgenant, C.M.C., *et al.*, Infection control in dental health care during and after the SARS-CoV-2 outbreak. *Oral Dis*, 2021. 27(3):674-683. DOI: 10.1111/odi.13408.
243. Centers for Disease Control and Prevention, Sterilization or disinfection of dental instruments. *JADA*, 2001. 132.
244. Schwartz, A., *et al.*, Decontamination and Reuse of N95 Respirators with Hydrogen Peroxide Vapor to Address Worldwide Personal Protective Equipment Shortages During the SARS-CoV-2 (COVID-19) Pandemic. *Journal of ABSA International*, 2020.
245. Hoffmann, M., *et al.*, SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*, 2020. 181(2): p. 271-280 e8.
246. Wolfel, R., *et al.*, Virological assessment of hospitalized patients with COVID-2019. *Nature*, 2020.
247. To, K.K.-W., *et al.*, Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *The Lancet*, 2020.
248. Liu, L., *et al.*, Epithelial cells lining salivary gland ducts are early target cells of severe acute respiratory syndrome coronavirus infection in the upper respiratory tracts of rhesus macaques. *J Virol*, 2011. 85(8): p. 4025-30.

249. Xu, H., *et al.*, High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci*, 2020. 12(1): p. 8.
250. Yan Y, *et al.*, Consensus of Chinese experts on protection of skin and mucous membrane barrier for health-care workers fighting against coronavirus disease 2019. *Dermatol Ther.*, 2020. 13(e13310).
251. Neher, R.A., *et al.*, Potential impact of seasonal forcing on a SARS-CoV-2 pandemic. *Swiss Med Wkly*, 2020. 150: p. w20224.
252. Dion, M., *et al.*, Big Data and the Global Public Health Intelligence Network (GPHIN). *Can Commun Dis Rep*, 2015. 41(9): p. 209-214.
253. Hollander, J.E. and Carr, B.G., Virtually Perfect? Telemedicine for Covid-19. *N Engl J Med*, 2020.
254. Perry, D.J., *et al.*, Guidelines for the management of patients on oral anticoagulants requiring dental surgery. *Br Dent J*, 2007. 203(7): p. 389-93.
255. Kirchhof, P., *et al.*, 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *European Journal of Cardio-Thoracic Surgery*, 2016. 50(5).



## Additional Bibliography

1. Yonenaga, K., et al., Summary of implications for clinical practices in dentistry under coronavirus disease 19 (COVID-19): a systematic review. Registo PROSPERO.
2. Kumar, G., et al., Guidelines and standard operating protocol for pediatric dental practice during COVID-19: a systematic review. Registo PROSPERO.
3. Mahdi, S.S., et al., Pivoting Dental Practice Management during the COVID-19 Pandemic - A Systematic Review. *Medicina (Kaunas)*, 2020. 56(12):644. DOI: 10.3390/medicina56120644.
4. Melo Neto C.L.M., et al., SARS-CoV-2 and Dentistry-Review. *Eur J Dent*, 2020. 14(S 01):S130-S139. DOI: 10.1055/s-0040-1716438.
5. Tysiąg-Miśta, M., et al., Air disinfection procedures in the dental office during the COVID-19 pandemic. *Med Pr*, 2021. 72(1):39-48. DOI: 10.13075/mp.5893.01005.
6. Soares, R., et al., Biosafety Guidelines in Dentistry for COVID-19 control/prevention: systematic review. Registo PROSPERO.
7. Mouth Rinses for Inactivation of COVID-19. NCT04584684. <https://clinicaltrials.gov/show/NCT04584684>, 2020.
8. Effect of oral health care and mouth rinse in treatment of Covid-19. IRCT20200527047581N1. <http://www.who.int/trialsearch/Trial2.aspx?TrialID=IRCT20200527047581N1>, 2020.